

A mis padres.

Gracias por vuestro amor, dedicación y apoyo incondicional.

Dr. José María Iriondo Alegría, Catedrático del Departamento de Biología y Geología, Física y Química Inorgánica y Analítica de la Universidad Rey Juan Carlos,

CERTIFICA:

Que los trabajos de investigación desarrollados en la memoria de tesis doctoral: "Evaluación de los factores que determinan la respuesta de las plantas de alta montaña mediterránea ante el cambio global. Una perspectiva integradora", han sido realizados bajo su supervisión y son aptos para ser presentados por el licenciado Carlos Lara Romero ante el tribunal que en su día se consigne, para aspirar al Grado de Doctor en el Programa de Doctorado de Conservación de Recursos Naturales por la Universidad Rey Juan Carlos.

VºBº Director de Tesis

Dr. José María Iriondo Alegría



Universidad
Rey Juan Carlos

Departamento de Biología y
Geología, Física y Química
Inorgánica y Analítica

**Evaluación de los factores que determinan la respuesta de las
plantas de alta montaña mediterránea ante el cambio global.
Una perspectiva integradora**

Determinants of Mediterranean high-mountain plant responses to
global change. An integrative approach

Tesis Doctoral
Carlos Lara Romero
Septiembre 2014

Director: José María Iriondo Alegría

“Many of my colleagues in other fields are surprised to learn that the study of biodiversity is still largely in a Linnaean phase of discovering and naming new species. Although our tools are more advanced, in many ways the science of biodiversity is not much farther along than medicine was in the Middle Ages. We are still at the state, as it were, of cutting open bodies to find out what organs are inside. The low investment in and slow pace of biodiversity research might be tolerable were it not for the overwhelmingly rapid destruction of the natural world. Without hyperbole we can truthfully say that we are almost out of time to save much of the diversity of life on Earth”.

Stephen P. Hubbell, 2001

The Unified Neutral Theory of Biodiversity and Biogeography

ÍNDICE

RESUMEN

Síntesis	15
Summary (English versión)	19
Antecedentes	23
Objetivos	32
Metodología general	34
Estructura general de la Tesis	45
Conclusiones	51
Bibliografía	53

CAPÍTULOS/CHAPTERS

Capítulo 1/ Chapter 1	61
Assessing intraspecific variation in effective dispersal along an altitudinal gradient: a test in two Mediterranean high-mountain plants	
Capítulo 2/ Chapter 2	107
Variation in fine-scale genetic structure of <i>Silene ciliata</i> (Caryophyllaceae) across populations with different spatial aggregation patterns	
Capítulo 3/ Chapter 3	141
Population spatial patterns conditioned by dispersal and heterogeneity reveal negative density dependence on reproductive output in spite of positive spatial associations	
Capítulo 4/ Chapter 4	173
Genetic variation in flowering phenology and reproductive performance in a Mediterranean high-mountain specialist	
Capítulo 5/ Chapter 5	203
Direct and indirect effects of shrub encroachment on alpine grasslands by altering the assemblage of pollinators	
CONCLUSIONS	247
COAUTORES/COAUTHORS	251
AGRADECIMIENTOS/ACKNOWLEDGMENTS	255

RESUMEN

SUMMARY

SÍNTESIS

Dada la elevada vulnerabilidad de los ecosistemas de montaña mediterráneos al cambio climático y otros motores de cambio global, resulta de primordial interés evaluar su capacidad de respuesta en el contexto actual y futuro de cambio ambiental. Existen, sin embargo, lagunas significativas en el conocimiento disponible sobre algunos de los mecanismos que están gobernando las respuestas que permiten su supervivencia en el actual escenario de cambio. Teniendo en cuenta estas consideraciones el objetivo general de la tesis doctoral es completar el conocimiento sobre los principales factores que determinan la respuesta de las plantas de alta montaña mediterránea al cambio global. Para ello, se ha seleccionado la comunidad de pastos psicroxerófilos de la Sierra de Guadarrama, de la cual se conocen numerosos aspectos de la biología, estado demográfico y éxito reproductivo de dos de sus especies más representativas: *Silene ciliata* y *Armeria caespitosa*. La tesis ha seguido un enfoque integral que contemple conjuntamente el patrón espacial del hábitat junto a la capacidad de dispersión de las especies, las interacciones bióticas, la estructura genética espacial y la base genética de la variación fenotípica. Todos estos factores se evaluaron a lo largo de gradientes ambientales para recoger la elevada variabilidad ambiental a escala local que caracteriza a los ambientes de alta montaña y que puede afectar de manera significativa a los factores estudiados en la tesis.

En el **capítulo 1** hemos utilizado una aproximación de modelización inversa para evaluar los patrones de reclutamiento en cinco poblaciones de *Silene ciliata* y *Armeria caespitosa* a lo largo de un gradiente altitudinal. La distancia media de dispersión efectiva fue corta en ambas especies (rango: 0,23 – 0,71m), probablemente debido a que procesos post-dispersivos relacionados con la sequía estival reducen su distancia de dispersión primaria. Ambas especies mostraron variación interpoblacional en el tipo de kernel que proporciona mejores ajustes y en el parámetro de fecundidad. Sin embargo, solo *A. caespitosa* exhibió diferencias significativas entre poblaciones en la distancia media de dispersión. La variación

ambiental altitudinal condicionó la fecundidad de los individuos en ambas especies pero no afectó a la escala espacial de la dispersión. Estos resultados resaltan la complejidad de la variación intraespecífica en los procesos que controlan el reclutamiento y sugieren la necesidad de análisis específicos en un amplio espectro de especies y ambientes para evaluar la prevalencia y magnitud de la variación intraespecífica de la dispersión. Otra conclusión significativa es que la baja distancia de dispersión de estas especies podría restringir su capacidad migratoria frente al cambio climático y limitar el flujo genético entre poblaciones.

En el **capítulo 2** aplicamos herramientas de análisis de autocorrelación espacial para cuantificar la estructura espacial demográfica y genética en cinco poblaciones de *Silene ciliata* estables demográficamente pero espacialmente heterogéneas. Encontramos estructura espacial genética a escala local (FSGS por sus siglas en inglés) significativa en tres de las cinco poblaciones estudiadas, confirmando la existencia de variación intra-poblacional. Se detectó una asociación positiva entre la intensidad de la FSGS y el nivel de agregación espacial de las plantas. Este resultado avala a los estudios teóricos que muestran que los patrones espaciales agregados tienden a aumentar los niveles de FSGS en comparación con distribuciones aleatorias. Los índices de diversidad genética y endogamia observados en las poblaciones estudiadas fueron homogéneos. Consecuentemente, no se encontró relación entre la intensidad de la FSGS y dichos parámetros. Esto sugiere que la variabilidad ambiental a escala local no está afectando a la variabilidad genética y endogamia de las poblaciones estudiadas.

En el **capítulo 3** aplicamos técnicas de análisis de patrón de puntos para describir la estructura espacial de *Silene ciliata* a lo largo de un gradiente altitudinal y evaluar la contribución que la dispersión, la heterogeneidad de hábitat y las interacciones intraespecífica tienen en la conformación del patrón observado. Las plántulas mostraron una estructura espacial agregada que parece estar controlada por la disponibilidad heterogénea de micrositios adecuados para la germinación y el establecimiento. La agregación inducida por la dispersión y el reclutamiento de

plántulas persistió en la clase adulta. Pese a la tendencia de la especie a distribuirse de forma agregada detectamos evidencias de densodependencia negativa en su éxito reproductivo. Los resultados en su conjunto demuestran que los patrones de co-ocurrencia no tienen por qué estar mediados por interacciones de facilitación, por lo que antes de establecer relaciones causales entre éstas y la distribución espacial de las plantas hay que evaluar otros factores como la dispersión o la heterogeneidad ambiental.

El objetivo del **capítulo 4** fue evaluar si la variación fenotípica observada en un conjunto de rasgos reproductivos y vegetativos en las poblaciones naturales de la especie tiene una base genética. Con este propósito utilizamos una aproximación de jardín común (*Common garden*) para cuantificar la variación en este conjunto de rasgos en cuatro poblaciones situadas en los límites elevacionales superior e inferior del rango de distribución de la especie. Las plantas procedentes del límite de elevación superior florecieron antes y produjeron más frutos viables, pero crecieron menos en comparación con las plantas procedentes del límite de elevación inferior. Estos resultados confirman que la variación inter-poblacional observada en algunos de los rasgos ligados a la fenología y el éxito reproductivo tiene base genética. El estudio también mostró que la respuesta selectiva en favor de una floración temprana detectada en el límite de elevación inferior, no ha generado una mayor proporción de fenotipos de floración temprana. Sin embargo, sugerimos que la variabilidad genética asociada al inicio de la floración, podría ser relevante para hacer frente al impacto ocasionado por el cambio climático en curso.

En el **capítulo 5** hemos utilizado técnicas de análisis redes complejas para cuantificar el efecto de la matorralización de los pastos de alta montaña en la red de interacciones planta-polinizador. Construimos matrices de interacción cuantitativas para comparar parcelas matorralizadas frente a parcelas en las que el matorral no estaba presente. La matorralización de los pastos estudiados tuvo un efecto positivo sobre la riqueza y la frecuencia de visitas de los polinizadores a las plantas, pero incrementó la competencia entre plantas por los servicios de polinización. Esta

última situación podría reducir el éxito reproductivo en alguna de las especies de plantas estudiadas. Un amplio número de grupo funcionales de polinizadores visitaron a las especies de matorral en mayor proporción que a la mayoría de las especies de plantas de la comunidad, lo que otorgó a los matorrales un rol de súper-generalista. Pese a ello, la estructura de la red de interacciones planta-polinizador apenas mostró cambios significativos en sus propiedades estructurales. El efecto diferencial del matorral en las plantas y los polinizadores resalta la necesidad de utilizar un enfoque integrador para evaluar el efecto de los motores de cambio global sobre las interacciones interespecíficas y la estructura y el funcionamiento de las comunidades ecológicas.

SUMMARY

Global change drivers are causing relevant alterations in Mediterranean high-mountain ecosystems. Although some first attempts to evaluate the effects of these modifications have been made, we still far from understand in detail the factors and processes that determine how Mediterranean high-mountain plants respond to accelerated climate change and other important drivers of global change. Thus, in this doctoral dissertation we followed an integrative approach to study three large groups of factors which *a priori* are determinant: a) the spatial pattern of the habitat together with the dispersal capacity of each species, b) the biotic interactions and c) the spatial genetic structure and the genetic basis of the phenotypic variation. The study site chosen was high-mountain psicroxerophilous pastures of the Central System (Spain), as this ecosystem is one of the most vulnerable to climate change and other important global change drivers. Most studies were performed on five populations of each of two representative species of the community: *Silene ciliata* Poiret and *Armeria caespitosa* (Gómez Ortega) Boiss. in DC.

In **chapter 1** we applied an inverse modeling approach to measure the net reproductive rate and effective dispersal patterns in five populations of both *S. ciliata* and *A. caespitosa* along an altitudinal gradient. Results showed that no single kernel function provides the best fit across all populations, although estimated mean dispersal distances were short in all cases (range: 0.23 – 0.71 m). Post-dispersal factors related to summer drought appeared to restrict effective seed dispersal in these two species. The altitudinal gradient appears to influence effective seed dispersal patterns in *S. ciliata* and *A. caespitosa* through adult plant fecundity and seedling density, but not through the effective seed dispersal range. There were significant differences in the fecundity parameter among populations in both species and in the mean dispersal distance among populations of *A. caespitosa*, while differences in the mean dispersal distance among *S. ciliata* populations were only marginally significant. These results illustrate the complexity of intraspecific variation in the processes underlying recruitment and call for a case-by-case analysis

in a wider range of plant taxa and environments to assess the prevalence and magnitude of intraspecific dispersal variation. Additionally, we suggested that the limited dispersal range of the two studied species raises uncertainty on their ability to track their climatic niche through migration under ongoing rapid global warming.

In **chapter 2** we used spatial autocorrelation analysis to assess the spatial demographic and genetic structure of five demographically stable but spatially heterogeneous populations of *S. ciliata*. We found significant FSGS in three of the five populations studied, and therefore, revealed among-population variation in the intensity of FSGS. There was a significant positive association between the spatial demographic and genetic structure measured in each population. These findings support theoretical studies showing that stronger spatial aggregation of adults generally increases the magnitude of FSGS compared to random individual distributions. Finally, we did not detect any relationship between population genetic diversity and inbreeding with FSGS, suggesting that population genetic diversity and inbreeding are not related with fine-scale environmental variation.

In **chapter 3** we used techniques of spatial point pattern analysis to study the spatial pattern of *Silene ciliata* in the same five populations referred in chapters 1 and 2. We aimed to depict the spatial structure of the species and its relation to plant performance across different populations assessing the contribution of dispersal, environmental heterogeneity and conspecific plant interactions to the observed patterns. Recruits exhibited a pronounced small-scale aggregation and large-scale heterogeneity that were well characterized by a heterogeneous Poisson cluster process. The spatial pattern of recruits after seed germination and establishment was independent of the spatial pattern of adults. The aggregation induced by seed dispersal and subsequent seedling recruitment persisted in the adult class. Even though this aggregated structure might suggest the presence of facilitative interactions among adults, a negative density-dependent effect on reproductive output was found. We suggest that the aggregated pattern observed is the result of environmental heterogeneity and short distance dispersal of the species together with

the availability of safe-sites for establishment, with a reduced effect of plant-plant interactions. Thus, our results provide empirical evidence for the lack of a simple and direct relationship between plant population structure and the sign of plant-plant interactions (*i.e.*, competition and facilitation) and outlines the importance of including all relevant ecological factors involved in determining plant spatial patterns, such as seed dispersal and environmental heterogeneity, if we want to fully understand the underlying mechanisms of plant spatial patterns.

In **chapter 4** we focused on *A. caespitosa* to assess whether the observed phenotypic variation in a suite of reproductive and vegetative traits in natural population has a genetic basis or is the result of plastic responses shaped by heterogeneous environmental conditions. For this purposes we used a common garden approach to study the variation of such traits in four *A. caespitosa* populations at the high and low edges of the species distribution range. Our results confirmed that among-population variation in flowering phenology and some relevant traits related to reproductive performance is genetically based. The later flowering onset found in the plants of the low-edge of the species distribution did not correspond with the stronger selection response in favor of early-flowering individuals previously reported in natural populations. Finally, we suggested that the inherited capacity of the species to shift its flowering phenology coupled with the observed genetic variation within populations indicates that there is adaptive potential to respond to ongoing global warming.

In **chapter 5** we used network analysis tools to investigate the effects of shrub encroachment on plant-pollinator interactions at the community level. We constructed quantitative visitation networks to compare replicated shrub encroached and un-encroached plots. Results showed distinctive effects of shrub encroachment on plant and pollinator species because it increased richness and visitation patterns of pollinators, but also increased pollinator-mediated competition among plant species, which can potentially result in reduced reproductive success. In spite of the central role of shrub species the structure of the networks remained relatively

unchanged. This result is congruent with previous findings in the literature suggesting that plant-pollinator mutualism may be robust to disruptions.

ANTECEDENTES

Motores de cambio global y respuestas de las comunidades vegetales

El concepto de cambio global incluye todos aquellos impactos originados por actividades antrópicas que modifican la composición y funcionamiento de los ecosistemas (Vitousek, 1992). Como consecuencia de estas alteraciones, la biodiversidad mundial está cambiando a un ritmo sin precedentes (Sala *et al.*, 2000; Thuiller *et al.*, 2005; Parmesan *et al.*, 2006), lo que se ha convertido en una cuestión importante tanto para la investigación ecológica como para los programas gubernamentales sobre cambio global (Sala *et al.*, 2000; MEA, 2005; IPCC, 2007; Thuiller *et al.*, 2008; Thomas *et al.*, 2010; Pauli *et al.*, 2012). Los cambios en los usos del suelo y el cambio climático acelerado son probablemente los motores de cambio global que tienen mayor incidencia sobre los ecosistemas terrestres (Vitousek *et al.*, 1992; Sala *et al.*, 2000). De hecho, existen considerables evidencias sobre su efecto directo en alteraciones del rango geográfico de las especies y en su presencia y abundancia en un conjunto amplio de ecosistemas (Tilman *et al.*, 2002; Parmesan *et al.*, 2006; Reidsma *et al.*, 2006; Thomas, 2010; Eldridge *et al.*, 2011; Pauli *et al.*, 2012).

Un gran número de estudios empíricos y teóricos indican que la principal respuesta de las plantas al actual escenario de cambio global es modificar su distribución geográfica desplazándose a latitudes o altitudes mayores siguiendo su actual nicho ambiental, lo que en muchos casos implica una disminución del área de ocupación (Thuiller *et al.*, 2004; Thomas *et al.*, 2006; Thuiller *et al.*, 2008; Thomas, 2010; Gottfried *et al.*, 2012; Pauli *et al.*, 2012). Sin embargo, las respuestas específicas de las especies al cambio global son complicadas de predecir, ya que las diferentes especies no responden de igual manera a los motores de cambio y los factores que las condicionan se encuentran estrechamente relacionados (Figura 1). Enfrentarse al cambio desplazándose puede no ser factible para algunas especies porque la tasa de cambio ambiental será más rápida que su capacidad de dispersarse

siguiendo su nicho ambiental (Davis y Shaw *et al.*, 2001; Thuiller *et al.*, 2004; Thuiller *et al.*, 2005; Thomas *et al.*, 2006; Dullinger *et al.*, 2012b). Además, aunque la especie tenga capacidad para alcanzar un hábitat compatible con sus necesidades, los cambios no tienen por qué ocurrir de forma coordinada con otras especies (Memmot *et al.*, 2007; Tylianakis *et al.*, 2008; Hegland *et al.*, 2009; Svenning *et al.*, 2014). La viabilidad de las especies vegetales se encuentra condicionada en gran medida por la estructura y composición de las redes de interacciones bióticas de la comunidad a la que pertenecen (Tylianakis *et al.*, 2008; Hegland *et al.*, 2009; Svenning *et al.*, 2014). Por ello, otros factores relacionados con la interacción entre especies y sus efectos sobre parámetros demográficos y la configuración espacial de las comunidades vegetales podrían limitar su establecimiento en el nuevo ambiente (Brooker *et al.*, 2007; Svenning *et al.*, 2014). A consecuencia de estas limitaciones, existen diversos modelos predictivos que prevén un rápido incremento del riesgo de extinción a nivel mundial durante las próximas décadas (Tilman *et al.*, 2002; Thomas *et al.*, 2004; Thuiller *et al.*, 2005; Dullinger *et al.*, 2012a). En estos casos, y si la opción de desplazamiento es inviable, otra alternativa a la extinción es permanecer en el lugar ajustando el fenotipo a las nuevas condiciones ambientales mediante procesos de adaptación evolutiva o plasticidad fenotípica (Davis *et al.*, 2005; Miner *et al.*, 2005; Jump y Peñuelas, 2005; Gienapp *et al.*, 2008; Chuine *et al.*, 2010).

El concepto de adaptación evolutiva implica la actuación de una fuerza de selección que permite la evolución de un rasgo concreto, proporcionando una ventaja en un hábitat determinado (Reeve y Sherman, 1993; Kawecki y Ebert, 2004). La respuesta adaptativa a presiones selectivas direccionales derivadas de la aparición de cambios ambientales ha sido demostrada en un amplio rango de especies y sistemas, incluso en aquellos casos en los que los cambios ambientales ejercieron una presión selectiva muy intensa en poblaciones naturales (Turesson 1925; Clausen *et al.*, 1940; Linhart y Grant, 1996; Jump y Peñuelas, 2005; Parmesan 2006). La evolución implica por definición un cambio en las frecuencias alélicas, y por lo

tanto, resulta necesaria una diversidad genética adecuada y heredable, sobre la que puedan actuar las presiones selectivas (Kawecki y Ebert, 2004). La configuración espacial de los fragmentos de hábitat que contienen poblaciones de una especie y la acción de los agentes polinizadores y dispersores que residan en la comunidad determinan en gran medida los flujos genéticos entre poblaciones; lo que en última instancia, configura la estructura genética espacial de la especie y su variabilidad genética (Slatkin, 1987; Linhart y Grant, 1996; Hamrick y Trappnel, 2011). Además, la existencia de una elevada heterogeneidad ambiental en el nivel local puede impedir la existencia de presiones selectivas direccionales y dificultar la aparición de adaptaciones (Kawecki y Ebert, 2004).

La plasticidad fenotípica es la capacidad de un organismo de producir fenotipos diferentes en respuesta a cambios en su ambiente (Schlichting, 1986; Ghalambor *et al.*, 2007). Esta respuesta permite ajustes más rápidos y flexibles en los individuos, cuando la respuesta por una vía genética resulta insuficiente (Schlichting, 1986; Miner *et al.*, 2005). A largo plazo, las respuestas plásticas a cambios ambientales extremos tienen sus límites (Bradshaw y McNeilly, 1991; Jump y Peñuelas, 2005; Gienapp *et al.*, 2008). Sin embargo, en estos escenarios extremos, pueden promover el establecimiento y la persistencia en el nuevo ambiente situando a la especie cerca del nuevo fenotipo óptimo, lo que facilita que la evolución adaptativa ocurra en una escala de tiempo ecológica (Van Tienderen, 1997; Ghalambor *et al.*, 2007). La capacidad de ajustar el fenotipo a las nuevas condiciones ambientales está intrínsecamente ligada a una serie de procesos, tales como la tolerancia fisiológica, mecanismos a nivel demográfico u otros que implican cambios en las interacciones entre especies, y que por sí mismos, también pueden permitir a las poblaciones ser viables localmente sin necesidad de desplazarse (Doak y Morris, 2010; García-Camacho *et al.*, 2012; Lloret *et al.*, 2012; Svenning *et al.*, 2014).




Figura 1. Modelo conceptual de las respuestas específicas de las especies vegetales (elipses de color gris claro) a cambios ambientales derivados de los motores de cambio global. Los factores que inciden de forma más relevante sobre las respuestas aparecen representados por cuadros blancos. Algunas especies permanecerán en sus localidades originales adaptándose a las nuevas condiciones ambientales mediante procesos de adaptación evolutiva o plasticidad fenotípica. Otras modificarán su distribución geográfica al desplazarse a latitudes o altitudes mayores siguiendo el clima al cual están adaptadas, mientras que otras no podrán responder lo suficientemente rápido como para escapar de la extinción.

En definitiva, son numerosos los factores que pueden influir en la respuesta de las especies a los motores cambio global. Todos estos factores pueden actuar en la misma dirección o en direcciones opuestas y también pueden ver sus efectos potenciados o aminorados entre sí. Todo ello ocasiona que la respuesta de las especies a los cambios ambientales sea específica de cada especie, lo que dificulta

enormemente pronosticar el estado y la funcionalidad de los ecosistemas bajo los cambios físicos, biológicos y químicos que se están produciendo en la actualidad (Clark *et al.*, 2001; Higgins *et al.*, 2003; Svenning *et al.*, 2014). Es necesario, por tanto, adquirir una aproximación multidisciplinar para estudiar los principales determinantes de las respuestas de las plantas a los cambios ambientales acusados y evaluar su capacidad de respuesta ante un escenario de cambio global.

Vulnerabilidad de los ecosistemas de alta montaña mediterránea al cambio global

Las comunidades vegetales de alta montaña se encuentran entre los organismos más vulnerables al cambio global y particularmente a uno de sus motores más notables, el cambio climático (Thuiller *et al.*, 2005; Parmesan 2006; Nogues-Bravo *et al.*, 2007; Engler *et al.*, 2011). Pese a su escasa superficie, estos ecosistemas tiene un valor desproporcionado en términos de diversidad biológica y de servicios ecosistémicos (Körner *et al.*, 2007; Nogues-Bravo et al 2008; Nagy y Grabherr 2009; Barrio *et al.*, 2013), y es por ello que constituyen una prioridad de investigación. Su distribución se ve afectada de manera natural por la existencia de un gradiente altitudinal que pone gradualmente fin a las condiciones ambientales que permiten su existencia y marca los límites altitudinales de distribución. Esto otorga al ecosistema un carácter insular al configurar en el espacio una red de fragmentos de hábitat por encima de una determinada altitud, que supone el aislamiento de las poblaciones por encima del límite del bosque (Escudero *et al.*, 2004; Körner *et al.*, 2007; Nagy y Grabherr 2009). Esta situación restringe la capacidad de respuesta frente al cambio climático mediante migración latitudinal, y condiciona el flujo genético entre fragmentos, especialmente si la capacidad de dispersión es baja, lo que indirectamente también condiciona su respuesta mediante tolerancia o adaptación (Engler *et al.*, 2011; Dullinger *et al.*, 2012b).

Las evidencias de los cambios originados por el clima en las comunidades de alta montaña provienen principalmente de las regiones boreales y templadas. El

incremento de temperatura y precipitación asociado a la retirada de la nieve de forma temprana incrementa la producción primaria (Körner, 2003) y, en definitiva, el periodo de actividad vegetativa y la cantidad de recursos para poder invertir en reproducción. En general, la mayoría de los estudios han observado un patrón general de ascenso en altitud de especies de carácter más termófilo o peor adaptadas a los ambientes de alta montaña que implica incremento en la diversidad alfa acompañado de descensos en la diversidad beta. Esto ha supuesto una mayor homogeneidad de la comunidad en paralelo con un ascenso de los límites altitudinales superiores de las especies más orófilas de la comunidad (Paulí *et al.*, 2007; Jurasinski y Kreyling, 2007; Frei *et al.*, 2010; Gottfried *et al.*, 2012; Pauli *et al.*, 2012).

Las condiciones experimentadas por las comunidades vegetales en las montañas de la región mediterránea son marcadamente diferentes respecto a las de las regiones templadas y boreales, lo que provoca que el marco general de afección del cambio global sea diferente. Como en las formaciones de regiones más septentrionales, las montañas de gran altitud de la región mediterránea se ven sometidas a fríos inviernos que las cubren de una capa de nieve espesa que impide la actividad vegetativa en la época invernal (Körner *et al.*, 2007; Nagy y Grabherr 2009). Sin embargo, la sequía estival característica de climas mediterráneos, también se manifiesta en estas montañas. Esta situación reduce la ventaja asociada a un deshielo temprano para muchas especies tanto de carácter termófilo como orófilo, al restringir el periodo de crecimiento y reproducción al final del periodo estival (Escudero *et al.*, 2004; Giménez-Benavides *et al.*, 2007, 2011). Esta característica climática, ubicua a la región mediterránea, se puede ver agravada en el actual y futuro escenario de cambio climático, ya que la mayor parte de los modelos predictivos de clima pronostican que los cambios de temperatura y precipitación más intensos de Europa van a tener lugar en las montañas de clima mediterráneo (Nogues-Bravo *et al.*, 2008). Esto puede significar que las respuestas al cambio climático tengan que ser más intensas y rápidas que en otros sistemas montañosos

situados en regiones más septentrionales. A este efecto, además hay que asociarle los desequilibrios causados por la modificación de los usos del suelo motivada por el abandono de las prácticas ganaderas tradicionales. Este proceso está interactuando con el cambio climático para beneficiar el ascenso de las plantas leñosas por encima de sus límites altitudinales actuales (Wookey *et al.*, 2009; Brandt *et al.*, 2013). Un proceso de alcance global, pero que está afectando de manera significativa a las montañas mediterráneas (Peñuelas y Boada, 2003; Sanz-Elorza, 2003; García-Romero *et al.*, 2010). Finalmente, cabe destacar que las especies de carácter más orófilo ya habitan las cumbres más altas (Escudero *et al.*, 2004), por lo que resulta físicamente imposible responder mediante el desplazamiento de su límite superior (Marris, 2007). El resultado previsible es la extinción local de las poblaciones que habitan las alturas inferiores con la reducción del área de ocupación total de la especie (Marris, 2007). Existen estudios recientes que parecen confirmar esta idea, ya que han observado un patrón general de contracción del límite inferior de distribución y una reducción general de la riqueza de especies en las montañas mediterráneas del sur de Europa (Gottfried *et al.*, 2012; Paulí *et al.*, 2012).

Silene ciliata y Armeria caespitosa como especies de estudio representativas de las comunidades vegetales de alta montaña mediterránea

Dada la elevada vulnerabilidad de los ecosistemas de montaña mediterráneos al cambio climático y otros motores de cambio global, resulta de primordial interés evaluar su capacidad de respuesta frente al actual y futuro escenario de cambio global (Regato y Salman 2008). Sin embargo, existen pocos proyectos de investigación ecológica y conservación a largo plazo en estas zonas que incrementen el conocimiento sobre las especies que las habitan y los mecanismos que están gobernando las respuestas que permiten su supervivencia en el medio (Barrio *et al.*, 2013). Esta información es imprescindible para poder diseñar e interpretar estudios específicos sobre las respuestas a nivel de especie y/o comunidad al cambio global (Clark *et al.*, 2001; Thuiller *et al.*, 2004; Thuiller *et al.*, 2008; Svenning *et al.*, 2014).

Una excepción son los trabajos desarrollados durante la última década en la comunidad de pastos psicroxerófilos de alta montaña de la Sierra de Guadarrama (Sistema Central, España). Estos estudios se han centrado principalmente en dos especies representativas de la comunidad: *Armeria caespitosa* (Gómez Ortega) Boiss. In DC. y *Silene ciliata* Poiret, en las que se ha evaluado el cambio en las respuestas individuales (en términos vegetativos, reproductivos, demográficos y genéticos) a lo largo de gradientes altitudinales como un subrogado de lo que puede ocurrir cuando las condiciones climáticas cambian (revisado en Escudero *et al.*, 2012). *Silene ciliata* presenta dinámicas demográficas conducentes a la extinción en su límite inferior de distribución, dónde también se han detectado evidencias de una mayor intensidad de la depresión endogámica (Giménez-Benavides *et al.*, 2008, 2011a; García-Fernández *et al.*, 2012a). Paralelamente, en estas mismas poblaciones se han identificado procesos de adaptación local que permiten aumentar el éxito en el establecimiento de plántulas, uno de los cuellos de botella demográficos de la especie, lo que sin duda, contribuye a aumentar su éxito demográfico (Giménez-Benavides *et al.*, 2007a). También se han encontrado evidencias de adaptación local en individuos adultos según su población de origen frente a un periodo de sequía extrema y relativa tolerancia en la fase de plántula a la reducción del periodo de cobertura nival (García-Fernández *et al.*, 2013; García-Fernández *et al.*, 2014). Ambos resultados sugieren cierto nivel de tolerancia fisiológica a eventos climáticos extremos. Por otro lado, *Armeria caespitosa* parece estar respondiendo variando algunos rasgos relacionadas con el éxito reproductivo entre poblaciones (García-Camacho y Escudero, 2009; García-Camacho *et al.*, 2010), mediante un mecanismo de “compensación demográfica” que implica procesos de tolerancia fisiológica y morfológica o adaptativos (Doak y Morris, 2010, García-Camacho *et al.*, 2012). Finalmente, se han detectado presiones selectivas significativas hacia una floración más temprana en algunas poblaciones de ambas especies, aunque no se pudo determinar su potencial adaptativo (Giménez-Benavides *et al.*, 2011b).

De estos estudios, se deduce que *S. ciliata* y *A. caespitosa* podrían mostrar respuestas divergentes a las condiciones impuestas por el cambio climático (Escudero *et al.*, 2012). Existen, sin embargo, lagunas significativas en el conocimiento disponible sobre algunos de los factores que inciden en las respuestas de estas especies a los motores de cambio global. Alguno de estos factores serían los relacionados con las interacciones bióticas, la capacidad de dispersión o el potencial evolutivo de la variación genotípica y fenotípica encontrada en ambas especies. Resultan necesarias por tanto investigaciones adicionales que aborden estas lagunas y faciliten el pronóstico del estado y funcionalidad de las comunidades vegetales de alta montaña mediterránea en el contexto actual y futuro de cambio ambiental (Clark *et al.*, 2001; Higgins *et al.*, 2003; Svenning *et al.*, 2014).

OBJETIVOS

El objetivo general de esta tesis es completar el conocimiento sobre los principales factores que determinan la respuesta de las plantas de alta montaña mediterránea al cambio global. Para ello, se ha seleccionado la comunidad de pastos psicroxerófilos de la Sierra de Guadarrama, de la cual se conocen numerosos aspectos de la biología, estado demográfico y éxito reproductivo de dos de sus especies más representativas: *Silene ciliata* y *Armeria caespitosa*. La tesis ha seguido un enfoque integral que contemple conjuntamente el patrón espacial del hábitat junto a la capacidad de dispersión de las especies, las interacciones bióticas, la estructura genética espacial y la base genética de la variación fenotípica. Todos estos factores se evaluaron a lo largo de gradientes ambientales para recoger la elevada variabilidad ambiental a escala local que caracteriza a los ambientes de alta montaña y que puede afectar de manera significativa a los factores estudiados en la tesis.

Más concretamente, para alcanzar el objetivo general anteriormente citado, se han planteado ocho objetivos específicos que se abordan en los cinco capítulos de esta tesis doctoral:

- 1) Evaluar la capacidad de dispersión de *Silene ciliata* y *Armeria caespitosa* cuantificando la fecundidad y el kernel de dispersión efectiva a lo largo de un gradiente altitudinal (Capítulo 1).
- 2) Estimar la variación intraespecífica de la dispersión efectiva bajo las distintas condiciones ambientales impuestas por el gradiente altitudinal (Capítulo 1).
- 3) Cuantificar la estructura genética espacial en cinco poblaciones de *Silene ciliata* y evaluar si su intensidad está relacionada con las distintas condiciones de heterogeneidad espacial existentes y con su capacidad de dispersión (*i.e.*, densidad y patrón espacial) (Capítulo 2).

4) Determinar si existe relación entre la intensidad de la estructura genética espacial y los índices de diversidad genética y endogamia de las poblaciones estudiadas (Capítulo 2).

5) Analizar el patrón espacial de *S. ciliata* y evaluar la contribución que la dispersión, la heterogeneidad del hábitat y las interacciones intraespecífica tienen en la conformación del patrón observado (Capítulo 3).

6) Estudiar si la variación fenotípica observada en un conjunto de rasgos reproductivos y vegetativos en las poblaciones naturales de *Armeria caespitosa* tiene base genética (Capítulo 4).

7) Evaluar si las presiones selectivas hacia una floración más temprana detectadas en las poblaciones del límite inferior de distribución de *A. caespitosa* están promoviendo la predominancia de fenotipos de floración temprana (Capítulo 4).

8) Evaluar el efecto de la matorralización de los pastos de alta montaña en la red de interacción mutualista planta-polinizador (Capítulo 5).

METODOLOGÍA GENERAL

Área de estudio

Las localidades en las que se ha realizado el trabajo de campo, así como la totalidad de los individuos y especies vegetales de la presente tesis doctoral forman parte de las comunidades de pastizales psicroxerófilos de alta montaña de la Sierra de Guadarrama, una de las más importantes del Sistema Central (Figura 2).




Figura 2. Situación de las localidades de estudio en la Sierra de Guadarrama (España). Las áreas sombreadas con tonos más oscuros indican mayor altitud. El mapa de la esquina superior izquierda indica la localización de la Sierra de Guadarrama en la Península Ibérica. Los números indican la posición de cada localidad y las clasifica de menor a mayor altitud (Ver tabla 1): (1) Sierra de los Porrones, (2) Collado de las Vacas, (3) Loma de Cabezas, (4) Laguna, (5) Najarra, (6) Nevero, (7) Cabezas de Hierro, (8) Peñalara.

La citada comunidad vegetal constituye la vegetación potencial de las cumbres de la Sierra de Guadarrama por encima de los 2100-2200 metros. La comunidad está dominada por la gramínea *Festuca curvifolia* Lag. ex Lange y un importante número de nanocaméfitos almohadillados o pulviformes (Figura 3 y 4). Desde un punto de vista fitosociológico la comunidad ha sido denominada como *Hieracio myriadeni-Festucetum curvifoliae* (Rivas-Martinez, 1963) y es endémica de las cumbres de la Sierra de Guadarrama (Rivas-Martinez *et al.*, 1990). La comunidad es particularmente rica en endemismos, como *Festuca curvifolia* Lag. ex Lange, *Pilosella vahlii* (Froel.) F.W. Schultz & Sch.Bip., *Armeria caespitosa* (Gómez Ortega) Boiss. *in DC.* o *Minuartia recurva* (All.) Schinz & Thell. Algunas especies relictas de origen artico-pirenaicas como *Agrostis rupestris* All. y *Phyteuma hemisphaericum* L. enriquecen aún más la composición de especies (Escudero *et al.*, 2004). La comunidad convive intercalada en una matriz arbustiva más o menos densa caracterizada por *Cytisus oromediterraneus* Rivas Mart. *et al.*, *Adenocarpus hispanicus* (Lam. DC.) y *Juniperus communis* subsp. *alpina* (Suter) Celak. Se diferencia de las comunidades genuinamente alpinas por presentar un periodo de sequía estival intensa característico de la alta montaña mediterránea, lo que, unido a la fuerte insolación recibida, confiere a las especies que los habitan caracteres propios de la vegetación xerofítica: cutículas gruesas, colores glaucos, abundancia de especies leñosas (Rivas-Martinez *et al.*, 1990). A consecuencia de su singularidad y riqueza florística, así como por el reducido área de distribución esta comunidad se encuentra incluida dentro de la Directiva Habitats de la Unión Europea (Directiva 92/43/CEE). Más información sobre la composición florística de esta comunidad puede consultarse en Escudero *et al.*, (2004).



Figura 3. Aspecto del pastizal psicroxerófilo en las proximidades del Alto de las Guerramillas (2.200 m). Los procesos de crioturbación y las condiciones extremas confieren escasa cobertura y porte al pastizal (Fotografía: Carlos Lara).



Figura 4. Ejemplares de *Armeria caespitosa* y *Festuca curvifolia* en las proximidades de la cumbre de Peñalara (2.430 m) (Fotografía: Alfredo García).

La estación meteorológica más próxima a la zona se encuentra en el Puerto de Navacerrada ($40^{\circ} 46'N$, $4^{\circ}19'O$; 1890 msnm). La precipitación media anual registrada es de 1330 mm y la temperatura media anual es de $6,3^{\circ}C$ (<http://aemet.es>). La temperatura media mensual oscila desde $-1^{\circ}C$ en Enero hasta $16^{\circ}C$ en Julio. Existe un acusado periodo seco en verano (Junio – Septiembre), con menos del 10% de las precipitaciones, que reduce drásticamente el periodo vegetativo (Giménez-Benavides *et al.*, 2007b). La caracterización ecogeográfica de las localidades de estudio se incluye en la tabla 1 y figura 2. Todas las localidades se encuentran incluidas dentro del Parque Nacional de la Sierra de Guadarrama.

Tabla 1. Caracterización ecogeográfica de las localidades de estudio.

Localidad de estudio	Coordinadas geográficas	Altitud (m)	Pm (mm)	Tm ($^{\circ}C$)
Porrones	$40^{\circ} 45'N$, $3^{\circ} 56'O$	1750	1133 (25 – 130)	8,2 (5,2 – 18,3)
Collado de las Vacas	$40^{\circ} 6'N$, $3^{\circ} 56'O$	1880	1210 (28 – 142)	7,3 (-0,9 – 18,9)
Loma de Cabezas	$40^{\circ} 46'N$, $3^{\circ} 56'O$	1950	1351 (30 – 153)	5,7 (-0,7 – 15,1)
Laguna	$40^{\circ} 50'N$, $3^{\circ} 57'O$	1980	1347 (31 – 179)	6,6 (-1,1 – 14,8)
Najarra	$40^{\circ} 49'N$, $3^{\circ} 49'O$	2080	1335 (27 – 181)	7,3 (-0,8 – 15,7)
Nevero	$40^{\circ} 58'N$, $3^{\circ} 50'O$	2190	1245 (30 -144)	6,6 (-1,3 – 14,6)
Cabezas de Hierro	$40^{\circ} 48'N$, $3^{\circ} 55'O$	2305	1594 (34 – 193)	4,2 (-2,8 – 13,7)
Peñalara	$40^{\circ} 51'N$, $3^{\circ} 57'O$	2405	1457 (32 – 183)	4,9 (-2,6 – 13,8)

Pm: precipitación anual media, *Tm:* temperatura anual media. Las temperaturas y precipitaciones mínimas y máximas mensuales se muestran entre paréntesis. Datos climáticos obtenidos de Gonzalo-Jiménez (2011).

Especies de estudio: *Silene ciliata* y *Armeria caespitosa*

Silene ciliata Poiret (Caryophyllaceae) es un caméfito pulviniforme perene que se distribuye en los sistemas de montaña mediterráneos del sur de Europa, desde la Península Ibérica hasta los Balcanes (Tutin, 1995). La especie alcanza uno de sus límites de distribución más meridionales en las cumbres de la Sierra de Guadarrama, donde habita un área restringida entre 1900 y 2430 m de altitud. Tiene un porte de pequeño tamaño, alcanzando los 25-30 cm de diámetro, mientras que las inflorescencias con 1 a 5 flores alcanzan hasta 15 cm de altura (Figura 5). Es una especie geitonogámica (en la planta aparecen flores hermafroditas y femeninas) con una fuerte protandria (estructuras femeninas se desarrollan posteriormente a las masculinas), lo cual limita la posibilidad de autocruzamientos, aunque la planta es autocompatible (García-Fernández *et al.*, 2012b). Tiene síndrome de polinización nocturna, por lo que es polinizada principalmente por polillas (Lepidoptera: Noctuidae), aunque también puede ser polinizada por insectos diurnos, principalmente sírfidos (Himenoptera: Syrphidae). Florece a finales del verano, con un pico de floración en Agosto. El fruto es una cápsula dehiscente que contiene hasta 100 semillas dispersadas por viento entre agosto y septiembre. Las semillas, que no tienen ninguna estructura específica para promover su dispersión, tienen una masa media de 0,59 mg y su diámetro oscila entre 1,1 y 1,5 mm (Giménez-Benavides *et al.*, 2005). Su dotación cromosómica en la Sierra de Guadarrama es diploide ($2n=24$, García-Fernández *et al.*, 2012b).



Figura 5. Ejemplar de *Silene ciliata* con las flores abiertas durante la noche. Debido a su síndrome de polinización nocturna, la especie es polinizada principalmente por polillas (Fotografía: Luis Giménez).

Armeria caespitosa (Gómez Ortega) Boiss. in DC. es un caméfito pulviniforme de escaso porte que pertenece a la familia Plumbaginaceae. Es una planta endémica del Sistema Central Ibérico (Sierra de Guadarrama, Ayllón y Este de Gredos), apareciendo entre los 1600 y 2430 m de altitud (Nieto-Feliner, 1996). Se desarrolla en dos hábitats bien diferenciados, pero que coexisten: en pastos psicroxerófilos sobre litosuelos pobres, poco desarrollados, y en fisuras y pequeñas repisas de afloramientos rocosos de granito y gneis (Figura 6). Sus flores rosadas se agrupan en pequeños capítulos címosos (19 ± 8 flores por capítulo) sustentados por escapos cortos. Cada flor contiene un único ovulo, que produce una única semilla que se dispersa entre junio y agosto dentro del cáliz, engrosado durante el proceso de maduración. El peso medio de la semilla es de 1.18 mg y su diámetro oscila entre 2 y 4,7 mm (Giménez-Benavides *et al.*, 2005). Es una de las especies de floración más temprana de la comunidad, abarcando desde finales de abril en las poblaciones de menor altitud hasta finales de junio en las poblaciones más elevadas (García-Camacho y Escudero, 2009). Es polinizada por insectos generalistas, principalmente

abejas (Himenoptera: Apoidea) y sírfidos (Himenoptera: Syrphidae). Tiene dotación diploide y, como la mayoría de las especies del género *Armeria* es autoincompatible estricta, debido a un sistema dimórfico de autoincompatibilidad (García-Camacho y Escudero, 2009).



Figura 6. Principales microhábitats que ocupa *A. caespitosa*. Foto superior: individuos en afloramiento rocoso. Foto inferior: individuos en pasto psicroxerófilo de *Festuca curvifolia* (Fotografía: Raúl García Camacho).

Técnicas de análisis estadístico

En cada capítulo de esta memoria se incluye información detallada sobre la metodología y técnicas de análisis estadístico utilizadas para responder a los objetivos concretos planteados. Este apartado incluye una breve descripción de las principales técnicas utilizadas.

Modelización inversa y funciones kernel de dispersión

La fecundidad de las plantas adultas reproductoras y la distribución de las distancias de dispersión son características fundamentales del proceso de dispersión (Ribbens *et al.*, 1994; Clark *et al.*, 1999; Nathan y Muller-Landau, 2000; Bullock *et al.*, 2006). Sin embargo, en muchos sistemas es un reto estimar la producción de semillas y las distancias de dispersión porque la trayectoria de las semillas dispersadas no puede ser establecida, y como consecuencia no se puede determinar la planta de origen (Bullock *et al.*, 2006). La modelización inversa aplica métodos de máxima verosimilitud para estimar la fecundidad de las plantas adultas y la función kernel de dispersión sin identificar la fuente exacta de la que procede cada semilla o plántula (Ribbens *et al.*, 1994; Clark *et al.*, 1999). Esta técnica asume que la contribución de cada planta adulta a la lluvia de semillas en un área dada depende sólo de la fecundidad de la planta y de su distancia a dicha área (Ribbens *et al.*, 1994; Clark *et al.*, 1999, Nathan *et al.*, 2012).

La curva o “kernel” de dispersión es una función de probabilidad de densidad que caracteriza la distribución espacial de los propágulos (semillas o plántulas) dispersados desde una fuente común (Nathan *et al.*, 2012). La mayoría de las funciones kernel de dispersión que se aplican en ecología se definen en base al parámetro de forma u y de escala a . u describe la forma de la curva del kernel de dispersión; mientras que a proporciona la extensión espacial a la que se producen los eventos de dispersión (Nathan *et al.*, 2012).

Herramientas de análisis basados en patrones de puntos

El patrón espacial de organización de las comunidades vegetales está recibiendo un interés creciente en ecología en los últimos 20 años (Wiegand *et al.*, 2012). Este interés se debe principalmente a que a través del análisis de estos patrones y de su dinámica podemos entender los procesos subyacentes que los están generando y testar hipótesis ecológicas (Legendre y Fortin 1989; Legendre *et al.*, 2002; Wiegand *et al.*, 2012). Los patrones de puntos son definidos como la posición geográfica del conjunto individuos de una especie en un área (Wiegand *et al.*, 2012). El fundamento que subyace detrás de las técnicas de análisis de patrones de puntos es que algunos procesos ecológicos tienen un fuerte componente espacial que crea patrones espaciales identificables (Brown *et al.*, 2011; Wiegand *et al.*, 2012). El estadístico espacial más utilizado es la función K de Ripley ($K(r)$, Ripley, 1976), que es el número de puntos (*i.e.*, plantas) dentro de un círculo de radio r alrededor de cualquier punto del patrón. Otros estadísticos básicos ampliamente usados en ecología son la función de correlación de par $g(r)$, derivada de $K(r)$ (Stoyan y Stoyan, 1994); y la función de la distancia al vecino más cercano $D(r)$, que estima la probabilidad de tener la planta más próxima dentro de una distancia r (Loosmore y Ford, 2006).

Análisis de autocorrelación espacial genética

Los análisis de autocorrelación espacial genética permiten analizar si la diversidad genética se encuentra espacialmente estructurada (*i.e.*, distribución no aleatoria de los genotipos en el espacio). El interés de la estructura espacial de la diversidad genética se debe a que la distribución no aleatoria de genotipos en el espacio influencia varios procesos evolutivos como la pérdida de diversidad genética por endogamia, el tamaño efectivo poblacional o los procesos de evolución adaptativa. Como consecuencia, puede tener un efecto significativo en la eficacia biológica de los individuos y las poblaciones (Linhart y Grant, 1996; Escudero *et al.*, 2004). Los análisis de autocorrelación espacial genética permiten evaluar la

existencia de la estructura espacial genética y cuantificarla (Smouse y Peakall, 1999). El estadístico utilizado para establecer el grado de emparentamiento genético en los análisis de autocorrelación espacial ha sido el coeficiente de parentesco genético, F_{ij} , propuesto por Loiselle *et al.*, (1995). Se define como la probabilidad de que un gen muestreado al azar de un individuo i sea idéntico a un gen muestreado al azar del individuo j (Vekemans y Hardy, 2004).

Experimentos de jardín común

Cultivar las plantas en un único ambiente permite controlar el efecto de factores ambientales en la expresión de los caracteres de interés, asegurando que cualquier diferencia observada entre los tratamientos experimentales tiene base genética (Turesson, 1922; Clausen *et al.*, 1948; Kawecki y Ebert, 2004). Los experimentos de jardín común (*common garden*, en inglés) utilizan esta técnica para detectar diferenciación genética en rasgos fenotípicos de plantas procedentes de diferentes regiones o poblaciones (Turesson, 1922; Clusen *et al.*, 1948). No son adecuados para establecer sin incertidumbre el origen de la variación genética, por lo que no permiten establecer si las diferencias en la expresión de los caracteres estudiados han sido causadas por fuerzas evolutivas neutrales (*e.g.*, deriva genética) o por procesos adaptativos (Kawecki & Ebert 2004). Sin embargo, una respuesta diferencial en función de la población de origen a diferentes condiciones experimentales (*e.g.*, diferentes tratamientos de riego), o una relación estadística significativa entre los rasgos fenotípicos medidos en el jardín común y las condiciones ambientales de los sitios de origen pueden indicar adaptación (Linhart y Grant 1996).

Redes complejas de interacciones mutualistas

El enfoque de redes complejas al estudio de interacciones ecológicas parte del supuesto de que no es posible comprender el funcionamiento de las comunidades ecológicas a partir del estudio individualizado de las especies que lo componen, sino que es necesario estudiar la manera en que se organizan las interacciones entre las

especies (Jordano *et al.*, 2003). Las redes de interacciones mutualistas pueden ser representadas mediante la utilización de matrices de interacción (Jordano, 1987). En ellas, los vectores fila representan especies del nivel trófico superior (*e.g.*, polinizadores o frugívoros) y los vectores columna representan especies del nivel trófico inferior (*e.g.*, plantas con flores o frutos carnosos). Para cada especie animal o de planta, podemos mantener un registro de las especies con las que interactúa y la intensidad con la que se produce la interacción. A partir de la información contenida en estas matrices, podemos describir las redes de interacción mediante un conjunto de métricas y estadísticos que aportan información sobre la estructura, funcionamiento y estabilidad de la comunidad estudiada (revisado en Bascompte y Jordano, 2007). La tesis doctoral se ha centrado en las redes de interacción planta-polinizador. Como consecuencia de su potencial, estas redes han sido ampliamente utilizadas para analizar la estabilidad de los ecosistemas (Bascompte *et al.*, 2006), las consecuencias de las perturbaciones originadas por el cambio climático (Hegland *et al.*, 2009), el efecto de la introducción de especies invasoras en los ecosistemas (Aizen *et al.*, 2008, Bartomeus *et al.*, 2008; Traveset *et al.*, 2013) y el éxito de la restauración ecológica de ecosistemas degradados (Devoto *et al.*, 2012).

ESTRUCTURA DE LA TESIS

La tesis se estructura en cinco capítulos que han sido escritos en inglés para su publicación en revistas científicas de ámbito internacional. Por ello, se presentan los manuscritos originales de dichos artículos. A continuación se incluye la traducción del título al español, la lista de coautores y el estado de publicación de cada capítulo y una breve síntesis de cada uno de ellos.

Capítulo 1. Evaluación de la variación intraespecífica de la dispersión efectiva a lo largo de un gradiente altitudinal: un análisis en dos especies de alta montaña mediterránea.

Carlos Lara-Romero, Juan J. Robledo-Arnuncio, Alfredo García-Fernández, José M. Iriondo.

Manuscrito publicado en PLoS ONE 9(1): e87189.

Las condiciones ambientales y su variación espacial controlan en gran medida los patrones de reclutamiento de las comunidades vegetales. Para poder entender cómo esta variación ambiental afecta a los procesos que controlan el reclutamiento es necesario adoptar un enfoque multi-poblacional. Hemos utilizado una aproximación de modelización inversa para evaluar los patrones de reclutamiento en cinco poblaciones de *Silene ciliata* y *Armeria caespitosa* a lo largo de un gradiente altitudinal. Ambas especies son representativas de uno de los ecosistemas ibéricos más vulnerables al cambio climático: la comunidad de pastos psicroxerófilos de alta montaña mediterránea. Consideramos cuatro kernels de dispersión de semilla diferentes: Exponencial-Potencial, 2Dt, WALD y Log-normal. Los resultados mostraron una distancia media de dispersión reducida (<1m) en todas las poblaciones. Ambas especies mostraron variación interpoblacional en el tipo de kernel que proporciona mejores ajustes y en el parámetro de fecundidad. Sin embargo, solo *A. caespitosa* exhibió diferencias significativas entre poblaciones en la distancia media de dispersión. La variación ambiental altitudinal condicionó la fecundidad de los individuos en ambas especies. Junto a la limitación por dispersión

primaria, la dispersión efectiva podría estar limitada por procesos post-dispersivos relacionados con la sequía estival. Estos resultados resaltan la complejidad de la variación intraespecífica en los procesos que controlan el reclutamiento, mostrando que el kernel de dispersión puede permanecer relativamente invariante entre poblaciones de una misma especie, incluso si existe fuerte variación en la estructura demográfica y/o el ambiente físico entre poblaciones; mientras que la asunción de invariancia intraespecífica del kernel de dispersión podrían no mantenerse para otras especies en el mismo ambiente. Por lo tanto, los resultados de este estudio sugieren la necesidad de análisis específicos en un amplio espectro de especies y ambientes para evaluar la prevalencia y magnitud de la variación intraespecífica de la dispersión. La baja distancia de dispersión de estas especies podría restringir la capacidad migratoria frente al cambio climático y limitar el flujo genético entre poblaciones.

Capítulo 2. Variación de la estructura genética espacial a escala local entre poblaciones de *Silene ciliata* (Caryophyllaceae) con diferente grado de agregación espacial.

Carlos Lara-Romero, Alfredo García-Fernández, Juan J. Robledo-Arnuncio, Javier Morente-López, Ángela López, Marie Roumet, José M. Iriondo.

Manuscrito en revisión en *Molecular Ecology*.

El patrón y la intensidad de la estructura espacial de la variación genética a escala local (FSGS por sus siglas en inglés) pueden variar entre poblaciones de una especie como consecuencia de la interacción entre diferentes factores demográficos y ambientales. Los modelos teóricos predicen que en hábitats o ambientes heterogéneos, la intensidad de la FSGS se incrementará como consecuencia de la variación local de la densidad. Sin embargo, existen pocos trabajos empíricos que hayan examinado las diferencias en la intensidad de la FSGS entre poblaciones con diferente grado de heterogeneidad espacial. En este estudio usamos métodos de autocorrelación espacial para cuantificar la estructura espacial demográfica y

genética en cinco poblaciones estables demográficamente pero espacialmente heterogéneas de *Silene ciliata*. Encontramos FSGS significativa en tres de las cinco poblaciones estudiadas, confirmando la existencia de variación intra-poblacional en la intensidad de la estructura espacial genética. Se detectó una asociación positiva entre la intensidad de la FSGS y el nivel de agregación espacial de las plantas. Este resultado avala a los estudios teóricos que muestran que los patrones espaciales agregados tienden a aumentar los niveles de FSGS en comparación con distribuciones aleatorias. Contrariamente a lo observado en otras especies, la población con mayor densidad de plantas fue también la población con mayor intensidad de FSGS, probablemente debido a una reducida distancia de dispersión de polen. Estos resultados muestran el potencial que tiene incorporar explícitamente información sobre la distribución espacial a escala local de los individuos y el hábitat en estudios de la variación espacial de la diversidad genética.

Capítulo 3. Los patrones de distribución espacial de *S. ciliata*, condicionados por la dispersion y la heterogeneidad ambiental, revelan densodependencia negativa en el éxito reproductivo pese a la existencia de asociación espacial positiva.

Carlos Lara-Romero, Marcelino de la Cruz, Gema Escribano-Ávila, Alfredo García-Fernández, José M. Iriondo.

Manuscrito inédito.

Los patrones de distribución espacial en plantas pueden estar generados por un conjunto amplio de factores ecológicos, lo que dificulta distinguir la contribución que los efectos denso-dependientes tienen en la conformación de los patrones espaciales observados, respecto a otros factores como la existencia de heterogeneidad ambiental o la dispersión. En este estudio analizamos la estructura espacial de *Silene ciliata* a lo largo de un gradiente altitudinal y evaluamos la contribución que la dispersión, la heterogeneidad del hábitat y las interacciones intraespecífica tienen en la conformación del patrón observado. Primero utilizamos

funciones de correlación de par $g(r)$ homogéneas y heterogéneas para describir el patrón espacial de las plántulas y adultos en cada población y evaluar la contribución de la heterogeneidad ambiental al patrón observado. A continuación, utilizamos un diseño de “caso-control” junto a un modelo nulo de “etiquetado aleatorio” para evaluar si el patrón espacial resultante del proceso de dispersión y establecimiento de plántulas persiste en la clase adulta, así como para analizar la relación espacial entre ambas clases de edad (*i.e.*, agregación o repulsión). Finalmente, analizamos el efecto de la densidad de conspecíficos en la producción de flores de cada planta, con el objetivo de evaluar si existe denso-dependencia en el éxito reproductivo. Tanto las plántulas como los adultos mostraron una estructura espacial agregada que parece estar controlada por la heterogeneidad del hábitat y el patrón de dispersión y establecimiento de semillas. Además pese a la tendencia de la especie a establecerse de forma agregada detectamos evidencias de densodependencia negativa en su éxito reproductivo. Los resultados en su conjunto demuestran que los patrones de co-ocurrencia en agregación no tienen por qué estar mediados por interacciones de facilitación, por lo que antes de establecer relaciones causales entre éstas y la distribución espacial de las plantas, hay que evaluar otros factores como la dispersión o la heterogeneidad ambiental.

Capítulo 4. Variación genética en la fenología y el éxito reproductivo en un especialista de alta montaña mediterránea.

Carlos Lara-Romero, Raúl García-Camacho, Adrian Escudero, Jose M. Iriondo.

Manuscrito aceptado en *Botanical Journal of the Linnean Society*.

Las respuestas adaptativas al clima pasado podrían ser importantes para la persistencia de las plantas de alta montaña en el actual escenario de cambio climático acelerado. *Armeria caespitosa* es una planta de alta montaña, endémica del Sistema Central Ibérico. Las diferencias encontradas en el ambiente abiótico a lo largo del gradiente altitudinal someten a la especie a dos gradientes de estrés

opuestos relacionados con el estrés hídrico y la duración de la temporada de crecimiento vegetativo. Además, la especie tiene la capacidad de subsistir en dos micro hábitats contrastados (afloramientos rocosos y pastos psicroxerófilos) que, dependiendo de la altitud, tienen diferentes efectos sobre su dinámica poblacional. Esta variación ambiental provoca que la especie muestre gran diferenciación inter-poblacional en un conjunto de rasgos reproductivos y vegetativos. En este estudio, utilizamos una aproximación de jardín común para evaluar si la variación fenotípica observada en las poblaciones naturales tiene una base genética, o si por el contrario, es el resultado de respuestas plásticas moduladas por el conjunto condiciones ambientales heterogéneas descrito anteriormente. Las plantas procedentes del límite de elevación superior y de los pastos psicroxerófilos florecieron antes y produjeron más frutos viables, pero crecieron menos en comparación con las plantas procedentes del límite de elevación inferior y de los afloramientos rocosos. Estos resultados confirman que la variación inter-poblacional observada en algunos de los rasgos ligados a la fenología y el éxito reproductivo tiene base genética. El estudio también mostró que la respuesta selectiva en favor de una floración temprana detectada en el límite de elevación inferior, no ha generado una mayor proporción de fenotipos de floración temprana. Sin embargo, la variabilidad genética asociada al inicio de la floración, podría ser relevante para hacer frente al impacto ocasionado por el cambio climático en curso.

Capítulo 5. Efectos directos e indirectos de la matorralización de los pastos de alta montaña sobre la red de interacciones planta-polinizador.

Carlos Lara-Romero, Cristina García, Javier Morente-López, José M. Iriondo.

Manuscrito inédito.

Las interacciones mutualistas son claves para la estructura y el funcionamiento de las comunidades ecológicas y son especialmente sensibles a los motores de cambio global. En este estudio utilizamos técnicas de análisis redes complejas para

cuantificar el efecto de la matorralización de los pastos de alta montaña en la red de interacciones planta-polinizador. Realizamos el estudio en los pastos psicroxerófilos de la Sierra de Guadarrama porque el ascenso del matorral motivado por el cambio global está ocasionando una reducción del hábitat disponible para la comunidad que los habita, lo que constituye una de las mayores amenazas para su persistencia. Construimos matrices de interacción cuantitativas para comparar parcelas matorralizadas frente a parcelas en las que el matorral no estaba presente. Basándonos en la hipótesis de la Variación del Nicho (HVN) y la hipótesis de la Diversidad Funcional (HDF), nos preguntamos si el incremento en el recurso nutricional, como consecuencia de la expansión de los matorrales, modifica la amplitud de nicho y la riqueza de los polinizadores; así como las consecuencias de estos cambios para las interacciones interespecíficas entre plantas. Como predice la HDF, la matorralización de los pastos estudiados tuvo un efecto positivo sobre la riqueza y la frecuencia de visitas de los polinizadores a las plantas, pero no modificó su amplitud de nicho, contrariamente a las expectativas de la HVN. Por otro lado, la matorralización incrementó la competencia entre plantas por los servicios de polinización. Esta última situación podría reducir el éxito reproductivo en alguna de las especies de plantas estudiadas. Un amplio número de grupo funcionales de polinizadores visitaron a las especies de matorral en mayor proporción que a la mayoría de las especies de plantas de la comunidad, lo que otorgó a los matorrales un rol de súper-generalista. Pese a ello, la estructura de la red de interacciones planta-polinizador apenas mostró cambios significativos en sus propiedades estructurales, aunque la red de los pastos matorralizados fue ligeramente más generalista. Este resultado coincide con estudios anteriores y sugiere que el mutualismo planta-polinizador podría ser relativamente permeable y robusto a las modificaciones provocadas por los motores de cambio global. Este estudio resalta la necesidad de utilizar un enfoque integrador en los estudios del efecto de los motores de cambio global sobre las interacciones bióticas y la estructura y el funcionamiento de las comunidades ecológicas.

CONCLUSIONES

De los cinco capítulos en los que se ha estructurado esta tesis doctoral se pueden extraer las siguientes conclusiones:

- 1) La distancia media de dispersión efectiva de *Silene ciliata* y *Armeria caespitosa* es reducida (0,23 – 0,71 m). Junto a la limitación por dispersión primaria, la dispersión efectiva podría estar limitada por procesos post-dispersivos relacionados con la sequía estival. Esta situación podría restringir la capacidad migratoria frente al cambio climático y limitar el flujo genético entre poblaciones.
- 2) La variación ambiental altitudinal condicionó la fecundidad de los individuos *S. ciliata* y *A. caespitosa* pero no afectó a la escala espacial a la que se produce la dispersión. Ambas especies mostraron variación inter-poblacional en el tipo de kernel que proporciona mejores ajustes y en el parámetro de fecundidad. Sin embargo, solo *A. caespitosa* exhibió diferencias significativas entre poblaciones en la distancia media de dispersión. Estos resultados sugieren la necesidad de análisis específicos en un amplio espectro de especies y ambientes para evaluar la prevalencia y magnitud de la variación intraespecífica de la dispersión.
- 3) Se ha detectado una asociación positiva entre la intensidad de la estructura genética espacial (EGS) y el nivel de agregación espacial de las poblaciones de *Silene ciliata*. Este resultado avala a los estudios teóricos que muestran que los patrones espaciales agregados tienden a aumentar los niveles de EGS en comparación con distribuciones aleatorias.
- 4) Los índices de diversidad genética y endogamia observados en las poblaciones estudiadas de *Silene ciliata* fueron homogéneos. Consecuentemente, no se encontró relación entre la intensidad de la EGS y dichos parámetros.
- 5) El patron de agregación de *S. ciliata* parece estar controlada por la heterogeneidad del hábitat y el patrón de dispersión y establecimiento de semillas, mientras que el efecto de las interacciones intraespecífica sobre el patrón es

reducido. Estos resultados minimizan la importancia de las interacciones intraespecífica para el patrón espacial de la especie, a la vez que destacan la importancia de evaluar todos los factores que promueven la co-ocurrencia agregada de las plantas si queremos conocer plenamente la dinámica de las poblaciones.

6) La variación inter-poblacional observada en el inicio de la floración y en el número de inflorescencias y de frutos viables en poblaciones naturales de *Armeria caespitosa* tiene base genética.

7) La respuesta selectiva en favor de una floración temprana detectada en el límite de elevación inferior de distribución de *A. caespitosa* no ha generado una mayor proporción de fenotipos de floración temprana. Sin embargo, la variabilidad genética asociada al inicio de la floración, podría ser relevante para hacer frente al impacto ocasionado por el cambio climático en curso debido a su potencial adaptativo.

8) Como predice la hipótesis de la Diversidad Funcional, la matorralización de los pastos estudiados tuvo un efecto positivo sobre la riqueza y la frecuencia de visitas de los polinizadores a las plantas, pero no modificó su amplitud de nicho, contrariamente a las expectativas de la hipótesis de la Variación del Nicho, lo que incrementó la competencia entre plantas por los servicios de polinización. Esta última situación podría reducir el éxito reproductivo en alguna de las especies de plantas estudiadas.

BIBLIOGRAFÍA

- Aizen MA, Morales CL, Morales JM. 2008. Invasive mutualists erode native pollination webs. *PLoS biology*, 6: e31.
- Barrio IC, Bueno CG, Nagy L, Palacio S, Grau O, Munilla I, García M-B, Garcia-Cervign AI, Gartzia M, Gazol A. 2013. Alpine Ecology in the Iberian Peninsula: What Do We Know, and What Do We Need to Learn? *Mountain Research and Development*, 33: 437-442.
- Bartomeus I, Vilá M, Santamaría L. 2008. Contrasting effects of invasive plants in plant-pollinator networks. *Oecologia*, 155: 761-770.
- Bascompte J, Jordano P. 2007. Plant-animal mutualistic networks: the architecture of biodiversity. *Annual Review of Ecology, Evolution, and Systematics*: 567-593.
- Bradshaw AD, McNeilly T. 1991. Evolutionary response to global climatic change. *Annals of Botany*, 67: 5-14.
- Brandt JS, Haynes MA, Kuemmerle T, Waller DM, Radeloff VC. 2013. Regime shift on the roof of the world: Alpine meadows converting to shrublands in the southern Himalayas. *Biological Conservation*, 158: 116-127.
- Brooker RW, Travis JMJ, Clark EJ, Dytham C. 2007. Modelling species-range shifts in a changing climate: the impacts of biotic interactions, dispersal distance and the rate of climate change. *Journal of Theoretical Biology*, 245: 59-65.
- Brown C, Law R, Illian JB, Burslem DFRP. 2011. Linking ecological processes with spatial and non-spatial patterns in plant communities. *Journal of Ecology*, 99: 1402-1414.
- Bullock JM, Shea K, Skarpaas O. 2006. Measuring plant dispersal: an introduction to field methods and experimental design. *Plant Ecology*, 186: 217-234.
- Clark JS, Carpenter SR, Barber M, Collins S, Dobson A, Foley JA, Lodge DM, Pascual M, Pielke Jr R, Pizer W. 2001. Ecological forecasts: an emerging imperative. *Science*, 293: 657-660.
- Clark JS, Silman M, Kern R, Macklin E, HilleRisLambers J. 1999. Seed dispersal near and far: patterns across temperate and tropical forests. *Ecology*, 80: 1475-1494.
- Clausen J, Keck DD, Hiesey WM. 1948. *Environmental responses of climatic races of Achillea. Experimental studies on the nature of species*. Washington D.C: Carnegie Institute of Washington.
- Chuine I. 2010. Why does phenology drive species distributions? *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 365: 3149-3160.
- Davis MB, Shaw RG. 2001. Range shifts and adaptive responses to Quaternary climate change. *Science*, 292: 673-679.
- Davis MB, Shaw RG, Etterson JR. 2005. Evolutionary responses to changing climate. *Ecology*, 86: 1704-1714.
- Devoto M, Bailey S, Craze P, Memmott J. 2012. Understanding and planning ecological restoration of plant-pollinator networks. *Ecology letters*, 15: 319-328.
- Doak DF, Morris WF. 2010. Demographic compensation and tipping points in climate-induced range shifts. *Nature*, 467: 959-962.

- Dullinger S, Gatterer A, Thuiller W, Moser D, Zimmermann NE, *et al.*, 2012a. Extinction debt of high-mountain plants under twenty-first-century climate change. *Nature Climate Change*, 2: 619-622.
- Dullinger S, Willner W, Plutzar C, Englisch T, Schratt-Ehrendorfer L, Moser D, Ertl S, Essl F, Niklfeld H. 2012b. Post-glacial migration lag restricts range filling of plants in the European Alps. *Global Ecology and Biogeography*, 21: 829-840.
- Engler R, Randin CF, Thuiller W, Dullinger S, Zimmermann NE, *et al.*, 2011. 21st century climate change threatens mountain flora unequally across Europe. *Global Change Biology*, 17: 2330-2341.
- Escudero A, García-Camacho R, García Fernández A, Gavilán RG, Giménez-Benavides L, Iriondo JM, Lara-Romero C, Morente J, Pescador DS. 2012. Vulnerability to global change in Mediterranean high mountains plants - Vulnerabilidad al cambio global en la alta montaña mediterránea. *Ecosistemas*, 21: 63-72.
- Escudero A, Gimenez-Benavides L, Iriondo JM, Rubio A. 2004. Patch dynamics and islands of fertility in a high mountain Mediterranean community. *Arctic Antarctic and Alpine Research*, 36: 518-527.
- Garcia-Camacho R, Albert MJ, Escudero A. 2012. Small-scale demographic compensation in a high-mountain endemic: the low edge stands still. *Plant Ecology & Diversity*, 25: 37-44.
- Garcia-Camacho R, Escudero A. 2009. Reproduction of an early-flowering Mediterranean mountain narrow endemic (*Armeria caespitosa*) in a contracting mountain island. *Plant Biology*, 11: 515-524.
- Garcia-Camacho R, Iriondo JM, Escudero A. 2010. Seedling dynamics at elevation limits: complex interactions beyond seed and microsite limitations. *American Journal of Botany*, 97: 1791-1797.
- García-Fernández A, Escudero A, Lara-Romero C, Iriondo JM. 2014. Snow cover duration effects on early life stages of *Silene ciliata*, a locally adapted mountain plant. *Plant Biology*, In press.
- García-Fernández A, Iriondo JM, Bartels D, Escudero A. 2013. Response to artificial drying until drought-induced death in different elevation populations of a high-mountain plant. *Plant Biology*, 15: 93-100.
- Garcia-Fernandez A, Iriondo JM, Escudero A. 2012a. Inbreeding at the edge: does inbreeding depression increase under more stressful conditions? *Oikos*, 121: 1435-1445.
- García-Fernández A, Segarra-Moragues JG, Widmer A, Escudero A, Iriondo JM. 2012b. Unravelling genetics at the top: mountain islands or isolated belts? *Annals of botany*, 110: 1221-1232.
- Garcia-Romero A, Munoz J, Andres N, Palacios D. 2010. Relationship between climate change and vegetation distribution in the Mediterranean mountains: Manzanares Head Valley, Sierra de Guadarrama (Central Spain). *Climatic Change*, 100: 645-666.

- Ghalambor CK, McKay JK, Carroll SP, Reznick DN. 2007. Adaptive versus non adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Functional ecology*, 21: 394-407.
- Gienapp P, Teplitsky J, Alho S, Mills A, Merila J. 2008. Climate change and evolution: disentangling environmental and genetic responses. *Molecular Ecology*, 17: 167-178.
- Gimenez-Benavides L, Albert MJ, Iriondo JM, Escudero A. 2011a. Demographic processes of upward range contraction in a long-lived Mediterranean high mountain plant. *Ecography*, 34: 85-93.
- Gimenez-Benavides L, Escudero A, Iriondo JM. 2007a. Local adaptation enhances seedling recruitment along an altitudinal gradient in a high mountain Mediterranean plant. *Annals of Botany*, 99: 723-734.
- Gimenez-Benavides L, Escudero A, Iriondo JM. 2007b. Reproductive limits of a late-flowering high-mountain Mediterranean plant along an elevational climate gradient. *New Phytologist*, 173: 367-382.
- Gimenez-Benavides L, Escudero A, Perez-Garcia F. 2005. Seed germination of high mountain Mediterranean species: altitudinal, interpopulation and interannual variability. *Ecological Research*, 20: 433-444.
- Gimenez-Benavides L, Garcia-Camacho R, Iriondo JM, Escudero A. 2011b. Selection on flowering time in Mediterranean high-mountain plants under global warming. *Evolutionary Ecology*, 25: 777-794.
- Gonzalo-Jiménez J. 2011. *Diagnosis fitoclimática de la España peninsular*. Madrid: Organismo Autónomo de Parques Nacionales.
- Gottfried M, Pauli H, Futschik A, Akhalkatsi M, Barancok P, et al., 2012. Continent-wide response of mountain vegetation to climate change. *Nature Climate Change*, 2: 111-115.
- Hamrick JL, Trapnell DW. 2011. Using population genetic analyses to understand seed dispersal patterns. *Acta Oecologica*, 37: 641-649.
- Hegland SJ, Nielsen A, Lázaro A, Bjerknes A-L, Totland Ö. 2009. How does climate warming affect plant-animal-pollinator interactions? *Ecology Letters*, 12: 184-195.
- Higgins SI, Clark JS, Nathan R, Hovestadt T, Schurr F, Fragoso JMV, Aguiar MR, Ribbens E, Lavorel S. 2003. Forecasting plant migration rates: managing uncertainty for risk assessment. *Journal of Ecology*, 91: 341-347.
- IPCC, Intergovernmental Panel on Climate Change. 2007. *Climate change 2007. The physical science basis*. Cambridge: Cambridge University Press.
- Jordano P. 1987. Patterns of mutualistic interactions in pollination and seed dispersal: connectance, dependence asymmetries, and coevolution. *American naturalist*: 657-677.
- Jordano P, Bascompte J, Olesen JM. 2003. Invariant properties in coevolutionary networks of plant-animal interactions. *Ecology letters*, 6: 69-81.
- Jump AS, Peñuelas J. 2005. Running to stand still: adaptation and the response of plants to rapid climate change. *Ecology Letters*, 8: 1010-1020.

- Jurasinski G, Kreyling Jr. 2007. Upward shift of alpine plants increases floristic similarity of mountain summits. *Journal of Vegetation Science*, 18: 711-718.
- Korner C. 2003. *Alpine Plant Life: Functional Plant Ecology of High Mountain Ecosystems; with 47 Tables*: Springer.
- Korner C. 2007. The use of 'altitude' in ecological research. *Trends in Ecology & Evolution*, 22: 569-574.
- Legendre P, Dale MRT, Fortin MJ, Gurevitch J, Hohn M, Myers D. 2002. The consequences of spatial structure for the design and analysis of ecological field surveys. *Ecography*, 25: 601-615.
- Legendre P, Fortin M. 1999. Spatial pattern and ecological analysis. *Vegetatio*, 80: 107-138.
- Linhart YB, Grant MC. 1996. Evolutionary significance of local genetic differentiation in plants. *Annual Review of Ecology Evolution and Systematics*, 27: 237-277.
- Loiselle BA, Sork VL, Nason J, Graham C. 1995. Spatial genetic structure of a tropical understory shrub, *Psychotria officinalis* (Rubiaceae). *American Journal of Botany*: 1420-1425.
- Loosmore NB, Ford ED. 2006. Statistical inference using the *G* or *K* point pattern spatial statistics. *Ecology*, 87: 1925-1931.
- Lloret F, Escudero A, Iriondo JM, Martínez-Vilalta J, Valladares F. 2012. Extreme Climatic Events and Vegetation: The Role of Stabilizing Processes. *Global Change Biology*, 18: 797-805.
- Marris E. 2008. The escalator effect. *Nature Reports Climate Change*: 94-96.
- MEA. 2005. Ecosystems and human well-being: biodiversity synthesis. *World Resource Institute*. Whasington, DC.
- Memmott J, Craze PG, Waser NM, Price MV. 2007. Global warming and the disruption of plant-pollinator interactions. *Ecology letters*, 10: 710-717.
- Miner BG, Sultan SE, Morgan SG, Padilla DK, Relyea RA. 2005. Ecological consequences of phenotypic plasticity. *Trends in Ecology & Evolution*, 20: 687-692.
- Nagy L, Grabberr G. 2009. *The biology of alpine habitats*: Oxford University Press.
- Nathan R, Klein E, Robledo-Arnuncio JJ, Revilla E. 2012. Dispersal kernels: review. In: Clobert J, Baguette M, Benton TG, Bullock JM, eds. *Dispersal ecology and evolution*. Oxford: Oxford Univesity Press.
- Nathan R, Muller-Landau HC. 2000. Spatial patterns of seed dispersal, their determinants and consequences for recruitment. *Trends in Ecology & Evolution*, 15: 278-285.
- Nieto-Feliner GN, Izuzquiza A, Lansac AR. 1996. Natural and experimental hybridization in Armeria (Plumbaginaceae): *Armeria villosa* subsp *carratracensis*. *Plant Systematics and Evolution*, 201: 163-177.
- Nogues-Bravo D, Araujo MB, Errea MP, Martinez-Rica JP. 2007. Exposure of global mountain systems to climate warming during the 21st Century. *Global Environmental Change - Human and Policy Dimensions*, 17: 420-428.
- Nogues-Bravo D, Araujo MB, Lasanta T, Moreno JIL. 2008. Climate change in Mediterranean mountains during the 21st century. *AMBIO: A Journal of the Human Environment*, 37: 280-285.

- Parmesan C. 2006. Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology Evolution and Systematics*.
- Pauli H, Gottfried M, Dullinger S, Abdaladze O, Akhalkatsi M, et al., 2012. Recent plant diversity changes on Europe's mountain summits. *Science*, 336: 353-355.
- Pauli H, Gottfried M, Reiter K, Klettner C, Grabherr G. 2007. Signals of range expansions and contractions of vascular plants in the high Alps: observations (1994-2004) at the GLORIA master site Schrankogel, Tyrol, Austria. *Global change biology*, 13: 147-156.
- Peñuelas J, Boada M. 2003. A global change-induced biome shift in the Montseny mountains (NE Spain). *Global Change Biology*, 9: 131-140.
- Reeve HK, Sherman PW. 1993. Adaptation and the goals of evolutionary research. *Quarterly Review of Biology*, 68: 1-32.
- Regato P, Salman R. 2008. *Mediterranean Mountains in a Changing World: Guidelines for developing action plans*: IUCN.
- Reidsma P, Tekelenburg T, Van den Berg M, Alkemade R. 2006. Impacts of land-use change on biodiversity: an assessment of agricultural biodiversity in the European Union. *Agriculture, ecosystems & environment*, 114: 86-102.
- Ribbens E, Silander JA, Pacala SW. 1994. Seedling recruitment in forests: calibrating models to predict patterns of tree seedling dispersion. *Ecology*, 75: 1794-1806.
- Ripley BD. 1976. The second-order analysis of stationary point processes. *Journal of applied probability*: 255-266.
- Rivas-Martínez S. 1963. *Estudio de la vegetación y flora de las Sierras de Guadarrama y Gredos*. Madrid: Instituto Botánico AJ Cavanilles.
- Rivas-Martínez S, Díaz TE, Fernández-González, Sánchez-Mata D, Pizarro J. 1990. Vegetación de la Sierra de Guadarrama. *Itinera Geobotanica*, 4.
- Sala OE, Chapin FS, Armesto JJ, Berlow E, Bloomfield J, Dirzo R, Huber-Sanwald E, Huenneke LF, Jackson RB, Kinzig A. 2000. Global biodiversity scenarios for the year 2100. *science*, 287: 1770-1774.
- Schlücht CD. 1986. The evolution of phenotypic plasticity in plants. *Annual review of ecology and systematics*: 667-693.
- Slatkin M. 1987. Gene flow and the geographic structure of natural. *Science*, 236: 787-792.
- Smouse PE, Peakall R. 1999. Spatial autocorrelation analysis of individual multiallele and multilocus genetic structure. *Heredity*, 82: 561-573.
- Stoyan D, Stoyan H. 1994. *Fractals, random shapes, and point fields: methods of geometrical statistics*. Chichester: Wiley.
- Svenning J-C, Gravel D, Holt RD, Schurr FM, Thuiller W, Münkemüller T, Schiffers KH, Dullinger S, Edwards TC, Hickler T. 2014. The influence of interspecific interactions on species range expansion rates. *Ecography*.
- Thomas CD. 2010. Climate, climate change and range boundaries. *Diversity and Distributions*, 16: 488-495.
- Thomas CD, Cameron A, Green RE, Bakkenes M, Beaumont LJ, Collingham YC, Erasmus BFN, De Siqueira MF, Grainger A, Hannah L. 2004. Extinction risk from climate change. *Nature*, 427: 145-148.


- Thomas CD, Franco A, Hill JK. 2006. Range retractions and extinction in the face of climate warming. *Trends in Ecology & Evolution*, 21: 415-416.
- Thuiller W. 2004. Patterns and uncertainties of species' range shifts under climate change. *Global Change Biology*, 10: 2020-2027.
- Thuiller W, Albert C, Araujo MB, Berry PM, Cabeza M, Guisan A, Hickler T, Midgely GF, Paterson J, Schurr FM, Sykes MT, Zimmermann NE. 2008. Predicting global change impacts on plant species' distributions: Future challenges. *Perspectives in Plant Ecology Evolution and Systematics*, 9: 137-152.
- Thuiller W, Lavorel S, Araujo MB, Sykes MT, Prentice IC. 2005. Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences of the United States of America*, 102: 8245-8250.
- Tilman D, Cassman KG, Matson PA, Naylor R, Polasky S. 2002. Agricultural sustainability and intensive production practices. *Nature*, 418: 671-677.
- Traveset A, Heleno R, Chamorro S, Vargas P, McMullen CK, Castro-Urgal R, Nogales M, Herrera HW, Olesen JM. 2013. Invaders of pollination networks in the Galápagos Islands: emergence of novel communities. *Proceedings of the Royal Society B: Biological Sciences*, 280: 20123040.
- Turesson G. 1922. The genotypical response of the plant species to the habitat. *Hereditas*, 3: 211-350.
- Tutin TG, Heywood VH, Burges NA, Valentine DH, Walters SM, Webb Da. 1995. *Flora Europaea*. Cambridge: Cambridge University Press.
- Tylianakis JM, Didham RK, Bascompte J, Wardle DA. 2008. Global change and species interactions in terrestrial ecosystems. *Ecology letters*, 11: 1351-1363.
- Van Tienderen PH. 1997. Generalists, specialists, and the evolution of phenotypic plasticity in sympatric populations of distinct species. *Evolution*: 1372-1380.
- Vekemans X, Hardy OJ. 2004. New insights from fine-scale spatial genetic structure analyses in plant populations. *Molecular Ecology*, 13: 921-935.
- Vitousek PM. 1992. Global environmental change: an introduction. *Annual Review of Ecology and Systematics*: 1-14.
- Wiegand K, Fanglinag H, Hubbel S. 2012. A systematic comparison of summary characteristic for quantifying point pattern in ecology. *Ecography*, 35: 1-12.
- Wookey PA, Aerts R, Bardgett RD, Baptist F, Brathen KA, Cornelissen JHC, Gough L, Hartley IP, Hopkins DW, Lavorel S. 2009. Ecosystem feedbacks and cascade processes: understanding their role in the responses of Arctic and alpine ecosystems to environmental change. *Global Change Biology*, 15: 1153-1172.

CAPÍTULOS

CHAPTERS

CAPÍTULO 1/CHAPTER 1

Assessing intraspecific variation in effective dispersal along an altitudinal gradient: a test in two Mediterranean high-mountain plants



Carlos Lara-Romero, Juan J. Robledo-Arnuncio, Alfredo García-Fernández;
Jose M. Iriondo

Manuscrito publicado en PLoS ONE 9(1): e87189

ABSTRACT

Background: Plant recruitment depends among other factors on environmental conditions and their variation at different spatial scales. Characterizing dispersal in contrasting environments may thus be necessary to understand natural intraspecific variation in the processes underlying recruitment. *Silene ciliata* and *Armeria caespitosa* are two representative species of cryophilic pastures above the tree line in Mediterranean high mountains. No explicit estimations of dispersal kernels have been made so far for these or other high-mountain plants. Such data could help to predict their dispersal and recruitment patterns in a context of changing environments under ongoing global warming.

Methods: We used an inverse modelling approach to analyse effective seed dispersal patterns in five populations of both *Silene ciliata* and *Armeria caespitosa* along an altitudinal gradient in Sierra de Guadarrama (Madrid, Spain). We considered four commonly employed two-dimensional seedling dispersal kernels: exponential-power, 2Dt, WALD and log-normal.

Key results: No single kernel function provided the best fit across all populations, although estimated mean dispersal distances were short (< 1 m) in all cases. *S. ciliata* did not exhibit significant among-population variation in mean dispersal distance, whereas significant differences in mean dispersal distance were found in *A. caespitosa*. Both *S. ciliata* and *A. caespitosa* exhibited among-population variation in the fecundity parameter and lacked significant variation in kernel shape.

Conclusions: This study illustrates the complexity of intraspecific variation in the processes underlying recruitment, showing that effective dispersal kernels can remain relatively invariant across populations within particular species, even if there are strong variations in demographic structure and/or physical environment among populations, while the invariant dispersal assumption may not hold for other species in the same environment. Our results call for a case-by-case analysis in a wider range of plant taxa and environments to assess the prevalence and magnitude of intraspecific dispersal variation.

INTRODUCTION

Dispersal has important implications for plant species. It determines gene flow rates and the spatial distribution of genetic diversity (Govindaraju, 1988; Ouborg *et al.*, 1999; Moran and Clark, 2012), and it influences processes like colonization (Escribano-Avila *et al.*, 2012), range expansion (Clark *et al.*, 1998), local adaptation (Blanquart *et al.*, 2012) and recruitment patterns (Nathan and Muller-Landau, 2000). It also affects metapopulation dynamics and species coexistence and diversity (Hanski, 2001; Levine and Murrell, 2003).

Dispersal studies distinguish between primary and effective dispersal. Primary dispersal is the initial movement of seeds to the location where they are first deposited, whereas effective dispersal is the complex result of primary dispersal and the post-dispersal processes that take place after the seed is deposited on the soil surface until it is a successfully established seedling (Cousens *et al.*, 2008; Nathan and Muller-Landau, 2000). Effective dispersal is more relevant to population dynamics, because it comprises the environmental factors that are needed for seedling recruitment (Cousens *et al.*, 2008; Nathan and Muller-Landau, 2000; Nathan *et al.*, 2003).

The fecundity of reproductive adults and the distribution of dispersal distances are fundamental characteristics of the dispersal process which have been extensively used as basic descriptors of this process (Clark *et al.*, 1999b; Nathan and Muller-Landau, 2000; Ribbens *et al.*, 1994; Schurr *et al.*, 2008). However, in many systems, measuring seed production and dispersal distances poses a challenge, because dispersing seeds cannot be tracked easily, and seed or seedling shadows of neighboring plants typically overlap, making it difficult to identify mother plants (Bullock *et al.*, 2006). Inverse modelling is a useful methodology that allows the fecundity of adult plants and the shape of the dispersal kernel (the probability density function of propagule dispersal distances from an individual plant) to be estimated without identifying the exact source of each seed or seedling (Clark, 1998;

Clark *et al.*, 1999b; Martinez and Gonzalez-Taboada, 2009; Ribbens *et al.*, 1994; Uriarte *et al.*, 2005); reviewed in (Nathan *et al.*, 2012).

Most seed dispersal research, employing inverse modelling or other methods, has been carried out in single populations, disregarding potentially relevant environmental variation across space that could affect the natural processes underlying the recruitment of particular species (Cousens *et al.*, 2008; Nathan *et al.*, 2012); but see (Clark *et al.*, 1998; LePage *et al.*, 2000). Heterogeneity in environmental conditions and landscape properties (*e.g.* intra and interspecific plant density, fragmentation, soil moisture, wind conditions) may indeed influence the dispersal kernel even over local geographic scales (Cousens *et al.*, 2008; Martinez and Gonzalez-Taboada, 2009; Nathan *et al.*, 2002; Nathan *et al.*, 2012). However, this variation is implicitly ignored in spatially unreplicated studies, which may lead to local results that are unrepresentative of the average dispersal pattern of the species. Hence, more comparative empirical studies are needed to test whether effective dispersal kernels exhibit intraspecific variation across contrasting environments and to find environmental correlates of potential variation (Cousens *et al.*, 2008; Klein and Oddou-Muratorio, 2011; Nathan *et al.*, 2012). Such studies could assess the validity of kernel-based approaches that assume a sole kernel for each species (Klein and Oddou-Muratorio, 2011).

High-mountain habitats have been identified as one of the most fragile environments in the world, and global warming is thought to be especially critical for plant populations in mountain systems (Nogues-Bravo *et al.*, 2007; Thuiller *et al.*, 2005). The movement of plants to higher elevations tracking their climatic niche seems to be one of the main responses to ongoing global warming (Gottfried *et al.*, 2012; Pauli *et al.*, 2012), although phenotypic plasticity and adaptation may also play an important role in their response (Davis *et al.*, 2005; Jump and Peñuelas, 2005; Miner *et al.*, 2005). Although all these processes are significantly conditioned by dispersal and recruitment patterns (Clark *et al.*, 1998; Cousens *et al.*, 2008; Davis *et al.*, 2005; Escribano-Avila *et al.*, 2012; Blanquart *et al.*, 2012), no explicit

measurements of dispersal kernels have been made for high-mountain plants so far. A better understanding of dispersal and recruitment patterns of high-mountain plants, through the study of effective dispersal kernels, would, therefore, provide greater insight into high-mountain plant species response to climate warming.

We used an inverse modelling approach to measure the net reproductive rate and effective dispersal kernel parameters along an altitudinal gradient in a threatened Mediterranean high-mountain pasture community in central Spain. Analyses were performed on five populations of each of two representative species of the community: *Silene ciliata* Poiret and *Armeria caespitosa* (Gómez Ortega) Boiss. *in DC*. Previous studies carried out on these species showed that the altitudinal gradient is associated with an environmental stress gradient, with the lowest population experiencing the most stressful conditions, constraining seedling establishment and reproductive performance (Garcia-Camacho *et al.*, 2010; Garcia-Camacho and Escudero, 2009; Gimenez-Benavides *et al.*, 2007b; Gimenez-Benavides *et al.*, 2007c; Gimenez-Benavides *et al.*, 2008). We expected that variation in population structure and physical environment along the altitudinal gradient could modify the spatial recruitment pattern and dispersal distance of the species. Specifically, we addressed two main questions: (1) What is the spatial range of effective seed dispersal in the two high-mountain species? (2) Do effective seed dispersal and fecundity parameters vary among populations at different altitudes?

METHODS

Ethics Statement

All necessary permits were obtained for the described field studies. Mr. Juan Antonio Vielva from the Administration Bureau of the Natural Park of Cumbres, Circo y Lagunas de Peñalara and Mr. Antonio Sanz from the Administration Bureau of the Regional Park of Cuenca Alta del Manzanares gave their permission to work in the protected natural areas. Field studies did not involve any endangered or protected species.

Study site and species

The study was carried out in the orophyllous cryophilic pastures of Sierra de Guadarrama, a mountain range located in central Spain. Mean annual precipitation, measured at the Navacerrada Pass ($40^{\circ} 46'N$, $4^{\circ} 19'W$; 1860 m a.s.l.), is 1330 mm with a pronounced dry season (< 10% of total annual rainfall) from May to October. Mean annual temperature is $6.3^{\circ}C$, with mean monthly temperatures ranging from $-1^{\circ}C$ in January to $16^{\circ}C$ in July (www.aemet.es). Dry cryophilic pastures occur in the higher summits above the tree line between 1900 and 2430 m.a.s.l and are dominated by *Festuca curvifolia* Lag. ex Lange and other perennial plants interspersed in a shrub matrix characterized by *Cytisus oromediterraneus* Rivas Mart. & al. and *Juniperus communis* subsp. *alpina* (Suter) Čelak.

Silene ciliata Poiret (Caryophyllaceae) is a chamaephytic cushion perennial plant that occurs in the Mediterranean mountain ranges of southern Europe (Tutin *et al.*, 1995). One of its southernmost distribution limits is found in Sierra de Guadarrama at altitudes from 1900 to 2430 m, where it grows in dry cryophilic pastures dominated by *Festuca curvifolia*. It blooms in late summer, with a peak in early August (Gimenez-Benavides *et al.*, 2011b). Flowering stems reach 15 cm in height and have 1–5 flowers. Fruit capsules have up to 100 seeds which are wind dispersed in August-September. *Silene ciliata* seeds have an average mass of 0.59 mg and their diameter range between 1.1 and 1.5 mm (Gimenez-Benavides *et al.*, 2005). The species is essentially barochorous (seeds lack any specific structure to promote dispersal) and is pollinated by syrphid flies, bumblebees and moth species (Gimenez-Benavides *et al.*, 2007a). It is self-compatible, although autogamy is restricted by pronounced protandry (Garcia-Fernandez *et al.*, 2012a). In Sierra de Guadarrama, genetic diversity is quite homogeneous across *S. ciliata* populations, which show significant levels of inbreeding (Garcia-Fernandez *et al.*, 2012a).

Armeria caespitosa (Gómez Ortega) Boiss. in DC. is a high-mountain dwarf chamaephytic cushion plant, endemic to the mountains of central Spain (Sierra de Guadarrama, Ayllón and East Gredos), which occurs at altitudes from 1600 to 2430 m. It grows in the same dry cryophilic pastures dominated by *Festuca curvifolia*,

although isolated individuals have also been found on the ledges of granite and gneiss rocks. Its flowers are grouped in short-scaped flowerheads with 19 ± 8 flowers each. Each flower has a single ovule, which yields one seed that remains enclosed in the papyraceous calyx. Average seed mass is 1.18 mg and its diameter ranges between 2.0 and 4.7 mm (Garcia-Camacho *et al.*, 2010). The seeds have a papyraceous calyx that may facilitate wind dispersal (Garcia-Camacho *et al.*, 2010). *A. caespitosa* is self-incompatible (Garcia-Camacho and Escudero, 2009) and pollinated by bees, bumblebees and syrphid flies (Garcia-Camacho and Escudero, 2009). Previous studies on the population genetics of both species revealed relatively low genetic differentiation and a complex genetic structure among populations (García-Fernández *et al.*, 2013; Garcia-Fernandez *et al.*, 2012b).

Field data

In August and September 2010, we established 10x10m plots in five populations of both *S. ciliata* and *A. caespitosa* distributed along an altitudinal gradient in the study site (Table 1). For each plot, we estimated a set of climate variables for the growing season (April-September). We calculated the mean, minimum and maximum rainfall and temperature for each population using the Digital Climatic Atlas of Spain of the Spanish Ministry of Environment (Gonzalo-Jiménez, 2011) (Table 1), and the mean wind speed at 80 m height using the Spanish Wind Energy Atlas of the Spanish Ministry of Industry (<http://atlaseolico.idae.es>) (Table 1).

Table 1. Description of *A. caespitosa* and *S. ciliata* sampling sites.

Population	UTM Coordinates (x,y)		Altitude m	Pm (min, max) mm	Tm (min, max) °C	Ws m/s	Seedling	Reproductive
<i>Silene ciliata</i>								
Peñalara	419427.30	4522814.15	2405	82.50 (32, 149)	8.77 (0.2, 14.1)	10.62	469	1068
Cabezas de Hierro	421169.50	4516859.61	2305	85.17 (34, 151)	8.37 (0.2, 13.7)	10.87	243	1085
Nevero	428863.74	4537167.57	2190	71.00 (30, 121)	9.67 (2.1, 14.6)	8.97	19	245
Najarra	430196.59	4518886.29	2080	72.17 (27, 112)	10.45 (2.8, 15.7)	9	89	523
Laguna	419931.16	4521082.09	1946	77.00 (31, 134)	10.27 (2.8, 15.2)	7.17	6	147
<i>Armeria caespitosa</i>								
Cabezas de Hierro	420950.54	4516685.32	2336	85.2 (34, 151)	8.37 (0.2, 13.7)	10.87	145	318
Najarra	430196.79	4518886.33	2080	72.2 (27, 112)	10.45 (2.8, 15.7)	9	498	356
Loma de Cabezas	420013.12	4514794.49	1970	77.0 (30, 131)	9.78 (2.5, 14.8)	8.45	53	132
Collado de las Vacas	419168.95	4513371.38	1882	66.3 (25, 110)	11.80 (4.0, 17.2)	8.46	49	169
Sierra de los Porrones	420580.85	4512251.64	1647	64.5 (26, 102)	12.88 (4.4, 18.3)	7.4	32	52

Geographical coordinates, altitude, climatic variables for the growing season (April-September) and seedling and reproductive plant density (individuals per 10x10 m plot) in study populations. Pm mean monthly precipitation, Tm mean monthly temperature, Ws wind speed at 80 m height. Minimum and maximum monthly precipitation and temperatures are provided between brackets.

In each study plot, we mapped every *S. ciliata* and *A. caespitosa* reproductive plant and seedling using two high-resolution Differential Global Navigation Satellite System (DGNSS) receivers (Viva GS15, Leica, Switzerland) with an accuracy of 5 cm for *x* and *y* coordinates. We also measured plant diameter and the number of inflorescences of every reproductive plant. To account for potential seedling sources outside the study area, we also recorded the location, diameter and number of inflorescences of reproductive plants in a buffer zone of two meters around the plots. The seedling stage in *S. ciliata* and *A. caespitosa* includes up to one-year individuals (*i.e.* plants that had germinated in fall or spring and had survived their first full summer), because the seedlings that survive a second growth period grow into the reproductive stage (Garcia-Camacho *et al.*, 2012; Gimenez-Benavides *et al.*, 2011a). Summer is critical to the survival of Mediterranean plants due to the incidence of severe droughts (Beniston *et al.*, 2007; Larcher, 2000). This is also the case in Mediterranean high-mountain plants (Castro *et al.*, 2005) such as *S. ciliata* and *A. caespitosa* (Garcia-Camacho *et al.*, 2010; Gimenez-Benavides *et al.*, 2007c). With this framework, we included all critical stages of effective dispersal: seed dispersal, seed germination and seedling survival.

The dataset used for this study is available from the institutional repository of Rey Juan Carlos University (<http://ecienca.urjc.es/handle/10115/11835>).

Statistical analysis

We used inverse modelling to estimate seedling dispersal kernel parameters for each population of each species and to test for dispersal parameter variation among populations within species. Following Ribbens *et al.*, (1994), the expected number of seedling recruits \hat{s}_j at a given plot or trap *j* of area A_j (\hat{s}_j) equals the sum of the seedling shadows across this plot of all *m* maternal plants, in the form:

$$\hat{s}_j(\mathbf{b}, \mathbf{r}_j; \beta, \delta, u) = \sum_{i=1}^m Q(b_i; \beta) \int_{A_j} f\left(\sqrt{(x_i - x_j)^2 + (y_i - y_j)^2}; \delta, u\right) dx_j dy_j \quad \text{eqn. 1a}$$

where \mathbf{b} and \mathbf{r}_j are, respectively, m -length vectors of individual plant size measurements b_i and spatial distances r_{ij} between the spatial coordinates (x_j, y_j) of plant i , and the coordinates of plot j (x_i, y_i) , Q is an allometric function with parameter β yielding the number of seedlings produced by a single plant, and f is a two-dimensional isotropic seedling dispersal probability density function (dispersal kernel) as a function of dispersal distance $r_{ij} = \sqrt{(x_i - x_j)^2 + (y_i - y_j)^2}$ with mean δ and shape parameter u . The integral in *eqn. 1a* provides the exact probability of dispersal from plant i into the whole surface A_j of plot j . In practice, all studies of which we are aware have assumed that the probability of dispersal from any given plant into any of the points lying within the area A_j of each plot is constant, yielding the usual simplified expression:

$$\hat{s}_j(\mathbf{b}, \mathbf{r}_j; \beta, \delta, u) = \sum_{i=1}^m Q(b_i; \beta) f(r_{ij}; \delta, u) A_j, \quad \text{eqn 1b}$$

where r_{ij} is the spatial distance between plant i and the centre of plot j . The approximation in *eqn 1b* will be closer to the exact values predicted by *eqn. 1a* the smaller the seedling plot areas and the larger the distances between maternal plants and seedling plots.

We assumed a linear relationship between seedling production and plant diameter, $Q_i = \beta b_i$, because previous studies showed that plant size is linearly related with seed production in both species (Garcia-Camacho and Escudero, 2009; Gimenez-Benavides *et al.*, 2007c). Furthermore, as found in previous studies (Clark *et al.*, 1999a; Schurr *et al.*, 2008), alternative allometric functions worsened inverse modelling fits, including those using the number of inflorescences (results not shown).

For every population of each species, we divided the 100-m² study area in $c = 100L^2$ equal adjacent cells of area L^2 , and for each cell j , we computed the observed (s_j) and predicted (\hat{s}_j) seedling densities. We used both equations 1a and

1b to estimate \hat{s}_j , given that in our case, with relatively large cell areas compared to some of the mother-cell pairwise distances (occasionally with maternal plants lying within the target cell), it was not obvious *a priori* that the approximation in equation 1b would be sufficiently good. We obtained however virtually identical dispersal and fecundity estimates for all populations and assumed kernels with both *equation 1a* or *1b* (results not shown), so we only present the results for the latter, whose computational efficiency enabled calculating confidence intervals and conducting hypothesis tests (see below) in a reasonable number of CPU hours. We also tested the effect on parameter estimates of three different values for cell side length, $L = 0.125, 0.25$ and 0.50 m, with lower and upper values approximating DGNSS accuracy and dispersal range, respectively. Distribution of seedling densities across cells exhibited overdispersion relative to a Poisson distribution ($p < 0.05$) in all populations excepting *S. ciliata*'s Nevero, and we thus modelled seedling counts using a negative binomial distribution (Clark *et al.*, 1998):

$$L(\mathbf{S} | \beta, \delta, u, \theta) = \prod_{j=1}^c \frac{\Gamma(s_j + \theta)}{\Gamma(s_j + 1)\Gamma(\theta)} \frac{\hat{s}_j^{s_j} (\mathbf{b}, \mathbf{r}_j; \beta, \delta, u) \theta^\theta}{[\hat{s}_j (\mathbf{b}, \mathbf{r}_j; \beta, \delta, u)]^{s_j + \theta}}, \quad eqn\ 2$$

where \mathbf{S} is the set of c observed seedling counts s_j , Γ is the gamma function, and θ is the negative binomial parameter, determining the distribution variance (values of $\theta < 1$ indicate overdispersed distributions, while $\theta \gg 1$ tend to the Poisson distribution). Maximum-likelihood estimates for β , δ , u and θ were obtained for each population of each species by maximizing *eqn 2*. Confidence intervals were computed using the profile-likelihood method (Edwards, 1992).

As most studies show that no single dispersal function provides consistently superior fits across species (Greene *et al.*, 2004; Nathan *et al.*, 2012, Schurr *et al.*, 2008; Uriarte *et al.*, 2005), it is advisable to always compare the performance of alternative dispersal kernels (Martinez and Gonzalez-Taboada, 2009; Schurr *et al.*, 2008). We considered four commonly employed two-dimensional seedling dispersal

kernels, with scale parameter a and shape parameter u : the 2Dt kernel (Clark *et al.*, 1999b); see (Austerlitz *et al.*, 2004) for the parameterization used here.

$$f_{2\text{Dt}}(r_{ij} | a, u) = \frac{u-1}{\pi a^2} \left(1 + \frac{r_{ij}^2}{a^2} \right)^{-u}, \quad \text{eqn 3}$$

the log-normal kernel (Stoyan and Wagner, 2001)

$$f_{LN}(r_{ij} | a, u) = \frac{1}{(2\pi)^{3/2} u r_{ij}^2} \exp \left[-\frac{(\ln(r_{ij}) - a)^2}{2u^2} \right], \quad \text{eqn 4}$$

the WALD kernel (Katul *et al.*, 2005, Schurr *et al.*, 2008)

$$f_{WALD}(r_{ij} | a, u) = \left(\frac{u}{8\pi^3 r_{ij}^5} \right)^{1/2} \exp \left[-\frac{u(r_{ij} - a)^2}{2a^2 r_{ij}} \right], \quad \text{eqn 5}$$

and the exponential-power kernel (Clark *et al.*, 1998)

$$f_{EXPOW}(r_{ij} | a, u) = \frac{u}{2\pi a^2 \Gamma(2/u)} \exp \left[-\left(\frac{r_{ij}}{a} \right)^u \right]. \quad \text{eqn 6}$$

Mean seedling dispersal distances for each kernel are given by $\delta_{2\text{Dt}} = a\Gamma(3/2)\Gamma(u-3/2)/\Gamma(u-1)$, $\delta_{LN} = \exp(a + 0.5u^2)$, $\delta_{WALD} = a$ and $\delta_{EXPOW} = a\Gamma(3/u)/\Gamma(2/u)$, respectively. Given that δ provides a more intuitive description of dispersal range than a , independently of the assumed kernel, we parameterized the model (*eqns 1* and *2*) in terms of δ and u , rather than in terms of a and u , facilitating estimation of profile-likelihood confidence intervals for δ (Clark *et al.*, 1998; Oddou-Muratorio *et al.*, 2005) and tests of its variation across populations (see below). We used the Akaike Information criterion (AIC) (Akaike, 1973) to assess method performance and calculated the goodness of fit of the most parsimonious model by measuring Pearson's product-moment correlation between observed vs. predicted seedling densities.

We tested for potential differences in dispersal and fecundity parameters across populations using likelihood ratio statistics as in Clark *et al.* (Clark *et al.*, 1998). Assuming that mean dispersal distance δ is constant across populations within species, we obtained parameter estimates based on the likelihood of the whole species data set \mathbf{S}_q , incorporating information from all q populations of the species:

$$L_\delta(\mathbf{S}_q | \boldsymbol{\beta}_q, \delta, \mathbf{u}_q, \boldsymbol{\theta}_q) = \prod_{k=1}^q L(\mathbf{S}_k | \beta_k, \delta, u_k, \theta_k), \quad eqn\ 7$$

where $\boldsymbol{\beta}_q$, \mathbf{u}_q , and $\boldsymbol{\theta}_q$ are q -length vectors of population-specific fecundity parameters β_k , kernel shape parameters u_k , and negative binomial parameters θ_k . This model has $3q$ degrees of freedom and provides an average species-level estimate of δ . Note that, unlike Clark *et al.* (1998), we allowed for different values of the negative binomial parameter (*i.e.* different degrees of spatial seedling clumping) across populations. We assumed the same kernel family for all populations for conducting this test, choosing the one yielding the best average fit over all populations (*i.e.* the one with the smallest average AIC across the population-level fits obtained using *eqn. 2*). We then obtained parameter estimates assuming that all parameters (including δ) are variable among populations within species (still assuming a single consensual kernel family), based on the likelihood

$$L(\mathbf{S}_q | \boldsymbol{\beta}_q, \boldsymbol{\delta}_q, \mathbf{u}_q, \boldsymbol{\theta}_q) = \prod_{k=1}^q L(\mathbf{S}_k | b_k, \delta_k, u_k, \theta_k), \quad eqn\ 8$$

where $\boldsymbol{\delta}_q$ is the q -length vector of population-specific mean dispersal distances δ_k . The model in *eqn 8* has $4q-1$ degrees of freedom. The deviance

$$D = -2 \ln \left[\frac{L_\delta(\mathbf{S}_q | \boldsymbol{\beta}_q, \delta, \mathbf{u}_q, \boldsymbol{\theta}_q)}{L(\mathbf{S}_q | \boldsymbol{\beta}_q, \boldsymbol{\delta}_q, \mathbf{u}_q, \boldsymbol{\theta}_q)} \right] \quad eqn\ 9$$

is asymptotically distributed as χ^2 with $q-1$ degrees of freedom. Large deviances mean that stand-specific δ improves the likelihood of the data so substantially that the null hypothesis of invariant δ across populations can be rejected (Clark *et al.*, 1998). We first tested for significant differences in δ across all populations of each species. When the overall test was positive, we tested for specific populations with δ -values significantly different from the global species-average by subsequently conducting q deviance tests, one for each population i (χ^2 with 1 degree of freedom), using:

$$D_i = -2 \ln \left[\frac{L_\delta(\mathbf{S}_q | \boldsymbol{\beta}_q, \delta, \mathbf{u}_q, \boldsymbol{\theta}_q)}{L(\mathbf{S}_q | \boldsymbol{\beta}_q, \delta_i, \delta_{-i}, \mathbf{u}_q, \boldsymbol{\theta}_q)} \right], \quad eqn\ 10$$

where the numerator is the same likelihood as before, assuming that the mean dispersal distance δ is constant across all populations, while the likelihood in the denominator assumes that the i -th population has a mean dispersal distance δ_i , while the remaining $q-1$ populations have a constant mean dispersal distance δ_{-i} . We applied the sequential Bonferroni correction to correct for multiple testing (Holm, 1979). Analogous tests were conducted to investigate among-population variation in the dispersal kernel's shape parameter (assuming constant u and variable δ in *eqn 7*) and in the fecundity parameter β (assuming constant β and variable δ in *eqn 7*). Although the field survey was exhaustive in all study sites, the sample sizes were unbalanced and relatively small in some sites due to natural density variation (Table 1), which may compromise the large numbers approximation under which the χ^2 distribution is expected. It has been shown that small samples can indeed make likelihood ratio (LR) tests relatively lax in the case of unequal male reproductive success analysis (Elle and Meagher, 2000). Smouse *et al.* (1999), proposed the use of nonparametric (permutational) tests as an alternative to LR tests, showing that the former can perform better than LR tests when parent-offspring genealogical information is available. Similar non-parametric tests have not been developed so far in the different statistical framework of inverse modeling, for which LR tests, such

as those used here, are still widely employed as feasible approximations for exploratory assessment of variation in dispersal and fecundity parameters (Caspersen and Saprunoff, 2005; Clark *et al.*, 1998; Debain *et al.*, 2007; Sagnard *et al.*, 2007). All statistical analyses were performed in the open source software package R (R Core Team, 2013).

RESULTS

We mapped a total of 4095 reproductive individuals and 1603 seedlings across all populations of the two species (Table 1; Figure S1 in Supporting Information). Reproductive plant and seedling densities varied between populations in both species (Table 1). We focused on the results obtained with a 0.125 m cell side length (Tables 2 and 3), because some of the *S. ciliata* populations did not converge when 0.25 or 0.5 m cell side lengths were used (Table S1 in Supporting Information). In any case, similar results were obtained for both species for all cell side lengths (Tables 2, 3, S1 and S2). As none of the four models converged for *A. caespitosa* in the Najarra population, this population was excluded from further analyses.

Differences in AIC values indicated that no single kernel function provided the best fit across all populations of any of the two species (Table 2 and 3). For *S. ciliata*, the WALD kernel was among the most parsimonious models (*i.e.* models whose AIC values differ $\Delta\text{AIC} < 2$ from the model with the lowest AIC; Burnham & Anderson 2002) in all five populations, while the log-normal model was among the most parsimonious in four populations, the exponential-power model in three populations and the 2Dt model in two populations (Table 2). Results did not support a particular kernel shape in *S. ciliata*: the mesokurtic, thin-tailed exponential-power with $u > 1$ showed similarly good fits ($\Delta\text{AIC} < 2$) as leptokurtic functions with either exponential (WALD) or fat (2Dt and log-normal) tails, within the same populations. Using the WALD kernel across all populations of *S. ciliata*, species-level estimates of the mean dispersal distance and the shape and fecundity parameters were $\delta = 0.39$

m (95% CI: 0.35 – 0.45 m), $u = 0.80$ (0.63 – 1.13) and $\beta = 0.0032$ seedling/cm (0.0003 - 0.0061). For *A. caespitosa*, it was the log-normal kernel that ranked among the best models in all four populations where estimation converged, whereas the WALD, 2Dt and exponential-power models showed a similarly good performance within each of three populations (Table 3). Supported models in *A. caespitosa* always included leptokurtic functions with exponential or fat tails (including the exponential-power with $u < 1$). Using the log-normal model across all populations of *A. caespitosa*, the species-level estimates kernel's parameters were $\delta = 0.43$ m (0.32 – 0.60 m), $u = 0.98$ (0.87 – 1.14) and $\beta = 0.0032$ seedling/cm (0.0003 - 0.0061).

Table 2. Estimated parameters for the models fitted to the seedling recruitment data of *S. ciliata*.

Population	Kernel	Mean dispersal (δ) (m)	Shape parameter (u)	Fecundity parameter (β) (seedlings/cm)	Negative binomial parameter (ϕ)	- logL	ΔAIC	r^2
Peñalara	Log-norm	0.32 (0.27 - 0.39)	0.64 (0.52 - 0.77)	0.009 (0.008 - 0.010)	0.359 (0.260 - 0.523)	1624.80	-	0.12
	WALD	0.33 (0.28 - 0.41)	0.64 (0.44 - 0.99)	0.009 (0.008 - 0.010)	0.360 (0.260 - 0.525)	1624.82	0.04	
	2Dt	0.32 (0.27 - 0.46)	3.80 (1.99 - na)	0.009 (0.009 - 0.011)	0.351 (0.255 - 0.509)	1627.77	5.94	
	Exp-pow	0.31 (0.27 - 0.37)	1.51 (0.97 - 2.34)	0.009 (0.008 - 0.010)	0.349 (0.253 - 0.507)	1628.76	7.92	
Cabezas de Hierro	Exp-pow	0.37 (0.32 - 0.45)	9.86 (2.34 - na)	0.003 (0.003 - 0.004)	0.449 (0.235 - 1.198)	1029.04	-	0.05
	Log-norm	0.43 (0.35 - 0.60)	0.43 (0.23 - 0.70)	0.003 (0.003 - 0.004)	0.460 (0.240 - 1.246)	1029.04	0.01	
	WALD	0.43 (0.35 - 0.62)	2.26 (0.84 - 8.70)	0.003 (0.003 - 0.004)	0.459 (0.239 - 1.244)	1029.17	0.27	
	2Dt	0.43 (0.33 - 0.58)	172.54 (3.65 - na)	0.003 (0.003 - 0.004)	0.450 (0.235 - 1.205)	1031.17	4.25	
Nevero	Exp-pow	0.23 (0.17 - 0.50)	2.84 (0.74 - 21.91)	0.001 (0.001 - 0.002)	142.062 (0.045 - na)	108.04	-	0.08
	2Dt	0.23 (0.17 - 0.42)	171.06 (1.74 - na)	0.001 (0.001 - 0.002)	142.162 (0.082 - na)	108.12	0.17	
	WALD	0.30 (0.22 - 0.60)	1.23 (0.41 - 3.27)	0.001 (0.001 - 0.002)	141.865 (0.054 - na)	109.04	1.99	
	Log-norm	0.30 (0.21 - 0.86)	0.47 (0.29 - 1.35)	0.001 (0.001 - 0.002)	50.393 (0.053 - na)	109.14	2.19	
Najarra	WALD	0.71 (0.37 - na)	0.55 (0.27 - 1.24)	0.002 (0.003 - 0.003)	0.280 (0.092 - 8.456)	461.85	-	0.04
	Log-norm	0.62 (0.36 - 3.01)	0.86 (0.58 - 1.46)	0.002 (0.002 - 0.003)	0.280 (0.092 - 8.401)	462.14	0.59	
	Exp-pow	0.61 (0.36 - 2.45)	0.83 (0.27 - 2.49)	0.002 (0.002 - 0.003)	0.275 (0.091 - 7.395)	463.15	2.59	
	2Dt	nc	nc	nc	nc	nc	nc	
Laguna	Exp-pow	0.29 (0.17 - 0.40)	1411.75 (0.03 - na)	0.001 (0.0003 - 0.004)	0.003 (0.001 - 0.035)	38.74	-	0.01
	WALD	0.24 (0.14 - na)	1.00 (na - na)	0.001 (0.0003 - 0.029)	0.004 (0.001 - 0.049)	38.91	0.34	
	Log-norm	0.24 (0.16 - na)	0.47 (0.23 - na)	0.001 (0.0003 - na)	0.004 (0.001 - 0.048)	38.95	0.42	
	2Dt	0.23 (0.15 - 0.86)	165.18 (1.00 - na)	0.001 (0.0003 - 0.006)	0.003 (0.001 - 0.035)	39.12	0.76	

For each parameter, 95% confidence intervals are presented along with the mean value. Log-norm, log-normal; Exp-pow, exponential-power. na denotes that the confidence interval limit is not available because of flat likelihood function. nc denotes models that did not converge. – logL, log-likelihood; AIC, Akaike's information criterion; r, Pearson correlation coefficient between observed and predicted seedling density values. Models with the lowest AIC are marked in bold for each population.

The fits of the most parsimonious models ranged from good in *A. caespitosa* populations ($r^2 = 0.17\text{--}0.35$) to fair or poor ($r^2 = 0.01\text{--}0.12$) in *S. ciliata* populations (Tables 2 and 3), and were highly significant ($p < 0.0001$) in all cases except in the population with the lowest seedling number ($p = 0.32$ for Laguna in *S. ciliata*, which had only 6 seedlings). Population estimates of mean dispersal distance (δ) and fecundity (β) parameters were generally consistent when assuming different kernel families (Table 2 and 3). No significant correlation was found between $\log \beta$ and $\log \delta$ ($r = 0.35$, $p = 0.65$).

Table 3. Estimated parameters for the models fitted to the seedling recruitment data of *A. caespitosa*.

Population	Kernel	Mean dispersal (δ) (m)	Shape parameter (u)	Fecundity parameter (β) (seedlings/cm)	Negative binomial parameter (θ)	- logL	ΔAIC	r^2
Cabezas de Hierro	Log-norm	0.33 (0.24 - 0.56)	0.97 (0.78 - 1.29)	0.024 (0.019 - 0.029)	0.122 (0.076 - 0.208)	584.41	-	0.17
	2Dt	0.80 (0.30 - na)	1.60 (1.24 - 2.48)	0.024 (0.020 - 0.030)	0.122 (0.076 - 0.207)	584.78	0.73	
	WALD	0.35 (0.24 - 1.44)	0.21 (0.13 - 0.35)	0.024 (0.020 - 0.030)	0.121 (0.076 - 0.206)	584.80	0.79	
	Exp-pow	0.29 (0.23 - 0.45)	0.60 (0.31 - 1.03)	0.024 (0.020 - 0.029)	0.120 (0.075 - 0.202)	585.01	1.19	
Loma de Cabezas	Exp-pow	0.66 (0.46 - 1.39)	0.69 (0.24 - 2.37)	0.014 (0.010 - 0.019)	0.061 (0.028 - 0.162)	247.54	-	0.12
	Log-norm	0.78 (0.48 - 2.57)	1.02 (0.69 - 1.64)	0.014 (0.010 - 0.019)	0.063 (0.028 - 0.169)	247.82	0.57	
	WALD	0.77 (0.46 - na)	0.44 (0.19 - 1.01)	0.014 (0.010 - 0.020)	0.064 (0.028 - 0.171)	247.96	0.85	
Collado de las Vacas	2Dt	0.73 (0.05 - na)	2.15 (0.12 - na)	0.014 (0.010 - 0.020)	0.058 (0.026 - 0.158)	248.64	2.20	
	WALD	0.28 (0.19 - 0.58)	0.25 (0.13 - 0.44)	0.011 (0.008 - 0.014)	0.286 (0.107 - 1.330)	206.84	-	0.22
	Log-norm	0.25 (0.17 - 0.43)	0.84 (0.63 - 1.18)	0.011 (0.008 - 0.014)	0.298 (0.109 - 1.444)	207.11	0.56	
	Exp-pow	0.27 (0.18 - 0.45)	0.56 (0.29 - 0.96)	0.011 (0.008 - 0.015)	0.249 (0.096 - 1.057)	209.86	6.05	
Sierra de los Porrones	Log-norm	0.23 (0.13 - 0.50)	1.07 (0.84 - 1.56)	0.021 (0.013 - 0.033)	0.369 (0.118 - 2.966)	110.01	-	0.35
	2Dt	0.28 (0.16 - na)	1.75 (1.27 - 3.12)	0.021 (0.013 - 0.033)	0.371 (0.117 - 3.291)	110.02	0.01	
	Exp-pow	0.21 (0.13 - 0.40)	0.48 (0.19 - 0.93)	0.020 (0.013 - 0.033)	0.370 (0.118 - 3.227)	110.15	0.27	
	WALD	0.20 (0.10 - na)	0.07 (0.03 - 0.12)	0.024 (0.015 - 0.043)	0.280 (0.098 - 1.231)	111.27	2.52	

For each parameter, 95% confidence intervals are presented along with the mean value. Log-norm, log-normal; Exp-pow, exponential-power. na denotes that the confidence interval limit is not available because of flat likelihood function. – logL, log-likelihood; AIC, Akaike's information criterion; r, Pearson correlation coefficient between observed and predicted seedling density values. Models with the lowest AIC are marked in bold for each population.

Focusing on the WALD dispersal model for *S. ciliata* cross-population comparisons, estimated mean dispersal distances ranged between $\delta = 0.23$ and 0.71 m (Table 2), although differences were only marginally significant ($\chi^2_{4\text{d.f.}} = 8.846$, $p = 0.065$). Estimated shape parameters ranged between $u = 0.55$ and 2.26, and were not significantly different from each other ($\chi^2_{4\text{d.f.}} = 7.022$, $p = 0.135$). Fecundity parameters ranged between $\beta = 0.001$ and 0.009 seedlings/cm across *S. ciliata* populations, being significantly different ($\chi^2_{4\text{d.f.}} = 254.26$, $p < 0.001$). Per-population tests indicated that all population estimates of β were significantly different from the species average (Table 4).

Table 4. Comparison of estimated fecundity parameters (β) among *S. ciliata* populations.

Population	β_i (seedlings/cm)	β_{-i} (seedlings/cm)	-logL	$\chi^2_{1\,d.f.}$ (<i>p</i> -value)
Peñalara	0.009	0.003	-3276.28	229.28 (0.0000)*
Cabezas de Hierro	0.003	0.007	-3365.25	51.35 (0.0000)*
Nevero	0.001	0.006	-3361.72	58.41 (0.0000)*
Najarra	0.002	0.006	-3366.10	49.64 (0.0000)*
Laguna	0.001	0.005	-3380.92	19.99 (0.0000)*
Average	0.005	-	-3390.92	-

β_i is the fecundity parameter for the corresponding population and β_{-i} is the average β across the remaining four populations. -logL is the log-likelihood of the model considering β_i and separately β_{-i} . Last column shows the deviance ($\chi^2_{1\,d.f.}$ -distributed) of the model considering β_i and separately β_{-i} relative to the model (shown in the last row) assuming constant β across all five populations. Asterisks denote a significant test after sequential Bonferroni correction.

Assuming a log-normal dispersal model for *A. caespitosa*, population estimates of δ varied between 0.23 and 0.78 m (Table 3) and were significantly different ($\chi^2_{3\,d.f.} = 14.022$, $p = 0.003$). Per-population tests showed that only the population with the largest δ (Loma de Cabezas with 0.78 m) differed significantly from the species average (Table 5). Estimated shape parameters ranged between $u = 0.84$ and 1.07 and were not significantly different from each other ($\chi^2_{3\,d.f.} = 1.426$, $p = 0.699$). Fecundity parameter estimates exhibited significant differences in *A. caespitosa* ($\chi^2_{3\,d.f.} = 22.51$, $p < 0.001$), ranging from $\beta = 0.011$ to 0.024 seedlings/cm. Per-population tests indicated that the maximum (at Cabezas de

Hierro) and minimum (at Collado de las Vacas) estimates of β were significantly larger and smaller than the species average, respectively (Table 6).

Table 5. Comparison of estimated mean dispersal distances (δ) among *A. caespitosa* populations.

Population	δ_i (m)	δ_i (m)	-logL	$\chi^2_{1\,d.f.}$ (<i>p-value</i>)
Cabezas de Hierro	0.33	0.49	1155.80	1.84 (0.1744)
Loma de Cabezas	0.78	0.29	1150.46	12.53 (0.0003)*
Collado de las Vacas	0.25	0.50	1154.47	4.50 (0.0338)
Sierra de los Porrones	0.23	0.47	1155.20	3.04 (0.0809)
Average	0.43	-	1156.72	-

δ_i is the mean dispersal distance for the corresponding population and δ_i is the average δ across the remaining three populations. -logL is the log-likelihood of the model considering δ_i and separately δ_i . Last column shows the deviance ($\chi^2_{1\,d.f.}$ -distributed) of the model considering δ_i and separately δ_i relative to the model (shown in the last row) assuming constant δ across all four populations. The asterisk denotes a significant test after sequential Bonferroni correction.

Table 6. Comparison of estimated fecundity parameters (β) among *A. caespitosa* populations.

Population	β_i (seedlings/cm)	β_{-i} (seedlings/cm)	-logL	$\chi^2_{1\text{d.f.}}$ (<i>p</i> -value)
Cabezas de Hierro	0.024	0.013	1152.64	16.65 (0.0000)*
Loma de Cabezas	0.014	0.020	1159.56	2.82 (0.0932)
Collado de las Vacas	0.011	0.021	1153.55	14.84 (0.0001)*
Sierra de los Porrones		0.021	1160.66	0.61(0.4337)
Average	0.017	-	1160.97	-

β_i is the fecundity parameter for the corresponding population and β_{-i} is the average β across the remaining three populations. $-\log L$ is the log-likelihood of the model considering β_i and separately β_{-i} . Last column shows the deviance ($\chi^2_{1\text{d.f.}}$ -distributed) of the model considering β_i and separately β_{-i} relative to the model (shown in the last row) assuming constant β across all four populations. Asterisks denote significant tests after sequential Bonferroni correction.

DISCUSSION

We used an inverse modelling approach to analyse the variation in effective seed dispersal patterns of two coexisting species along an altitudinal gradient in a high mountain pasture community of central Spain. Estimated mean dispersal distances were short in both species, with most seedlings established less than a meter from the source. There was not a single kernel function that provided the best fit across all populations. Furthermore, there were significant differences in the fecundity parameter among populations in both species and in the mean dispersal distance among populations of *A. caespitosa*, while differences in the mean dispersal

distance among *S. ciliata* populations were only marginally significant. The kernel's shape parameter did not vary significantly in any of the two species.

The inverse modelling approach proved to be useful in characterizing effective seed dispersal patterns in both species. Parameter estimates were consistent among populations and species independently of the assumed kernel. In addition, estimates conformed to independent fecundity measures for the same species (Garcia-Camacho *et al.*, 2010; Gimenez-Benavides *et al.*, 2008), and to primary seed dispersal estimates obtained with direct and indirect methods in congeneric species (*Silene latifolia*: 0.17-0.85 m, (Jongejans and Schippers, 1999); *Armeria maritima*: 0.6 m, (Philipp *et al.*, 1992). Although r^2 values were slightly lower than those obtained in analogous studies (LePage *et al.*, 2000; Ribbens *et al.*, 1994; Uriarte *et al.*, 2005), they were highly significant (except in the population with the lowest seedling number). This suggests that inverse modelling was able to provide information about the scale of dispersal in the high mountain plants studied, even though as frequently found in heterogeneous environments there seemed to be substantial distance-independent fluctuations in dispersal and establishment probabilities that cannot be characterized by simple kernel fits (Nathan *et al.*, 2012).

Most effectively dispersed seeds were established near conspecific adult plants (Figure S1). Consequently, estimated mean dispersal distances were low, suggesting that both *A. caespitosa* and *S. ciliata* have low effective seed dispersal ranges. Effective dispersal scale is influenced by factors operating at the level of both primary seed dispersal and germination and establishment. According to mechanistic wind dispersal models based on the WALD kernel, the seeds of six high-mountain species are transported no farther than a few meters by primary wind dispersal, probably due to low seed release height (Dullinger *et al.*, 2011). Release height, determined by inflorescence length, is indeed low (about 5-25 cm) in both *A. caespitosa* and *S. ciliata* (see Castroviejo *et al.*, for morphological characteristics). However, according to the mechanistic simulations in Dullinger *et al.* (Dullinger *et al.*, 2011), the primary seed dispersal range of these two species should be larger

than that estimated by our effective dispersal models. This suggests that effective seed dispersal is further limited by post-dispersal processes in our study species. Environmental harshness in summer might be the main factor in reducing primary seed dispersal distance. In the two species studied here, recruitment is mainly limited at the seedling emergence and survival stages by summer drought (Garcia-Camacho *et al.*, 2010; Gimenez-Benavides *et al.*, 2008). Furthermore, García-Camacho *et al.* (Garcia-Camacho *et al.*, 2010) found a positive effect of *A. caespitosa* adult cover on seedling emergence and survival in 0.5 x 0.5 m samplings subplots. This suggests that adults act as nurse plants, facilitating seedling emergence and survival along the altitudinal gradient (Garcia-Camacho *et al.*, 2010), as observed in other Mediterranean mountain regions (Cavieres *et al.*, 2006). The effect of adult cover on post-dispersal survival might lead to unrealistic conclusions of short effective dispersal distance because our modelling approach assumes a distance-invariant survival function. This assumption may be violated in the case of distance-dependent survival because the primary and effective seed dispersal kernel would attain different shapes (Tautenhahn *et al.*, 2012). Hence, the modelling approach will implicitly assume that seedlings close to a conspecific adult originate from this adult, even though facilitation can also favour seeds from other adults, after more substantial dispersal (Tautenhahn *et al.*, 2012). Nevertheless, given the mean distances between adult plants recorded in our study plots (mean \pm SD: 5.35 \pm 0.38 m for *S. ciliata*; 5.21 \pm 0.58 m for *A. caespitosa*) and the essentially barochorous seed dispersal mechanism, the ratio of seeds from other adults to seeds from the closest adult is likely to be very small, and, therefore, the effect of this bias negligible. Moreover, in *S. ciliata*, all the populations studied exhibited significant positive fine-scale spatial genetic structure (FSGS) at the 0-1 m distance class (Lara-Romero et al unpublished data), indicating that the spatial aggregation of genotypes is consistent with a low effective dispersal distance observed in our study.

Mean dispersal distance provides a limited characterization of dispersal range (Klein *et al.*, 2006; Nathan *et al.*, 2012). Other characteristics of the dispersal kernel such as the kurtosis and fatness of the distribution tail may have important

demographic, ecological and genetic consequences (Clark *et al.*, 1999b; Klein *et al.*, 2006; Nathan *et al.*, 2011; Nathan *et al.*, 2012). Our results did not support a particular kernel shape in the case of *S. ciliata*. Although the behavior near the origin and the tail is not independent in the assumed phenomenological kernels (see (Clark, 1998) and references therein), the virtual absence of seedlings beyond the close proximity of *S. ciliata* adults (Figure S1) may result in similarly good fits for kernels with very different tails but with sufficiently fast probability decay near the origin. That is, fat-tailed kernels could fit well in *S. ciliata*, because the actual dispersal probability decreased fast near the origin and not necessarily because the actual process had a fat tail. In *A. caespitosa*, by contrast, results supported leptokurtic functions with exponential or fat tails, with consistently more abundant isolated seedlings than for *S. ciliata* (Figure S1). This would suggest a larger proportion of long-distance effective dispersal events in *A. caespitosa* than in *S. ciliata*.

The few studies that have previously used inverse modelling with a multi-population approach found local variation in dispersal parameter estimates within species (Clark *et al.*, 1998; LePage *et al.*, 2000). In Clark *et al.* (1998) and LePage *et al.* (2000) variation in dispersal patterns within species was found across tree stands of temperate forests that spanned gradients in moisture and canopy openness. Similarly, we expected higher effective dispersal range in high-altitude populations because less stressful conditions could reduce the intensity of the adult nurse effect on the survival of emerged seedlings. Furthermore, the lower encroachment by subalpine shrubs at high elevation sites (Garcia-Romero *et al.*, 2010) would provide a more open habitat with greater exposure to wind that might favour longer-distance seed dispersal. However, contrary to our expectations, there were not significant differences in the shape parameter value among-populations in any of the two species, while only the population with the largest mean dispersal distance in *A. caespitosa* differed significantly from the species average value for this parameter. Despite the strong variation in demographic structure and/or physical environment along the altitudinal gradient, the dispersal parameter estimates tended to be

consistent among populations within the species. This implies that the inherent seed dispersal traits of the species (*e.g.*, inflorescence length, fruit morphology, seed size and shape), which are essentially the same in all populations, were the most important factors determining the effective dispersal kernel in our study sites during the study period. Post-dispersal processes controlling seedling emergence and establishment seem to operate similarly across populations, regardless of evident variation in local demographic and environmental conditions (intra and interspecific plant density, local topography, soil moisture, temperature regime, etc.). This result provides a new insight into the prevalence and magnitude of intraspecific dispersal variation because, although consistently with previous studies (Clark *et al.*, 1998; LePage *et al.*, 2000) it shows that effective mean dispersal distances vary among populations of particular species, it also indicates that effective dispersal kernels can remain invariant across populations of other co-occurring species, even if there is significant variation in the demographic structure and the environment.

Effective fecundity did vary significantly across populations in both species, and per-population tests identified that these differences were due to the greater fecundity values found in the populations located at the highest altitude. As expected, the fecundity parameter was related with the ratio of seedlings to reproductive adults ($\tau = 1$, $p < 0.001$ for *S. ciliata*; and $\tau = 0.67$, $p = 0.167$ for *A. caespitosa*), which represents a measure of the average number of successfully established seedlings produced by each mother. By contrast, the fecundity parameter was not associated with mean seed production per adult plant (as estimated with a seed-crop sample from a small number of individuals of each population; data not shown). Furthermore, seedlings density was positively related with altitude ($\tau = 0.8$, $p < 0.05$ for both species). These results suggest different seedling mortality rates at different altitudes, which have probably influenced the observed differences in the fecundity parameter among populations and which are congruent with previous studies on the demography of both species (Garcia-Camacho *et al.*, 2010; Gimenez-Benavides *et al.*, 2007c), indicating that more benign conditions at the populations located at the highest altitudes allow the emergence and establishment of larger

numbers of seedlings. Hence, the altitudinal gradient appears to influence effective seed dispersal patterns in *S. ciliata* and *A. caespitosa* through variation in effective seedling establishment probabilities, but not through variation in the effective seed dispersal range. The explicit incorporation of finer-scale environmental factors and landscape features (*e.g.* shrub cover) in our models (for instance through the application of the movement space concept; Schurr *et al.*, 2008) might help to estimate and model the potential effects of environmental variables on fine-scale spatial patterns of effective dispersal more accurately, and thus to increase the amount of explained variance in within-population recruitment patterns.

The estimates provided by empirical dispersal studies such as this one could help to explicitly incorporate real migration constraints in predictive species distribution models. This information is all the more relevant because accurate predictions about dispersal and migration capacities are considered to be among the most significant uncertainties in projecting climate impacts on plant species ranges (Thuiller *et al.*, 2008). Our results, together with previous seed dispersal mechanistic simulations of alpine plant species (Dullinger *et al.*, 2011) show that the majority of dispersal events occur within a few meters from the source. This is consistent with (Dullinger *et al.*, 2012), who found widespread post-glacial dispersal constraints on the current distribution of plants in the European Alps. These findings raise doubts about the capacity of high mountain plants to track their climatic niche under the rapid climate warming predicted for mountain systems during the 21st century (Nogues-Bravo *et al.*, 2007). However, spatial spread and colonization rates are not necessarily governed by the mean seed dispersal distance but by the frequency of rare long-distance events (LDD) (Bullock and Clarke, 2000; Nathan *et al.*, 2003; Schurr *et al.*, 2009). LDD are usually caused by extreme events in terms of horizontal wind speed or turbulence (Bullock and Clarke, 2000; Nathan *et al.*, 2002). High-mountain habitats are exposed to frequent and strong updrafts (Jackson *et al.*, 2013; Whiteman, 2000). Consequently, LDD may occur frequently in high-mountain environments (Tackenberg and Stocklin, 2008). These events could greatly increase the chance of threatened populations to track the altitudinal-zone displacement

induced by warming. Therefore, further research should accurately estimate the impact of LDD on seed dispersal patterns in mountain ecosystems using adequate methods (for instance through the application of recently developed genetic methods (Burczyk *et al.*, 2006; Moran and Clark, 2011; Robledo-Arnuncio and García, 2007).

Conclusions

This study shows that there is not a single kernel function that provides consistently the best fits across species and populations. More importantly, this study points out that effective dispersal kernels can remain invariant across populations of particular species under strong variation in demographic structure and/or physical environment, while they may vary among populations of other co-occurring species. These results call for a case-by-case analysis in a wider range of taxa and environments to assess the validity of approaches that assume invariant species-specific dispersal kernels (Klein and Oddou-Muratorio, 2011).

ACKNOWLEDGEMENTS

The authors specially thank Dr. María Jose Albert for her help on the design of the study and field work. They also thank two anonymous reviewers for their valuable comments on the draft version of the manuscript, Gema Escribano-Avila who helped in the fieldwork and made comments on the manuscript and Lori De Hond for linguistic assistance.

REFERENCES

- Akaike H. 1973. Information theory as an extension of the maximum likelihood principle. In: Petrov BN, Csaki F, eds. *Second International Symposium on Information Theory*. Budapest: Akademiai Kiado.
- Austerlitz F, Dick CW, Dutech C, Klein EK, Oddou-Muratorio S, Smouse PE, Sork VL. 2004. Using genetic markers to estimate the pollen dispersal curve. *Molecular Ecology*, 13: 937-954.

- Beniston M, Stephenson DB, Christensen OB, Ferro CAT, Frei C, Goyette S, Halsnaes K, Holt T, Jylha K, Koffi B, Palutikof J, Scholl R, Semmler T, Woth K. 2007. Future extreme events in European climate: an exploration of regional climate model projections. *Climatic Change*, 81: 71-95.
- Blanquart F, Gandon S, Nuismer SL. 2012. The effects of migration and drift on local adaptation to a heterogeneous environment. *Journal of Evolutionary Biology*, 21: 1351–1363.
- Bullock JM, Clarke RT. 2000. Long distance seed dispersal by wind: measuring and modelling the tail of the curve. *Oecologia*, 124: 506-521.
- Bullock JM, Shea K, Skarpaas O. 2006. Measuring plant dispersal: an introduction to field methods and experimental design. *Plant Ecology*, 186: 217-234.
- Burczyk J, Adams WT, Birkes DS, Chybicki IJ. 2006. Using genetic markers to directly estimate gene flow and reproductive success parameters in plants on the basis of naturally regenerated seedlings. *Genetics*, 173: 363-372.
- Caspersen JP, Saprunoff M. 2005. Seedling recruitment in a northern temperate forest: the relative importance of supply and establishment limitation. *Canadian Journal of Forest Research-Revue Canadienne De Recherche Forestiere*, 35: 978-989.
- Castro J, Zamora R, Hodar JA, Gomez JM. 2005. Alleviation of summer drought boosts establishment success of *Pinus sylvestris* in a Mediterranean mountain: an experimental approach. *Plant Ecology*, 181: 191-202.
- Castroviejo S, Laínz M, López González G, Montserrat P, Muñoz Garmendia G, Paiva J, Villar L. 1990. *Flora iberica. Plantas vasculares de la Península Ibérica e Islas Baleares. Vol. II* Madrid: Real Jardín Botánico. CSIC.
- Cavieres LA, Badano EI, Sierra-Almeida A, Gomez-Gonzalez S, Molina-Montenegro MA. 2006. Positive interactions between alpine plant species and the nurse cushion plant *Laretia acaulis* do not increase with elevation in the Andes of central Chile. *New Phytologist*, 169: 59-69.
- Clark JS. 1998. Why trees migrate so fast: confronting theory with dispersal biology and the paleorecord. *American Naturalist*, 152: 204-224.
- Clark JS, Beckage B, Camill P, Cleveland B, HilleRisLambers J, Lichter J, McLachlan J, Mohan J, Wyckoff P. 1999a. Interpreting recruitment limitation in forests. *American Journal of Botany*, 86: 1-16.
- Clark JS, Macklin E, Wood L. 1998. Stages and spatial scales of recruitment limitation in southern Appalachian forests. *Ecological Monographs*, 68: 213-235.
- Clark JS, Silman M, Kern R, Macklin E, HilleRisLambers J. 1999b. Seed dispersal near and far: patterns across temperate and tropical forests. *Ecology*, 80: 1475-1494.
- Cousens R, Dytham C, Law R. 2008. *Dispersal in plants. A population perspective*. Oxford: Oxford University Press.
- Davis MB, Shaw RG, Etterson JR. 2005. Evolutionary responses to changing climate. *Ecology*, 86: 1704-1714.
- Debain S, Chadoeuf J, Curt T, Kunstler G, Lepart J. 2007. Comparing effective dispersal in expanding population of *Pinus sylvestris* and *Pinus nigra* in calcareous grassland. *Canadian Journal of Forest Research*, 37: 705-718.

- Dullinger S, Mang T, Dirnbock T, Ertl S, Gatringer A, Grabherr G, Leitner M, Hulber K. 2011. Patch configuration affects alpine plant distribution. *Ecography*, 34: 576-587.
- Dullinger S, Willner W, Plutzar C, Englisch T, Schrott-Ehrendorfer L, Moser D, Ertl S, Essl F, Niklfeld H. 2012. Post-glacial migration lag restricts range filling of plants in the European Alps. *Global Ecology and Biogeography*, 21: 829-840.
- Edwards AWF. 1992. *Likelihood, expanded edn.* Baltimore: Johns Hopkins University Press.
- Elle E, Meagher TR. 2000. Sex allocation and reproductive success in the andromonoecious perennial *Solanum carolinense* (Solanaceae). II. Paternity and functional gender. *The American Naturalist*, 156: 622-636.
- Escribano-Avila G, Sanz-Pérez V, Pías B, Virgós E, Escudero A, Valladares F. 2012. Colonization of Abandoned Land by *Juniperus thurifera* Is Mediated by the Interaction of a Diverse Dispersal Assemblage and Environmental Heterogeneity. *Plos One*, 7: e46993.
- Garcia-Camacho R, Albert MJ, Escudero A. 2012. Small-scale demographic compensation in a high-mountain endemic: the low edge stands still. *Plant Ecology & Diversity*, 25: 37-44.
- Garcia-Camacho R, Escudero A. 2009. Reproduction of an early-flowering Mediterranean mountain narrow endemic (*Armeria caespitosa*) in a contracting mountain island. *Plant Biology*, 11: 515-524.
- Garcia-Camacho R, Iriondo JM, Escudero A. 2010. Seedling dynamics at elevation limits: complex interactions beyond seed and microsite limitations. *American Journal of Botany*, 97: 1791-1797.
- Garcia-Fernandez A, Iriondo JM, Escudero A. 2012a. Inbreeding at the edge: does inbreeding depression increase under more stressful conditions? *Oikos*, 121: 1435-1445.
- García-Fernández A, Iriondo JM, Escudero A, Fuerte J, Feliner GN. 2013. Genetic patterns of habitat fragmentation and past climate change effects in the Mediterranean high-mountain plant *Armeria caespitosa* (Plumbaginaceae). *American Journal of Botany*, 100: 1641-1650.
- Garcia-Fernandez A, Lara-Romero C, Segarra-Moragues JG, Iriondo JM, Widmer A, Escudero A. 2012b. Characterization of Microsatellites in the Mountain Plant *Armeria Caespitosa* (Plumbaginaceae) and Transferability to Congeners. *American Journal of Botany*, 99: E292-E294.
- Garcia-Romero A, Munoz J, Andres N, Palacios D. 2010. Relationship between climate change and vegetation distribution in the Mediterranean mountains: Manzanares Head Valley, Sierra de Guadarrama (Central Spain). *Climatic Change*, 100: 645-666.
- Gimenez-Benavides L, Albert MJ, Iriondo JM, Escudero A. 2011a. Demographic processes of upward range contraction in a long-lived Mediterranean high mountain plant. *Ecography*, 34: 85-93.
- Gimenez-Benavides L, Dotterl S, Jurgens A, Escudero A, Iriondo JM. 2007a. Generalist diurnal pollination provides greater fitness in a plant with nocturnal pollination

- syndrome: assessing the effects of a *Silene*-*Hadena* interaction. *Oikos*, 116: 1461-1472.
- Gimenez-Benavides L, Escudero A, Iriondo JM. 2007b. Local adaptation enhances seedling recruitment along an altitudinal gradient in a high mountain Mediterranean plant. *Annals of Botany*, 99: 723-734.
- Gimenez-Benavides L, Escudero A, Iriondo JM. 2007c. Reproductive limits of a late-flowering high-mountain Mediterranean plant along an elevational climate gradient. *New Phytologist*, 173: 367-382.
- Gimenez-Benavides L, Escudero A, Iriondo JM. 2008. What shapes the altitudinal range of a high mountain Mediterranean plant? Recruitment probabilities from ovule to seedling stage. *Ecography*, 31: 731-740.
- Gimenez-Benavides L, Escudero A, Perez-Garcia F. 2005. Seed germination of high mountain Mediterranean species: altitudinal, interpopulation and interannual variability. *Ecological Research*, 20: 433-444.
- Gimenez-Benavides L, Garcia-Camacho R, Iriondo JM, Escudero A. 2011b. Selection on flowering time in Mediterranean high-mountain plants under global warming. *Evolutionary Ecology*, 25: 777-794.
- Gonzalo-Jiménez J. 2011. *Diagnóstis fitoclimática de la España peninsular*. Madrid: Organismo Autónomo de Parques Nacionales.
- Gottfried M, Pauli H, Futschik A, Akhalkatsi M, Barancok P, Alonso JLB, Coldea G, Dick J, Erschbamer B, Calzado MRF, Kazakis G, Krajci J, Larsson P, Mallaun M, Michelsen O, Moiseev D, Moiseev P, Molau U, Merzouki A, Nagy L, Nakhutsrishvili G, Pedersen B, Pelino G, Puscas M, Rossi G, Stanisci A, Theurillat JP, Tomaselli M, Villar L, Vittoz P, Vogiatzakis I, Grabherr G. 2012. Continent-wide response of mountain vegetation to climate change. *Nature Climate Change*, 2: 111-115.
- Govindaraju DR. 1988. Relationship between dispersal ability and levels of gene flow in plants. *Oikos*, 52: 31-35.
- Greene DF, Canham CD, Coates KD, Lepage PT. 2004. An evaluation of alternative dispersal functions for trees. *Journal of Ecology*, 92: 758-766.
- Hanski I. 2001. Population dynamic consequences of dispersal in local populations and in metapopulations. In: Clobert E, Danchin A, Dhondt A, Nichols JD, eds. *Dispersal*. Oxford: Oxford University Press.
- Holm S. 1979. A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics, Theory and Applications*, 6: 65-70.
- Jackson PL, Mayr G, Vosper S. 2013. Dynamically-Driven Winds. In: Chow FK, De Wekker SFJ, Snyder BJ, eds. *Mountain Weather Research and Forecasting Recent Progress and Current Challenges*. New York: Springer.
- Jongejans E, Schippers P. 1999. Modeling seed dispersal by wind in herbaceous species. *Oikos*, 87: 362-372.
- Jump AS, Peñuelas J. 2005. Running to stand still: adaptation and the response of plants to rapid climate change. *Ecology Letters*, 8: 1010-1020.

- Katul GG, Porporato A, Nathan R, Siqueira M, Soons MB, Poggi D, Horn HS, Levin SA. 2005. Mechanistic analytical models for long-distance seed dispersal by wind. *American Naturalist*, 166: 368-381.
- Klein EK, Lavigne C, Gouyon P-H. 2006. Mixing of propagules from discrete sources at long distance: comparing a dispersal tail to an exponential. *BMC Ecology*, 6: 1-10.
- Klein EK, Oddou-Muratorio S. 2011. Pollen and seed dispersal inferred from seedling genotypes: the Bayesian revolution has passed here too. *Molecular Ecology*, 20: 1077-1079.
- Larcher W. 2000. Temperature stress and survival ability of Mediterranean sclerophyllous plants. *Plant Biosystems*, 134: 279-295.
- LePage PT, Canham CD, Coates KD, Bartemucci P. 2000. Seed abundance versus substrate limitation of seedling recruitment in northern temperate forests of British Columbia. *Canadian Journal of Forest Research*, 30: 415-427.
- Levine JM, Murrell DJ. 2003. The community-level consequences of seed dispersal patterns. *Annual Review of Ecology Evolution and Systematics*, 34: 549-574.
- Martinez I, Gonzalez-Taboada F. 2009. Seed dispersal patterns in a temperate forest during a mast event: performance of alternative dispersal kernels. *Oecologia*, 159: 389-400.
- Miner BG, Sultan SE, Morgan SG, Padilla DK, Relyea RA. 2005. Ecological consequences of phenotypic plasticity. *Trends in Ecology & Evolution*, 20: 687-692.
- Moran EV, Clark JS. 2011. Estimating seed and pollen movement in a monoecious plant: a hierarchical Bayesian approach integrating genetic and ecological data. *Molecular Ecology*, 20: 1248-1262.
- Moran EV, Clark JS. 2012. Between-site differences in the scale of dispersal and gene flow in red oak. *Plos One*, 7: e36492.
- Nathan R, Horvitz N, He YP, Kuparinen A, Schurr FM, Katul GG. 2011. Spread of North American wind-dispersed trees in future environments. *Ecology Letters*, 14: 211-219.
- Nathan R, Katul GG, Horn HS, Thomas SM, Oren R, Avissar R, Pacala SW, Levin SA. 2002. Mechanisms of long-distance dispersal of seeds by wind. *Science*, 418: 409-413.
- Nathan R, Klein E, Robledo-Arnuncio JJ, Revilla E. 2012. Dispersal kernels: review. In: Clobert J, Baguette M, Benton TG, Bullock JM, eds. *Dispersal ecology and evolution*. Oxford: Oxford University Press.
- Nathan R, Muller-Landau HC. 2000. Spatial patterns of seed dispersal, their determinants and consequences for recruitment. *Trends in Ecology & Evolution*, 15: 278-285.
- Nathan R, Perry G, Cronin JT, Strand AE, Cain ML. 2003. Methods for estimating long-distance dispersal. *Oikos*, 103: 261-273.
- Nogués-Bravo D, Araujo MB, Errea MP, Martínez-Rica JP. 2007. Exposure of global mountain systems to climate warming during the 21st Century. *Global Environmental Change - Human and Policy Dimensions*, 17: 420-428.
- Oddou-Muratorio S, Klein EK, Austerlitz F. 2005. Pollen flow in the wildservice tree, *Sorbus torminalis* (L.) Crantz. II. Pollen dispersal and heterogeneity in mating success inferred from parent-offspring analysis. *Molecular Ecology*, 14: 4441-4452.

- Ouborg NJ, Piquot Y, Van Groenendael JM. 1999. Population genetics, molecular markers and the study of dispersal in plants. *Journal of Ecology*, 87: 551-568.
- Pauli H, Gottfried M, Dullinger S, Abdaladze O, Akhalkatsi M, Alonso JLB, Coldea G, Dick J, Erschbamer B, Calzado RF, Ghosn D, Holten JI, Kanka R, Kazakis G, Kollar J, Larsson P, Moiseev P, Moiseev D, Molau U, Mesa JM, Nagy L, Pelino G, Puscas M, Rossi G, Stanisci A, Syverhuset AO, Theurillat JP, Tomaselli M, Unterluggauer P, Villar L, Vittoz P, Grabbherr G. 2012. Recent plant diversity changes on Europe's mountain summits. *Science*, 336: 353-355.
- Philipp M, Madsen HES, Siegismund HR. 1992. Gene Flow and Population-Structure in *Armeria maritima*. *Heredity*, 69: 32-42.
- R Core Team. 2013. R: A language and environment for statistical computing. Vienna, Austria R Foundation for Statistical Computing.
- Ribbens E, Silander JA, Pacala SW. 1994. Seedling recruitment in forests: calibrating models to predict patterns of tree seedling dispersion. *Ecology*, 75: 1794-1806.
- Robledo-Arnuncio JJ, García C. 2007. Estimation of the seed dispersal kernel from exact identification of source plants. *Molecular Ecology*, 16: 5098-5109.
- Sagnard F, Pichot C, Dreyfus P, Jordano P, Fady B. 2007. Modelling seed dispersal to predict seedling recruitment: recolonization dynamics in a plantation forest. *Ecological Modelling*, 203: 464-474.
- Schurr FM, Spiegel O, Steinitz O, Trakhtenbrot A, Tsoar A, Nathan R. 2009. Long-distance seed dispersal. *Annual Plant Reviews*, 38: 204-237.
- Schurr FM, Steinitz O, Nathan R. 2008. Plant fecundity and seed dispersal in spatially heterogeneous environments: models, mechanisms and estimation. *Journal of Ecology*, 96: 628-641.
- Smouse P, Meagher T, Kobak C. 1999. Parentage analysis in *Chamaelirium luteum* (L.) Gray (Liliaceae): why do some males have higher reproductive contributions? *Journal of Evolutionary Biology*, 12: 1069-1077.
- Stoyan D, Wagner S. 2001. Estimating the fruit dispersion of anemochorous forest trees. *Ecological Modelling*, 145: 35-47.
- Tackenberg O, Stocklin J. 2008. Wind dispersal of alpine plant species: A comparison with lowland species. *Journal of Vegetation Science*, 19: 109-118.
- Tautenhahn S, Heilmeier H, Jung M, Kahl A, Kattge J, Moffat A, Wirth C. 2012. Beyond distance-invariant survival in inverse recruitment modeling: A case study in Siberian *Pinus sylvestris* forests. *Ecological Modelling*, 233: 90-103.
- Thuiller W, Albert C, Araujo MB, Berry PM, Cabeza M, Guisan A, Hickler T, Midgely GF, Paterson J, Schurr FM, Sykes MT, Zimmermann NE. 2008. Predicting global change impacts on plant species' distributions: Future challenges. *Perspectives in Plant Ecology Evolution and Systematics*, 9: 137-152.
- Thuiller W, Lavorel S, Araujo MB, Sykes MT, Prentice IC. 2005. Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences of the United States of America*, 102: 8245-8250.
- Tutin TG, Heywood VH, Burges NA, Valentine DH, Walters SM, Webb Da. 1995. *Flora Europaea*. Cambridge: Cambridge University Press.

- Uriarte M, Canham CD, Thompson J, Zimmerman JK, Brokaw N. 2005. Seedling recruitment in a hurricane-driven tropical forest: light limitation, density-dependence and the spatial distribution of parent trees. *Journal of Ecology*, 93: 291-304.
- Whiteman DC. 2000. Terrain-Forces Flow. *Mountain Meteorology: Fundamentals and Applications*. New York: Oxford University Press.

SUPPORTING INFORMATION

Figure S1. Spatial distribution of adults (circles) and seedlings (crosses) in each study plot of *A. caespitosa* and *S. ciliata*.




Figure S1. (A) Spatial distribution of adults (circles) and seedlings (crosses) in the Sierra de los Porrones population of *Armeria caespitosa*. Axes show UTM coordinates in meters.




Figure S1. (B) Spatial distribution of adults (circles) and seedlings (crosses) in the Collado de las Vacas population of *Armeria caespitosa*. Axes show UTM coordinates in meters.




Figure S1. (C) Spatial distribution of adults (circles) and seedlings (crosses) in the Loma de Cabezas population of *Armeria caespitosa*. Axes show UTM coordinates in meters.




Figure S1. (D) Spatial distribution of adults (circles) and seedlings (crosses) in the Najarra population of *Armeria caespitosa*. Axes show UTM coordinates in meters.




Figure S1. (E) Spatial distribution of adults (circles) and seedlings (crosses) in the Cabezas de Hierro population of *Armeria caespitosa*. Axes show UTM coordinates in meters.




Figure S1. (F) Spatial distribution of adults (circles) and seedlings (crosses) in the Laguna population of *Silene ciliata*. Axes show UTM coordinates in meters.




Figure S1. (G) Spatial distribution of adults (circles) and seedlings (crosses) in the Najarra population of *Silene ciliata*. Axes show UTM coordinates in meters.




Figure S1. (H) Spatial distribution of adults (circles) and seedlings (crosses) in the Nevero population of *Silene ciliata*. Axes show UTM coordinates in meters.




Figure S1. (I) Spatial distribution of adults (circles) and seedlings (crosses) in the Cabezas de Hierro population of *Silene ciliata*. Axes show UTM coordinates in meters.




Figure S1. (J) Spatial distribution of adults (circles) and seedlings (crosses) in the Peñalara population of *Silene ciliata*. Axes show UTM coordinates in meters.

Table S1. Estimated parameters for the models fitted to the seedling recruitment data of *Silene ciliata* (cell size length 0.25 and 0.50 m)

populati	kernel	0.25 m					0.50 m				
		δ	u	β	θ	- logL	δ	u	β	θ	- logL
Peñalara	2Dt	0.33	3.20	0.010	0.511	1034.26	0.34	7.15	0.010	0.87	556.65
	Log-norm	0.34	0.66	0.010	0.511	1035.41	0.36	0.76	0.010	0.87	558.06
	WALD	0.34	0.68	0.010	0.510	1035.79	0.36	0.42	0.010	0.86	559.07
	Exp-pow	0.32	1.21	0.010	0.509	1034.49	0.34	1.54	0.010	0.87	556.60
Cabezas de Hierro	2Dt	0.42	172.46	0.003	0.523	708.02	0.41	5.16	0.003	0.91	420.42
	Log-norm	0.42	0.50	0.003	0.531	706.53	0.43	0.62	0.003	0.91	420.47
	WALD	0.42	1.49	0.003	0.531	706.48	0.43	1.09	0.003	0.91	420.69
	Exp-pow	0.41	2789.58	0.003	0.524	705.73	0.40	1.70	0.003	0.90	420.51
Nevero	2Dt	0.25	10.44	0.001	142.159	82.84	0.44	172.60	0.001	0.72	60.14
	log-norm	0.25	0.49	0.001	142.154	82.12	0.67	0.12	0.001	1.30	58.69
	WALD	0.25	0.89	0.001	142.150	82.07	0.67	47.81	0.001	1.48	58.69
	Exp-pow	0.25	2.10	0.001	142.053	82.85	0.56	398.50	0.001	0.96	59.05
Najarra	2Dt	nc	nc	nc	nc	nc	nc	nc	nc	nc	Nc
	Log-norm	nc	nc	nc	nc	nc	nc	nc	nc	nc	Nc
	WALD	nc	nc	nc	nc	nc	nc	nc	nc	nc	Nc
	Exp-pow	0.72	0.47	0.002	0.230	337.85	0.99	0.33	0.002	0.50	225.73
Laguna	2Dt	0.25	172.44	0.001	0.003	23.23	0.25	1.76	0.001	0.02	19.30
	Log-norm	0.28	0.36	0.001	0.003	23.24	0.24	0.76	0.001	0.02	19.29
	WALD	0.27	1.89	0.001	0.003	23.22	0.24	0.33	0.001	0.02	19.24
	Exp-pow	0.29	5.87	0.001	0.004	23.11	0.20	0.58	0.001	0.02	19.30

δ , mean dispersal distance (m); u shape parameter; β , fecundity parameter (seedlings/cm); θ , negative binomial parameter; – logL, log-likelihood; nc denotes models that did not converge.

Table S2. Estimated parameters for the four models fitted to the seedling recruitment data of *Armeria caespitosa*. (cell size length 0.25 and 0.50 m).

population	kernel	0.25 m					0.50 m				
		δ	u	β	θ	- logL	δ	u	β	θ	- logL
Cabezas de Hierro	2Dt	0.40	1.91	0.024	0.261	415.97	0.40	7.20	0.024	0.80	268.95
	Log-norm	0.34	0.84	0.024	0.259	415.96	0.43	0.61	0.024	0.80	268.62
	WALD	0.34	0.40	0.024	0.256	416.10	0.43	0.96	0.024	0.80	268.74
	Exp-pow	0.31	0.78	0.024	0.257	416.41	0.39	1.85	0.024	0.80	269.01
Najarra	2Dt	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc
	Log-norm	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc
	WALD	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc
	Exp-pow	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc
Loma Cabezas	2Dt	0.65	2.88	0.014	0.196	185.10	0.69	2.28	0.014	0.65	123.64
	Log-norm	0.82	1.14	0.014	0.225	184.52	0.92	1.21	0.014	0.76	123.30
	WALD	0.73	0.40	0.014	0.219	184.84	0.67	1.26	0.014	0.60	124.43
	Exp-pow	0.65	0.58	0.014	0.224	183.92	0.67	0.52	0.014	0.77	122.65
Collado de las Vacas	2Dt	0.41	1.75	0.011	0.463	151.63	0.43	1.85	0.010	0.75	99.30
	Log-norm	0.32	0.98	0.011	0.458	152.05	0.34	0.77	0.010	0.84	99.19
	WALD	0.33	0.18	0.011	0.424	152.79	0.36	0.41	0.010	0.85	98.63
	Exp-pow	0.31	0.54	0.011	0.455	152.64	0.37	0.66	0.010	0.68	100.90
Sierra de los Porrones	2Dt	0.25	1.84	0.022	0.387	75.93	0.26	3.58	0.024	0.46	54.49
	Log-norm	0.22	0.88	0.021	0.396	76.01	0.28	0.50	0.023	0.57	53.51
	WALD	0.23	0.20	0.021	0.417	75.79	0.28	0.99	0.023	0.57	53.59
	Exp-pow	0.20	0.43	0.024	0.350	76.60	0.27	1.32	0.023	0.45	55.04

δ , mean dispersal distance (m); u shape parameter; β , fecundity parameter (seedlings/cm); θ , negative binomial parameter; – logL, log-likelihood; nc denotes models that did not converge.

CAPÍTULO 2/CHAPTER 2

Variation in fine-scale genetic structure of *Silene ciliata* (Caryophyllaceae) across populations with different spatial aggregation patterns



Carlos Lara-Romero, Alfredo García-Fernández, Juan J. Robledo-Arnuncio, Javier Morente-López, Ángela López, Marie Roumet, José M. Iriondo

Manuscrito en revisión en *Molecular Ecology*

ABSTRACT

The pattern and intensity of fine-scale spatial genetic structure (FSGS) can vary among populations within species depending on the interplay between different demographic and environmental factors. Theoretical models predict that FSGS will increase with local density variation, especially when due to static habitat heterogeneities, but few empirical studies have analyzed differences in FSGS among populations with different degrees of spatial heterogeneity. In this study, we used spatial autocorrelation methods to assess the spatial demographic and genetic structures of five demographically stable but spatially heterogeneous populations of high-mountain specialist *Silene ciliata* Poiret (Caryophyllaceae). We mapped all individuals in each population and genotyped 96 of them using 8 microsatellite markers. We found significant FSGS in three of the five populations and a significant positive association between spatial demographic and genetic structures. Contrary to previous findings in other plant species, the population with the highest plant density had the strongest FSGS, probably due to reduced long-term gene dispersal rates in fragmented habitats and lower pollinator mobility in response to high plant densities and spatial aggregation. Overall our results support the importance of the effect of fine-scale plant aggregation on intraspecific FSGS variation, and highlight the usefulness of combining FSGS analyses and the explicit characterization of local spatial distribution of individuals and habitat to better understand the mechanisms which generate intraspecific variation in FSGS across landscapes.

Keywords: environmental variation, heterogeneity, landscape, spatial ecology, spatial point pattern analysis.

INTRODUCTION

Spatial genetic structure (SGS), *i.e.* the non-random spatial distribution of genotypes, influences various genetic and evolutionary parameters and processes, such as biparental inbreeding, kin competition, effective population size and local adaptation, which in turn affect individual and population fitness (Linhart & Grant 1996; Leblois *et al.* 2006). SGS within plant populations (*i.e.*, fine-scale SGS or FSGS) results from the complex interaction between limited gene flow through seed and pollen dispersal, genetic drift, and local selection (Epperson & Li 1997; Vekemans & Hardy 2004). Under restricted seed dispersal within a continuous plant population, local inbreeding and family structures build up across generations in the absence of homogenizing selection, leading to a genetic isolation-by-distance process (IBD, Wright 1943), which is enhanced if pollen dispersal is also restricted (Crawford 1984). IBD patterns can be found in most natural plant populations as mating preferentially occurs between neighbouring individuals and the majority of seed dispersal events occur over short distances (Hamrick & Trapnel 2011). The study of fine-scale FSGS and IBD patterns are of prime interest for evolutionary biologists, because they provide indirect information on the scale of dispersal within populations (revised in Hamrick & Trapnell 2011), allow the comparison of the effects of different dispersal vectors or landscape features on historical gene flow (Vekemans & Hardy 2004; Hardy *et al.* 2006, Born *et al.* 2008), and provide knowledge on the level of within-population inbreeding and genetic drift (Wang *et al.* 2011).

The equilibrium pattern of IBD within populations is conveniently measured by the rate of decrease of the probability of gene identity with spatial distance, which is determined by the product between the rate of effective gene dispersal, σ_e , and effective population density, D_e ('neighborhood size' N_b , sensu Wright 1943; Rousset 2000). As both σ_e and D_e can vary among species and populations within species, SGS patterns should exhibit variation at both hierarchical levels. Most empirical studies comparing plant FSGS patterns have focused on the influence of

interspecific life-history variation, and particularly on testing whether species with different life forms, mating systems, and seed and pollen dispersal mechanisms show different IBD patterns (Vekemans & Hardy 2004, Hardy *et al.* 2006, Hamrick & Trapnell 2011). The extent of intraspecific (among-population) variation in FSGS and its ecological and demographic determinants have been much less studied, although historic events such as founder effects or past disturbances, intraspecific competition and microenvironmental selection have been invoked (Tero 2005, Chung *et al.* 2007; Matesanz *et al.* 2011). Differences in population density and spatial aggregation may also generate intra-specific variation in FSGS, as effective population density directly affects N_b and may affect pollinator behaviour, which in turn may influence gene dispersal (Doligez *et al.* 1998, Vekemans & Hardy 2004; Hamrick & Trapnell 2011). Overall, the magnitude of FSGS is expected to be higher in low-density populations and/or in populations where individuals are spatially aggregated (Doligez 1998; Leblois *et al.* 2004). In this sense, several recent empirical studies have found stronger FSGS in conspecific plant populations in fragmented habitats than in continuous habitats (*e.g.*, Valbuena-Carabaña 2007, Wang *et al.* 2011; but see Born *et al.* 2008). Local aggregation patterns can be generated by species' life history traits and extrinsic biotic and abiotic factors (Getzin *et al.* 2008; Chung *et al.* 2011). Theoretical models predict that SGS will increase especially with local density variation due to temporally static habitat heterogeneities (Robledo-Arnuncio & Rousset 2010), while it may remain unaffected or even decrease (relative to random distribution of individuals) if aggregation is due to density-dependent regulation in homogeneous habitats (Barton *et al.* 2002; Robledo-Arnuncio & Rousset 2010).

Few empirical studies have examined variation in FSGS among populations with different densities and degrees of spatial heterogeneity (Hamrick & Trapnell 2011), and fewer yet have combined FSGS analyses with the explicit analysis of the local spatial distribution of individuals and habitat heterogeneities (*e.g.*, Chung *et al.* 2007, 2011; Jacquemyne *et al.* 2009). Further comparative studies of FSGS across populations with different demographic and habitat characteristics are needed to

better understand the mechanisms generating intraspecific variation in FSGS across landscapes (e.g., Born *et al.* 2008, Wang *et al.* 2011).

In this study, we described the spatial demographic and genetic structures of five demographically stable, but spatially heterogeneous populations of high-mountain specialist *Silene ciliata* Poiret (Caryophyllaceae) and assessed whether the intensity of FSGS increases in populations with lower plant density and stronger spatial heterogeneity (*i.e.*, aggregation). The low effective seed dispersal distance of this species, together with short-distance pollen transport and the possibility of selfing (García-Fernández *et al.* 2012b, Lara-Romero *et al.* 2014) are likely to yield a marked within-population spatial genetic structure in equilibrium populations of *S. ciliata*. Furthermore, previous studies have shown strong variation in population structure (e.g., density and spatial distribution of individuals) and physical environment (e.g., microhabitat fragmentation and water availability) among populations (Giménez-Benavides *et al.* 2007, 2008, Lara-Romero *et al.* 2014). Therefore, *Silene ciliata* populations constitute suitable systems to study the demographic and environmental determinants of intraspecific variation in FSGS. We addressed four main questions: (1) Does the study species have a fine-scale SGS? (2) Does FSGS vary among populations? (3) If so, is this variation associated with differences in spatial structure (*i.e.*, population density and individual spatial aggregation) and/or differences in the availability and spatial heterogeneity of suitable habitat? (4) Are population genetic diversity and inbreeding related to the intensity of FSGS?

METHODS

Study region and species

Silene ciliata Poiret (Caryophyllaceae) is a chamaephytic perennial cushion plant that occurs in the Mediterranean mountain ranges of southern Europe. It is self-compatible, although autogamy is restricted by pronounced protandry (Garcia-Fernandez *et al.* 2012a). It blooms in late summer with a peak in early August

(Giménez-Benavides *et al.* 2007). Flowering stems reach 15 cm in height and have 1–5 flowers. *S. ciliata* is pollinated at night by *Hadena consparcatoides* Schawerda (Lepidoptera, Noctuidae) but can also be pollinated by diurnal insects (Giménez-Benavides *et al.* 2007). Fruit capsules, dehisced by teeth, contain up to 100 seeds that are dispersed in August-September. Because seeds lack specialized structures, seed dispersal seems to be essentially barochorous.

This study was carried out in Sierra de Guadarrama, a mountain range located in central Spain, where *S. ciliata* reaches one of its southernmost distribution limits. Its populations here are considered relicts because of their isolation from populations in other mountain systems further north. It grows locally at altitudes from 1900 to 2430 m in dry cryophilic pastures dominated by *Festuca curvifolia* Lag ex Lange and other perennial plants that are interspersed in a shrub matrix characterized by *Cytisus oromediterraneus* Rivas Mart. & al. and *Juniperus communis* subsp. *alpina* (Suter) Čelak. Mean annual precipitation, measured at the Navacerrada Pass ($40^{\circ} 46'N$, $4^{\circ} 19'W$; 1860 m a.s.l.), is 1330 mm with a pronounced dry season (< 10% of total annual rainfall) from May to October. Mean annual temperature is $6.3^{\circ}C$, with mean monthly temperatures ranging from $-1^{\circ}C$ in January to $16^{\circ}C$ in July (www.aemet.es). Although *S. ciliata* presents variable ploidy levels in natural populations, all individuals from the studied Guadarrama populations were diploid ($2n = 24$, García-Fernández *et al.* 2012b).

Study plots

In August and September 2010, we established one 10x10m plot in five *S. ciliata* populations distributed along the Sierra de Guadarrama (Figure 1, Table 1). Additional information on the selected populations can be found in Lara-Romero *et al.* 2014. We mapped all adult *S. ciliata* individuals in each study plot using two high-resolution Differential Global Navigation Satellite System (DGNS) receivers (Viva GS15, Leica, Switzerland) with an absolute accuracy of 5 cm for *x* and *y* coordinates. We also mapped the spatial position and contour of all neighboring shrubs of *Cytisus oromediterraneus* and *Juniperus communis* as well as all rocky

outcrops and rocks greater than 50 cm in diameter, which are unsuitable microhabitats for *S. ciliata*. Based on the spatial position of each adult plant, shrub, and rock, we constructed a map for each plot (Figure 1) using Quantum GIS, version 1.8 (<http://www.qgis.org>). We then used this information to estimate the global density (D) and local spatial distribution (see below) of the adult plants in each study plot, as well as the availability of suitable habitat (SH , the proportion of area covered by dry cryophilic pasture) and the spatial heterogeneity of suitable habitat, calculated with the shape index (SI). SI is widely used in landscape ecology as a measure of landscape complexity and fragmentation (Berry 2007). It is calculated as a normalized ratio of patch perimeter to area in which the complexity of patch shape is compared to a standard shape (circle or square) of the same size. It has a value of one when the patch is a perfect circle/square and increases as the patch becomes more irregular (Berry 2007). The five populations spanned a gradient in adult density (D) and had different degrees of habitat heterogeneity (Table 1).




Figure 1. Spatial distribution of adult plants (circles), rocks (grey) and shrubs (dark grey) in each study plot of *S. ciliata*.

Table 1. Description of *S. ciliata* sampling sites.

Population	UTM coordinates (m)		D	D _e	Suitable habitat (%)	SI
Peñalara	419427.30	4522814.15	12.6	3.33	71.3	1.91
Cabezas de Hierro	421169.50	4516859.61	12.8	3.29	96.8	1.14
Najarra	430196.59	4518886.29	6.09	1.54	86.2	1.42
Nevero	428863.74	4537167.57	1.98	0.54	99	1.11
Laguna	419931.16	4521082.09	1.52	0.39	48.8	2.40

UTM, x,y geographical coordinates; D, census plant density (individuals/m²); D_e, effective plant density; SI, Shape index.

DNA extraction and microsatellite genotyping

We collected leaf material from 96 adult plants per plot for genetic analysis. In order to increase sample size of shorter inter-individual distances for autocorrelation analysis, plants were selected using a clustered random sampling design (*sensu* Storfer *et al.* 2007). Thus, we randomly selected 24 sample locations in each plot, where we collected leaf material from four individuals located in close proximity to each other. Portions of leaves with no sign of parasites, fungal infection or drought injuries were collected, cleaned and dried in silica gel.

DNA was extracted using the DNeasy Plant minikit (QIAGEN, Valencia, USA) with 10–20 mg of dried *S. ciliata* tissue. Ten microsatellite loci previously used in genetic studies of *S. ciliata* were selected for genotyping: *Sci1224*, *Sci1208*, *Sci0106*, *Sci1443*, EST-2HTS, EST-8HTS, EST-37HTS, EST-X4-3, EST-G34D06 and EST-G47A02 (García-Fernández *et al.* 2012b). Specifications of PCR reactions and amplification are detailed in García Fernández *et al.* (2012b). Samples were run on an automated DNA sequencer (ABI PRISM 3730, Applied Biosystems, California, USA) in Parque Científico de Madrid (Madrid, Spain). Fragment size

determination was conducted using GeneMarker version 1.85 (SoftGenetics, State College, Pennsylvania, USA).

We evaluated genotyping accuracy by re-amplifying and re-scoring 10% of the samples. We also estimated the occurrence of null alleles, large-allele dropout and stuttering using the software MICRO-CHECKER (Van Oosterhout *et al.* 2004). We discarded loci *Sci1443* and EST-SSR X4-3 because of inconsistent allelic scoring and the possible presence of null alleles, respectively. No consistent linkage disequilibrium was found between pairwise comparisons across loci or populations (Log-likelihood ratio G statistic based on 5000 permutations performed in GENEPOP v. 4.1, Rousset, 2008).

Spatial distribution of reproductive plants within populations

We employed spatial point pattern analyses to characterize the local spatial structure of adult plants in each of the five study populations. Specifically, we used the noncumulative O-ring statistic $O(r)$ (Wiegand & Moloney 2004), which is a derivative of the K -function that gives the expected number of points (*i.e.*, plants) in a ring (not a circle, as in the K -function) of radius r centred at an arbitrary point (Wiegand and Moloney 2004). We also used $O(r)$ to detect heterogeneities in the point pattern. These heterogeneities are found when the function departs from the expected value for a random pattern at scales where plant-plant interactions should become unimportant, in a process known as “virtual aggregation” (Wiegand and Moloney 2004). We set the threshold where plant-plant interactions are not relevant at 1.5 m, which is over twice the mean effective seed dispersal distance detected in the studied populations (Lara-Romero *et al.* 2014).

We calculated $O(r)$ considering concentric rings of constant width $w = 0.1$ m and an increasing (0.1 m lag) radius r centred at each plant, and plotted $O(r)$ as a function of the distance r to the plant. To avoid estimation bias, distance classes were defined between $r = 0.10$ and 2.5 m, *i.e.* up to a quarter of the plot width (Baddeley and Turner 2005). We used Ripley’s isotropic correction (Baddeley and

Turner 2005) to correct for edge effects. The association between $O(r)$ and r was tested by computing $O(r)$ and comparing these values with those expected under the null hypothesis of complete spatial randomness (CSR). The 95% simulation envelopes associated to the null hypothesis of CSR were generated using the 10th-lowest and 10th-highest value of 399 Monte Carlo simulations of the CSR null model. $O(r)$ values above and below the envelope indicate significant spatial clustering and repulsion compared to a CSR pattern, respectively.

Finally, to compare the overall strength of spatial structure among populations, we calculated the slope ($b_{O(r)}$) of the linear regression of $O(r)$ on the logarithm of the spatial distance r and its associated 95% confidence intervals (95% CI) for each population using least-squares regression theory (Zar 1999). We performed pairwise *t*-tests to determine whether $b_{O(r)}$ significantly differed among populations. All statistical analyses were performed in the open source software R v 3.0.2 using the package ‘spatstats’ (Baddely & Turner 2005).

Population genetic analyses

We used GENEPOP v. 4.1 (Rousset, 2008) to calculate standard population genetic measures, including total number of alleles (A), observed (H_O) and expected (H_E) heterozygosity, and inbreeding coefficients (F_{IS}). We tested for statistical differences in A , H_E and F_{IS} across populations using the non-parametric Friedman rank sum test (Hollander & Wolf 1973) implemented in the R package ‘stats’ (R Core Team, 2013). Deviations from Hardy Weinberg (HW) equilibrium within each population were tested using the Fisher’s exact test implemented in GENEPOP. Genetic differentiation between populations (F_{ST}) was calculated using FSTAT v. 2.9.3 (Goudet 1995). The significance of F_{ST} values was evaluated based on 10 000 random permutations of multilocus genotypes among populations (Excoffier *et al.* 1992).

Characterization of within-population SGS

Bayesian clustering analysis

Within-population spatial discontinuities in genetic variation can arise from biotic and abiotic landscape features that could act as a barrier to within-population gene flow (Storfer *et al.* 2007). To identify such genetic discontinuities in each plot (*e.g.*, due to the spatial distribution of shrub and rock microhabitats), we used a Bayesian clustering method implemented in the R package ‘GENELAND’ version 4.0.3 (Guillot *et al.* 2005a). This approach uses Markov chain Monte Carlo (MCMC) procedures to estimate the number of genetic clusters, which is treated as an unknown parameter (Guillot *et al.* 2005a,b, 2008). Five independent runs of the spatial D-model were performed with 100 000 MCMC iterations, of which every hundredth one was saved. The spatial D-model approximates the domain of each population by the union of a few polygonal domains. This model corresponds with the spatial patterns expected under limited gene flow induced by the presence of physical barriers (Guillot *et al.* 2005a,b, 2008). We assumed linkage equilibrium between alleles across loci.

Spatial Genetic Autocorrelation Analysis

We used the spatial autocorrelation approach implemented in SPAGeDI version 1.4 (Hardy & Vekemans 2002) to assess FSGS resulting from IBD. This approach, described by Vekemans and Hardy (2004), is based on genetic kinship coefficients between pairs of individuals i and j (F_{ij} , Loiselle *et al.* 1995). To test the overall pattern of SGS in each population, F_{ij} were regressed on the spatial distance between individuals, d_{ij} , and its natural logarithm, $\ln(d_{ij})$, giving the regression slopes (b_d and b_{Ld}) and the coefficients of determination (R^2_d and R^2_{Ld}) for each type of regression. Standard errors for F_{ij} , b_d and b_{Ld} were estimated by jackknifing over loci. We also obtained average pairwise F_{ij} over a set of distance intervals (d) comprised between 0.5 and 10m (above this threshold the number of pairwise comparisons was low in some populations). For each distance interval,

computed values were compared to a 95% confidence interval around the null hypothesis of absence of SGS (*i.e.*, F_{ij} and d_{ij} or $\ln(d_{ij})$ are uncorrelated). The 95% CI was constructed by performing 9999 random permutations of genotypes among spatial positions.

To compare the intensity of SGS among populations, we calculated the statistic Sp for each population. Sp reflects the rate of decrease of F_{ij} with distance and is robust to sampling design (Vekemans & Hardy 2004). We used the formula $Sp = -b_{dl}/(1-F_1)$, where F_1 is mean F_{ij} at the smallest distance interval d . Mean Sp and standard error (SE) were calculated for each population over all loci, and the approximate 95% CIs were obtained as ± 1.96 times the SE estimates obtained across loci. We carried out pairwise t -tests to determine the statistical significance of the differences found between populations.

We also used the software SPAGeDI to estimate Wright's neighbourhood size (Nb , Wright 1943) and the effective gene dispersal rate (σ_e). In two-dimensional populations, Nb is related to σ_e and the effective population density (D_e) in the form $Nb = 4\pi D_e \sigma_e^2$ (Rousset 2000). Under Wright's IBD model, Nb can be estimated as $Nb = -(1-F_1)/b_r$ where b_r is the restricted regression slope (b_{ld}) of F_{ij} on $\ln(d_{ij})$ in the range $\sigma_e < d < 20 \sigma_e$ (Vekemans and Hardy, 2004). D_e can be conveniently approximated as $D * N_e/N$ where N_e/N is the ratio of the effective over census population sizes. As suggested by Vekemans & Hardy 2004, we assumed that density was constant over time and computed N_e/N from $N_e/N \equiv 4/[2(1-F_1) + (1 + F_1)V]$ (Kimura & Crow 1963), where V corresponds to the variance of the lifetime reproductive success among individuals. To estimate the value of V , we used the demographic data reported by Giménez-Benavides *et al.* (2011) and applied the equation $V = p^2 X_M^2 V_L + p^2 X_L V_M + X_L X_M V_p$ (Barrowclough & Rockwee 1993), where L , M and p denote reproductive life span, individual fecundity within a breeding season and probability of survival to breeding status of each propagule unit, respectively; with mean X and variance V . Following this procedure, the ratio of effective to census population sizes (N_e/N) was estimated at 0.264, 0.257, 0.252,

0.257 and 0.248, in Peñalara, Cabezas de Hierro, Najarra, Nevero and Laguna, respectively. These N_e/N ratios yielded effective densities ranging from 0.39 to 3.33 ind/m² (Table 1). We used these estimates of D_e to compute σ_e and Nb , applying the iterative procedure implemented in SPAGeDI.

Approximate 95% confidence intervals around Nb estimates were computed as $(F_I - 1)/(b_r + 2SE_b)$ and $(F_I - 1)/(b_r - 2SE_b)$, respectively, where SE_b is the standard error of b_r estimated by jackknifing over loci. When $b_r > 2SE$, the upper bound was reported as infinite (Fenster *et al.* 2003, Hardy *et al.* 2006). The 95% CI of σ_e was obtained using an analogous procedure.

Additionally, we used F_{IS} and F_{ij} to obtain indirect estimates of the selfing rate adjusted for biparental inbreeding, $\hat{s} = 2(F_{IS} - F_X)/(1 + F_{IS} - 2F_X)$, where F_X is the coefficient of biparental inbreeding (Fenster *et al.* 2003). Following Chung *et al.* (2007), we used the intercept of the regression of pairwise F_{ij} on the natural logarithm of distance (b_{Ld}) as estimator of F_X . This is a conservative estimate of selfing rate because \hat{s} assumes that outcrossing is restricted to nearest neighbor mating, where biparental inbreeding is highest (Chung *et al.* 2007).

Association between the spatial demographic and genetic structure

$O(r)$ shares the ring-based approach with spatial genetic autocorrelation, making the results of our spatial demographic and genetic analyses directly comparable (Chung *et al.* 2007, 2011). To test for statistic association between the spatial demographic and genetic structure, we conducted correlation analysis between b_{Ld} and $b_{O(r)}$ (Chung *et al.* 2007; Chung 2008). We also assessed the correlation between Sp and adult density (D), the availability of suitable habitat (SH) and their shape index (SI).

RESULTS

Within-population spatial structure of reproductive plants

The O-ring statistic $O(r)$ showed a significant pattern of individual spatial aggregation at small scales in all of the studied populations (Figure 2A). Peñalara showed the strongest overall aggregation, with a slope of the O-ring statistic of $b_{O(r)} = -4.73$ (95% CI: $-5.22, -4.24$). Laguna and Nevero exhibited intermediate patterns ($b_{O(r)} = -2.28$, 95% CI = $(-2.71, -1.85)$ and $b_{O(r)} = -2.19$ ($-2.35, -2.03$), respectively), while Cabezas de Hierro and Najarra showed the flattest functions ($b_{O(r)} = -1.73$ ($-1.87, -1.59$) and $b_{O(r)} = -1.18$ ($-1.30, -1.06$), respectively). The comparison of pairwise t -tests showed significant differences in $b_{O(r)}$ values between all studied population pairs except between Nevero and Laguna (Supplementary data, Table S1). Heterogeneities in point pattern intensity were detected in all populations by the departure of $O(r)$ from CSR at large scales (*i.e.*, above 1.5 m) (Figure 2A).




Fig 2. Spatial demographic structure and spatial genetic structure in *Silene ciliata* study plots using the O-ring statistic for individual plants $O_{(r)}$ with ring radius r and ring width 0.1 m (Panel A) and the genetic kinship coefficient, F_{ij} , (Panel B), respectively. **Panel A:** $b_{O(r)}$ denotes the slope of the regression of $O_{(r)}$ on the logarithm of the spatial distance, $\ln(r)$; the thin solid line indicates average intensity of the point pattern λ (i.e., plant density), and dashed lines indicate 95% confidence envelopes under the null hypothesis of random spatial structure. Note that y-axes show different scales for clarity. **Panel B:** b_{Ld} denotes the slope of the regression of kinship coefficient values on the logarithm of the spatial distance, $\ln(dij)$; dashed lines represent 95% confidence envelopes constructed under the null hypothesis (no genetic autocorrelation exists)..

Population genetic analyses

We scored a total of 144 different alleles across eight loci in 474 individuals (Table 2), with a mean (\pm SE) number of 10 (0.8) alleles per locus. The average number of alleles ranged from 9 (\pm 1.9) in Najarra to 12 (\pm 2) in Nevero. Observed heterozygosities varied from 0.344 (\pm 0.090) in Laguna to 0.392 (\pm 0.079) in Cabezas de Hierro, while expected heterozygosities ranged from 0.504 (\pm 0.097) in Peñalara to 0.548 (\pm 0.098) in Nevero. Inbreeding coefficients (F_{IS}) varied between 0.235 (Peñalara) and 0.349 (Laguna). All populations showed significant departures from H-W equilibrium towards heterozygote deficiency, according to global tests across loci. The Friedman test did not detect significant differences among populations for any of the genetic diversity parameters. Indirect estimates of selfing rate \hat{s} were 0.40, 0.45, 0.48, 0.45 and 0.52 in Peñalara, Cabezas de Hierro, Najarra, Nevero and Laguna, respectively. Detailed genetic diversity estimates for each marker and population are provided as Supplementary Data (Table S2).

Population differentiation (F_{ST}) was low but significant ($F_{ST} = 0.052$, $p < 0.001$). Pairwise F_{ST} values were significantly different from zero for all population pairs (Supplementary data, Table S3), ranging from 0.030 (Peñalara - Nevero) to 0.082 (Najarra - Laguna).

Characterization of within-population SGS

All five independent runs of GENELAND gave a single genetic group within each of the five populations, indicating that there were no genetic discontinuities or spatially structured gene pools.

No significant relationship was found between genetic kinship (F_{ij}) and geographic distance between individuals in Cabezas de Hierro and Najarra (b_L and b_{Ld} were not statistically different from 0, Figure 2B, Table S4). However, the slopes of the linear regressions between F_{ij} and both geographic distance and the logarithm of geographic distance were significantly negative in Peñalara, Nevero and Laguna (Figure 2B, Table S4), as expected under IBD. In all three populations, the decrease

of F_{ij} with distance better fitted a logarithmic relationship than a linear relationship ($R^2_{dL} > R^2_d$) (Table S4). The three populations with significant IBD showed significantly positive F_{ij} values in the first distance class ($d \leq 0.5$ m), rapidly declining and becoming non-significant in the second or third distance classes (Figure 2B). The Sp statistic confirmed the differences in the intensity of FSGS among populations (Table S4). Peñalara showed the strongest SGS with $Sp = 0.0161$ (95% CI: 0.0127, 0.0221), followed by Laguna and Nevero, $Sp = 0.0117$ (0.029, 0.0205) and $Sp = 0.0096$ (0.0033, 0.0159), respectively. The lowest Sp values were found in Najarra and Cabezas de Hierro with $Sp = 0.0064$ (0.0015, 0.0113) and $Sp = 0.0009$ (0, 0.0054), respectively. Pairwise t -tests indicated that Sp at Peñalara was significantly greater than at Najarra and Cabezas de Hierro ($t = 2.27, p < 0.05$ and $t = 2.70, p < 0.01$; respectively).

Table 2. Genetic diversity estimates for *S. ciliata* populations.

Population	<i>N</i>	<i>A</i> (SE)	H_O (SE)	H_E (SE)	F_{IS}
Peñalara	96	10 (1.6)	0.384 (0.092)	0.504 (0.097)	0.235*** (5)
Cabezas de Hierro	94	10 (2.1)	0.392 (0.079)	0.548 (0.091)	0.285*** (6)
Najarra	95	9 (1.9)	0.360 (0.074)	0.524 (0.095)	0.316*** (4)
Nevero	96	12 (2)	0.391 (0.091)	0.548 (0.092)	0.283*** (5)
Laguna	93	11 (1.8)	0.344 (0.090)	0.531 (0.084)	0.349*** (6)

N, sample size; *A*, average number of alleles per locus; H_O and H_E , expected and observed heterozygosities; F_{IS} , average inbreeding coefficient; Number in parentheses in F_{IS} column are the number of loci that deviate significantly from H-W equilibrium after bonferroni correction; SE, standard error; *** $P < 0.001$. There were no significant differences among populations for any of the genetic diversity estimators.

Wright's neighbourhood size (Nb) and the extent of gene dispersal (σ_e) were assessed in the three populations which had a SGS consistent with IBD. Estimates of neighbourhood size were $Nb = 158$ individuals (95% CI: 121, 195) for Peñalara, $Nb = 103$ individuals (91, ∞) for Laguna and $Nb = 106$ individuals (81, ∞) for Nevero. Nb values for Peñalara were about one-tenth of the total number of individuals within the plot, while in Nevero and Laguna they accounted for about 40 and 60% of total plot size, respectively. Using the independently estimated effective densities D_e (Table 1), we obtained a notably lower σ_e estimate for Peñalara (1.94 m, 95% CI: 1.74, 2.14) than for Nevero ($\sigma_e = 3.9$ m, 95% CI: 3.69, ∞) and Laguna ($\sigma_e = 4.65$ m, 95% CI: 4.17, ∞).

Association between the spatial demographic and genetic structure

We found a significant strong association between spatial autocorrelation measures of genetic structure (b_{Ld}) and individual spatial aggregation ($b_{O(r)}$) (Pearson's product-moment correlation: $r = 0.92$, $p = 0.022$, $n = 5$). No significant associations were found between Sp and D ($r = -0.15$; $p = 0.805$, $n = 5$), SH ($r = -0.69$; $p = 0.198$, $n = 5$) or SI ($r = 0.65$; $p = 0.230$, $n = 5$).

DISCUSSION

This study found significant fine-scale spatial genetic structure (FSGS) consistent with an isolation-by-distance process (IBD) in three of the five studied populations of *S. ciliata*, as well as significant among-population variation in the intensity of FSGS. Spatial demographic and genetic structures were positively correlated, supporting the importance of differences in the fine-scale aggregation of plants for intraspecific FSGS variation. Contrary to our expectations, global population density was not significantly associated with FSGS. Population genetic diversity and inbreeding did not vary significantly as a function of the intensity of FSGS.

Effect of individual spatial aggregation on FSGS

Assuming dispersal-drift equilibrium, the observed differences in SGS among the study populations, as measured by Sp , should mainly result from differences in the product between effective dispersal rate and effective population density (*i.e.*, the genetic neighbourhood size, inversely related to Sp ; Wright 1943, Rousset 2000, Vekemans & Hardy 2004). Effective gene dispersal σ_e is properly defined as the asymptotic mean-square dispersal distance of gene lineages across generations (Rousset 2004). If favourable habitat is not continuous, and especially if the discontinuities remain static across generations, some gene lineages will remain between the edges of a favourable habitat patch over several generations until an inter-patch dispersal event occurs. This landscape resistance to gene diffusion will result in σ_e values that may be substantially lower than the mean-square parent-offspring distance (σ) (Robledo-Arnuncio & Rousset 2010). Therefore, SGS is expected to be stronger in spatially aggregated populations than in continuous populations, even if parent-offspring dispersal distances remain unchanged. The significant positive correlation found between spatial demographic and genetic structure in this study is consistent with these theoretical predictions and suggests that habitat-driven modifications in local density and σ_e generate intra-specific variation in FSGS, even if average seed dispersal distances are largely invariant across populations (Lara-Romero *et al.* 2014).

Although previous empirical studies have suggested that environmental heterogeneity could be a potential determinant of observed SGS variation (Valbuena-Carabaña 2007, Born *et al.* 2008, Wang *et al.* 2011), its precise association with local density heterogeneities and its ensuing effects on effective gene dispersal rates have rarely been measured. The few studies that have quantified both spatial and genetic structure across plant populations found significant FSGS in spatially heterogeneous populations (Jacquemyne *et al.* 2009; Chung *et al.* 2011), while weak or no FSGS was found in continuously distributed populations (Chung *et al.* 2007; Chung *et al.* 2008). However, these studies compared stable continuous

populations to expanding and senescent spatially heterogeneous populations, or seedlings to adult plants within the same population. As far as we know, this is the first empirical study reporting evidence of a possible relationship between the spatial aggregation of plants and intra-specific FSGS among stable plant populations.

Influence of habitat availability and global plant density on FSGS

Although Peñalara had higher census and effective population densities, it had a significantly stronger FSGS than Cabezas de Hierro and Najarra. The stronger plant aggregation in this population may be due to low suitable habitat availability ($SH = 71.3\%$) and high habitat heterogeneity ($SI = 1.71$) compared to Cabezas de Hierro ($SH = 96.8 \%, SI = 1.14$) and Najarra ($SH = 86.2\%, SI = 1.42$). The habitat-driven fine-scale aggregation of adults in Peñalara would lead us to expect a notably lower effective gene dispersal rate in this population, which might partially explain its greater FSGS. However, FSGS in Peñalara was not significantly different than that found in Nevero and Laguna, even though Peñalara had a ten-fold larger population density and six-to-eight-fold larger effective density, which should have theoretically resulted in a much larger neighbourhood size (much weaker FSGS). We hypothesize that this lack of significant differences in FSGS may be due to the competing effects of higher effective population density (which tends to decrease FSGS) and lower effective dispersal rate (which tends to increase FSGS) in the former. Consistent with the hypothesized negative effect of individual aggregation on effective gene dispersal rates, σ_e was considerably lower in Peñalara ($\sigma_e = 1.94$ m) than in Nevero and Laguna ($\sigma_e = 3.9$ m and 4.65 m, respectively). In plants, σ_e can be decomposed into effective seed ($\sigma_{e,s}$) and pollen ($\sigma_{e,p}$) gene dispersal rates:

$$\sigma_e^2 = \sigma_{e,s}^2 + \frac{1}{2} \sigma_{e,p}^2 \quad (\text{Crawford 1984})$$

Assuming as a rough approximation that $\sigma_{e,s}$

equals contemporary field estimates of mother-seedling axial square dispersal distance (σ_s) obtained through inverse modelling, we obtained $\sigma_s = 0.27, 0.21$ and 0.20 m for Peñalara, Nevero and Laguna, respectively (Lara-Romero *et al.* 2014). Then, following Crawford (1984), the effective pollen gene dispersal rate $\sigma_{e,p}$ in

Peñalara was 1.35 m, less than half of the values calculated for Nevero and Laguna (2.75 m and 3.28 m, respectively). This might be due to greater pollinator mobility, resulting in higher pollen dispersal distances, in low-density and more continuous populations during foraging activities (Richards 1999; Born *et al.* 2008; Cörtes *et al.* 2013). Similar results were found for congeneric species *Silene tatarica*, where the *Sp* statistic was higher in denser populations as a consequence of restricted pollen dispersal (Tero *et al.* 2005).

Habitat heterogeneity may have also been produced by unaccounted factors in the study sites, such as interspecific competition. In particular, the fine-scale spatial aggregation of adults in Nevero, which is totally free of rocks and shrubs, could be explained by the presence of dominant patches of *Festuca curvifolia* and interspersed gaps where the rest of the plant species tend to occur (Sanchez-Pescador *et al.* 2014). This may explain the strong spatial aggregation of *S. ciliata* individuals observed in this population, as well as the lack of correlation between *Sp* and the availability (SH) and heterogeneity (SI) of suitable habitat.

Association between variation in FSGS and genetic diversity and inbreeding

We found homogeneous values of genetic diversity (H_E) and coefficient of inbreeding (F_{IS}) across populations, suggesting independence between FSGS and population genetic diversity and inbreeding. All populations were characterized by high, significant F_{IS} values. Such a pronounced level of inbreeding may be produced by selfing (*i.e.*, self-pollination) and/or biparental inbreeding. In the present case we observed rather high levels of self-pollination in all populations (between 0.40 and 0.52) confirming previous evidence of the mixed mating system of this species (García-Fernández *et al.* 2012a,b), but detected significant pattern of FSGS in only three of the populations. Thus, the occurrence of self-pollination in all populations (rather than biparental inbreeding induced by non-random spatial distribution of genotypes) seems to be the primary factor responsible for the high level of inbreeding observed in our study species.

Concerning the high level of diversity (H_E) observed in strongly structured population (characterized by a high value of sp) it may be related to the occurrence of inter-population gene flow. Indeed, García-Fernández *et al.* (2012b) previously found substantial gene flow among populations at the same study site, which is supported by the low F_{ST} values obtained between our study populations. Furthermore, as previously discussed, low plant density may enhance gene dispersal by modifying pollinator behaviour, preventing genetic erosion in small populations (Born *et al.* 2008, Còrtes *et al.* 2013). On the other hand, model predictions of the effect of IBD on the rate of decrease of allelic richness and heterozygosity have obtained higher or lower rates of decrease (relative to random mating populations) depending on factors such as population size, dispersal shape and range, habitat geometry and mutation (Leblois *et al.* 2006). All of this suggests that FSGS can be concurrent with a certain level of within and among-population gene dispersal and with maintained levels of genetic diversity within populations (Chung *et al.* 2007; Born *et al.* 2008; Wang *et al.* 2011). This may partially explain the lack of association between FSGS, plant density and genetic diversity found in this study.

Conclusions

This study found a significant positive relationship between the intensities of spatial demographic and genetic structures. These findings support theoretical studies showing that stronger spatial aggregation of adult individuals generally increases the magnitude of SGS compared to random or regular individual distributions (Doligez 1998, Barton *et al.* 2002; Robledo-Arnuncio & Rousset 2010). No significant association was found between global plant density and FSGS. Contrary to findings in other species (Vekemans and Hardy 2004, Hardy *et al.* 2006), the population with the highest plant density was also the population with strongest FSGS, probably due to reduced long-term gene dispersal rates in fragmented habitats, and possibly lower pollinator mobility in response to high plant density and aggregation. This study stresses the importance of combining FSGS analyses and the explicit characterization of the local spatial distribution of individuals and habitat to better

understand the mechanisms generating intraspecific variation in FSGS across landscapes.

ACKNOWLEDGEMENTS

We thank Alex Widmer (ETH-Zurich) for his valuable comments on the manuscript and Lori De Hond for linguistic assistance. We also thank Marcelino de la Cruz (ESCET, Universidad Rey Juan Carlos) who gave us helpful recommendations for analyzing spatial point pattern data and the staff of Sierra de Guadarrama National Park for permission to work in the field area. This work was supported by the project AdAptA (CGL2012-33528). CLR and JM were supported by a F.P.I. fellowship (Spanish Ministry of Science).

REFERENCES

- Baddeley A, Turner R (2005) Spatstat: an R package for analyzing spatial point patterns. *Journal of statistical software* **12**, 1-42.
- Barrowclough GF, Rockwell RF (1993) Variance of lifetime reproductive success: estimation based on demographic data. *American Naturalist*, 281-295.
- Barton NH, Depaulis F, Etheridge AM (2002) Neutral evolution in spatially continuous populations. *Theoretical Population Biology* **61**, 31-48.
- Berry JK (2007) *Map analysis: understanding spatial patterns and relationships* GeoTec Media, Los Angeles, CA.
- Born C, Hardy OJ, Chevallier MH, et al. (2008) Small-scale spatial genetic structure in the Central African rainforest tree species *Aucoumea klaineana*: a stepwise approach to infer the impact of limited gene dispersal, population history and habitat fragmentation. *Molecular Ecology* **17**, 2041-2050.
- Cortes MC, Uriarte M, Lemes MR, et al. (2013) Low plant density enhances gene dispersal in the Amazonian understory herb *Heliconia acuminata*. *Molecular Ecology* **22**, 5716-5729.
- Crawford TJ (1984) The estimation on neighbourhood parameters for plant populations. *Heredity* **52**, 273-283.
- Chung MY (2008) Variation in demographic and fine-scale genetic structure with population-history stage of *Hemerocallis taeanensis* (Liliaceae) across the landscape. *Ecological research* **23**, 83-90.

- Chung MY, Nason JD, Chung MG (2007) Effects of population succession on demographic and genetic processes: predictions and tests in the daylily *Hemerocallis thunbergii* (Liliaceae). *Molecular ecology* **16**, 2816-2829.
- Chung MY, Nason JD, Chung MG (2011) Significant demographic and fine-scale genetic structure in expanding and senescing populations of the terrestrial orchid *Cymbidium goeringii* (Orchidaceae). *American journal of botany* **98**, 2027-2039.
- Doligez A, Baril C, Joly HlnI (1998) Fine-scale spatial genetic structure with nonuniform distribution of individuals. *Genetics* **148**, 905-919.
- Epperson BK, Li T-Q (1997) Gene dispersal and spatial genetic structure. *Evolution* **51**, 672-681.
- Excoffier L, Smouse PE, Quattro JM (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* **131**, 479-491.
- Fenster CB, Vekemans X, Hardy OJ (2003) Quantifying gene flow from spatial genetic structure data in a metapopulation of *Chamaecrista fasciculata* (Leguminosae). *Evolution* **57**, 995-1007.
- Garcia-Fernandez A, Iriondo JM, Escudero A (2012a) Inbreeding at the edge: does inbreeding depression increase under more stressful conditions? *Oikos* **121**, 1435-1445.
- García-Fernández A, Segarra-Moragues JG, Widmer A, Escudero A, Iriondo JM (2012b) Unravelling genetics at the top: mountain islands or isolated belts? *Annals of botany* **110**, 1221-1232.
- Getzin S, Wiegand T, Wiegand K, He F (2008) Heterogeneity influences spatial patterns and demographics in forest stands. *Journal of Ecology* **96**, 807-820.
- Giménez-Benavides L, Escudero A, Iriondo JM (2007) Reproductive limits of a late-flowering high-mountain Mediterranean plant along an elevational climate gradient. *New Phytologist* **173**, 367-382.
- Giménez-Benavides L, Escudero A, Iriondo JM (2008) What shapes the altitudinal range of a high mountain Mediterranean plant? Recruitment probabilities from ovule to seedling stage. *Ecography* **31**, 731-740.
- Goudet J (1995) FSTAT (version 1.2): a computer program to calculate F-statistics. *Journal of heredity* **86**, 485-486.
- Guillot G, Estoup A, Mortier F, Cosson JF (2005a) A spatial statistical model for landscape genetics. *Genetics* **170**, 1261-1280.

- Guillot G, Mortier F, Estoup A (2005b) GENELAND: a computer package for landscape genetics. *Molecular Ecology Notes* **5**, 712-715.
- Guillot G, Santos F, Estoup A (2008) Analysing georeferenced population genetics data with Geneland: a new algorithm to deal with null alleles and a friendly graphical user interface. *Bioinformatics* **24**, 1406-1407.
- Hamrick JL, Trapnell DW (2011) Using population genetic analyses to understand seed dispersal patterns. *Acta Oecologica* **37**, 641-649.
- Hardy OJ, Maggia L, Bandou E, et al.. (2006) Fine-scale genetic structure and gene dispersal inferences in 10 Neotropical tree species. *Molecular ecology* **15**, 559-571.
- Hardy OJ, Vekemans X (2002) SPAGeDi: a versatile computer program to analyse spatial genetic structure at the individual or population levels. *Molecular ecology notes* **2**, 618-620.
- Jacquemyn H, wiegand T, Vandepitte K, Brys R, Roldán-Ruiz I, Honnay O (2009) Multigenerational analysis of spatial structure in the terrestrial, food-deceptive orchid *Orchis mascula*. *Journal of Ecology* **97**, 206-216.
- Kimura M, Crow JF (1963) The measurement of effective population number. *Evolution*, 279-288.
- Lara-Romero C, Robledo-Arnuncio JJ, García-Fernández A, Iriondo JM (2014) Assessing intraspecific variation in effective dispersal along an altitudinal gradient: a test in two Mediterranean high-mountain plants. *PLOS ONE* **9**, e87189.
- Leblois R, Rousset F, Estoup A (2004) Influence of spatial and temporal heterogeneities on the estimation of demographic parameters in a continuous population using individual microsatellite data. *Genetics* **166**, 1081-1092.
- Leblois R, Estoup A, Streiff R (2006) Genetics of recent habitat contraction and reduction in population size: does isolation by distance matter? *Molecular Ecology* **15**, 3601–3615.
- Linhart YB, Grant MC (1996) Evolutionary significance of local genetic differentiation in plants. *Annual Review of Ecology and Systematics* **27**, 237-277.
- Loiselle BA, Sork VL, Nason J, Graham C (1995) Spatial genetic structure of a tropical understory shrub, *Psychotria officinalis* (Rubiaceae). *American Journal of Botany*, 1420-1425.
- Matesanz S, Gimeno TE, de la Cruz M, Escudero A, Valladares F (2011) Competition may explain the fine-scale spatial patterns and genetic structure of two co-occurring plant congeners. *Journal of Ecology* **99**, 838-848.

- Pescador DS, Chacón-Labella J, de la Cruz M, Escudero A (2014) Maintaining distances with the engineer: patterns of coexistence in plant communities beyond the patch-bare dichotomy. *New phytologist* in press.
- Richards CM, Church S, McCauley DE (1999) The influence of population size and isolation on gene flow by pollen in *Silene alba*. *Evolution* **53**.
- Robledo-Arnuncio JJ, Rousset F (2010) Isolation by distance in a continuous population under stochastic demographic fluctuations. *Journal of evolutionary biology* **23**, 53-71.
- Rousset F (2000) Genetic differentiation between individuals. *Journal of Evolutionary Biology* **13**, 58-62.
- Rousset F (2008) genepop'007: a complete re-implementation of the genepop software for Windows and Linux. *Molecular ecology resources* **8**, 103-106.
- Storfer A, Murphy MA, Evans JS, et al.. (2007) Putting the 'landscape' in landscape genetics. *Heredity* **98**, 128-142.
- Tero N, Aspi J, Siikamäki P, Jäkäläniemi A (2005) Local genetic population structure in an endangered plant species, *Silene tatarica* (Caryophyllaceae). *Heredity* **94**, 478-487.
- Valbuena-Carabaña M, González-Martínez SC, Hardy OJ, Gil L (2007) Fine-scale spatial genetic structure in mixed oak stands with different levels of hybridization. *Molecular ecology* **16**, 1207-1219.
- Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* **4**, 535-538.
- Vekemans X, Hardy OJ (2004) New insights from fine-scale spatial genetic structure analyses in plant populations. *Molecular Ecology* **13**, 921-935.
- Wang R, Compton SG, Chen X-Y (2011) Fragmentation can increase spatial genetic structure without decreasing pollen-mediated gene flow in a wind-pollinated tree. *Molecular ecology* **20**, 4421-4432.
- Wiegand T, Moloney K (2004) Rings, circles, and null-models for point pattern analysis in ecology. *Oikos* **104**, 209-229.
- Wiegand T, Gunatilleke S, Gunatilleke N, Okuda T (2007) Analyzing the spatial structure of a Sri Lankan tree species with multiple scales of clustering. *Ecology* **88**, 3088-3102.
- Wright S (1943) Isolation by distance. *Genetics* **28**, 114.
- Zar JH (1999) *Biostatistical analysis* Prentice Hall, New Jersey.

SUPPORTING INFORMATION**Table S1.** *t* values for *S. ciliata* pairwise $b_{O(r)}$ comparisons and their statistical significance.

	Peñalara	Cabezas de Hierro	Nevero	Najarra	Laguna
Peñalara	-				
Cabezas de Hierro	11.56***	-			
Nevero	9.68***	4.33***	-		
Najarra	13.81***	5.97***	10.10***	-	
Laguna	7.36***	2.38**	0.38 ^{n.s.}	4.82***	-

ns $P > 0.05$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Table S2. Genetic diversity indices for each microsatellite loci and population of *S. ciliata*. Na , number of alleles per locus; H_O and H_E , observed and expected heterozygosities; F_{IS} , average inbreeding coefficient.

Population	Locus	Na	H_O	H_E	F_{IS}
Peñalara	SSR-Sci1224	10	0.677	0.736	0.08
	SSR-Sci1208	9	0.740	0.768	0.046
	SSR-Sci0106	19	0.221	0.788	0.721*
	SSR-G34D06	9	0.656	0.706	0.071*
	EST-G47A02	5	0.083	0.130	0.359*
	EST-37HTS	12	0.323	0.437	0.262*
	EST-2HTS	4	0.146	0.166	0.186
	EST-8HTS	8	0.240	0.327	0.268*
C. de Hierro	SSR-Sci1224	8	0.620	0.663	0.065
	SSR-Sci1208	7	0.456	0.708	0.358*
	SSR-Sci0106	20	0.376	0.864	0.579*
	SSR-G34D06	10	0.733	0.780	0.06*
	EST-G47A02	4	0.128	0.196	0.351*
	EST-37HTS	12	0.278	0.474	0.416*
	EST-2HTS	4	0.096	0.170	0.439*
	EST-8HTS	17	0.462	0.552	0.163*
Nevero	SSR-Sci1224	11	0.740	0.814	0.091*
	SSR-Sci1208	6	0.521	0.649	0.198*
	SSR-Sci0106	20	0.289	0.810	0.644*
	SSR-G34D06	13	0.763	0.737	-0.036
	EST-G47A02	4	0.042	0.082	0.489*
	EST-37HTS	8	0.209	0.331	0.37*
	EST-2HTS	18	0.250	0.585	0.574*
	EST-8HTS	13	0.313	0.397	0.213*
Najarra	SSR-Sci1224	7	0.522	0.624	0.165*
	SSR-Sci1208	8	0.653	0.717	0.09*
	SSR-Sci0106	21	0.358	0.847	0.579*
	SSR-G34D06	9	0.484	0.712	0.322*
	EST-G47A02	3	0.053	0.102	0.484*
	EST-37HTS	10	0.233	0.572	0.593*
	EST-2HTS	5	0.116	0.150	0.23*
	EST-8HTS	9	0.463	0.487	0.049*

Laguna	SSR-Sci1224	9	0.602	0.656	0.082
	SSR-Sci1208	6	0.688	0.744	0.076
	SSR-Sci0106	20	0.250	0.703	0.646*
	SSR-G34D06	12	0.641	0.768	0.166*
	EST-G47A02	3	0.054	0.143	0.625*
	EST-37HTS	8	0.130	0.358	0.636*
	EST-2HTS	14	0.215	0.616	0.652*
	EST-8HTS	12	0.174	0.280	0.38*
Mean	-	10	0.374	0.531	0.299*

Table S3. Matrix of pairwise differentiation (F_{ST}) among populations. * F_{ST} values significantly different from zero.

	Peñalara	Cabezas de Hierro	Nevero	Najarra	Laguna
Peñalara	-				
Cabezas de Hierro	0.033*	-			
Nevero	0.030*	0.037*	-		
Najarra	0.055*	0.045*	0.071*	-	
Laguna	0.043*	0.07*	0.054*	0.082*	-


Table S4. Estimates of within-population spatial genetic structure at five populations of *S. ciliata*.

Population	F_1 (SE)	b_d (R_d^2)	b_{Ld} (R_{Ld}^2)	Sp (SE)
Peñalara	0.059*** (0.012)	-0.0037*** (0.0045)	-0.0161*** (0.0086)	0.0161 (0.0042)
Cabezas de Hierro	0.006 ^{ns} (0.008)	-0.0011 ^{ns} (0.0004)	-0.0035 ^{ns} (0.0004)	0.0009 (0.0023)
Najarra	0.023 ^{ns} (0.019)	-0.0013 ^{ns} (0.0004)	-0.0048 ^{ns} (0.0005)	0.0064 (0.0025)
Nevero	0.038** (0.012)	-0.0013* (0.0008)	-0.0083** (0.0030)	0.0096 (0.0032)
Laguna	0.034*** (0.021)	-0.0025** (0.0022)	-0.0105*** (0.0032)	0.0117 (0.0045)

F_I , mean F_{ij} value between individuals separated at the minimum distance interval; b_d and b_{Ld} , denote slopes of the regression of kinship coefficient values on the spatial distance, dij , and its logarithm, $\ln(dij)$, respectively. The associated coefficient of determination R^2 is shown in brackets; Sp , strength of the genetic structure calculated for the minimum distance class in each population; SE, standard error. ns $P > 0.05$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

CAPÍTULO 3/CHAPTER 3

Population spatial patterns conditioned by dispersal and heterogeneity reveal negative density dependence on reproductive output in spite of positive spatial associations



Carlos Lara-Romero, Marcelino de la Cruz, Gema Escribano-Ávila, Alfredo García-Fernández, José M. Iriondo

Manuscrito inédito

ABSTRACT

1. The spatial patterns of plant species may be generated by an array of ecological features going from environmental constraints to intrinsic traits of the species. This makes difficult to distinguish density dependent effects from other process fostering plant co-occurrence, such as the existence of environmental heterogeneity or specific dispersal patterns.
2. Here, we studied the spatial pattern of the Mediterranean high-mountain specialist *Silene ciliata* Poiret (Caryophyllaceae) at five populations located along an altitudinal gradient. We aimed to depict the spatial structure of the species and its relation to plant performance across different populations assessing the contribution of dispersal, environmental heterogeneity and conspecific plant interactions to the observed patterns.
3. We used homogeneous and in-homogenous pair correlation functions to describe the spatial pattern of both recruits and adult plants and assess the contribution of environmental heterogeneity to the pattern. Then, we applied a case-control design with a null model of random-labelling to evaluate whether the spatial pattern induced by seed dispersal and recruitment persists in the adult class and to analyse the spatial relationships between both stage classes. Finally, we analysed the effects of conspecific neighbour density on individual reproductive performance to test for density dependence on reproductive output.
4. Recruits exhibited a pronounced small-scale aggregation and large-scale heterogeneity that were well characterized by a heterogeneous Poisson cluster process. The aggregation induced by seed dispersal and subsequent seedling recruitment persisted in the adult class.
5. The spatial pattern of recruits after seed germination and establishment was independent of the spatial pattern of adults. Even though the aggregated structure of adult plants might suggest the presence of facilitative interactions among them, a negative density-dependent effect on reproductive output was found.
6. *Synthesis.* This study provides empirical evidence for the lack of a simple and direct relationship between plant population structure and the sign of plant-plant

interactions (*i.e.*, competition and facilitation) and outlines the importance of include all relevant ecological factors involved in determinant plant spatial patterns, such as seed dispersal and environmental heterogeneity if we want to fully understand underlying mechanisms of plant spatial patterns.

Keywords: plant-plant interactions, point pattern analyses, microsite selection, spatial aggregation, competition, *Silene*, Mediterranean alpine grasslands.

INTRODUCTION

The strong role of plant-plant interactions in determining the spatial distribution of individuals within populations reveals the importance of the local neighbourhood context for plant fitness, demography and ecosystem processes (Theodore & Bowman, 1997; Jones & Comita, 2008; Reid & Lortie, 2010). Earlier studies have hypothesized that the outcome of plant interactions may vary inversely across gradients of abiotic stress, with positive interactions (facilitation) being more common in stressful environments and negative interactions (competition) prevailing in more benign abiotic conditions (The Stress-Gradient Hypothesis; Bertness & Callaway, 1994). This theoretical model has been verified in alpine and arctic ecosystems (Callaway *et al.*, 2002), but other studies have reported non-positive or even negative interactions under extremely stressful conditions (Maestre *et al.*, 2005; Dvorský *et al.*, 2013). These controversial results have led to refine the Stress-Gradient Hypothesis stating that the predicted outcome of plant-plant interactions would depend more frequently on the type of abiotic stress and the life history of the species (Maestre *et al.*, 2009; Soliveres & Maestre, 2014). Plant-plant interaction studies usually assume that positive spatial associations detected between plants at a certain scale are the result of a facilitative interaction (McIntire & Fajardo, 2009; Soliveres & Maestre, 2014; but see Callaway *et al.*, 2002; Jones & Comita, 2008). Nevertheless, the mechanisms underlying observed spatial patterns may be generated by an array of ecological features going from environmental constraints to intrinsic traits of the species (Schupp & Fuentes, 1995; Plotkin *et al.*, 2000; Turnbull *et al.*, 2004). This makes difficult to distinguish the effects of biotic interactions from other process driving plant co-occurrence, such as the existence of spatial variation in habitat quality (*i.e.*, habitat heterogeneity) or specific dispersal patterns (Schupp & Fuentes, 1995; North & Ovaskainen, 2007; Getzin *et al.*, 2008).

Limited seed dispersal leads to spatial aggregation of seedlings by generating higher local seed density around mother plants (Howe, 1989; Seidler & Plotkin, 2006). The distribution of dispersal distances is controlled by inherent seed dispersal traits of the species, such as inflorescence length, fruit morphology, seed size and

shape (Howe, 1989; Schupp & Fuentes, 1995; Seidler & Plotkin, 2006). Spatial aggregation initially caused by short-distance dispersal and subsequent seedling recruitment can be modified by habitat heterogeneity and density dependent effects on plant performance (Seidler & Plotkin, 2006; North & Ovaskainen, 2007; Lara-Romero *et al.*, 2014). Habitat heterogeneity has been previously highlighted as a possible confounding signal of positive plant–plant interactions as reinforces plant aggregation by promoting higher density of conspecifics in locally favourable environments (Getzin *et al.*, 2008; Chacón-Labolla *et al.*, 2014). Conversely, negative density dependence tends to reduce the aggregation of conspecifics seedlings when attacks by natural enemies (*e.g.*, pathogens, herbivores) and/or competition for limited resources increase with local conspecific density (Jones & Comita, 2008; Comita & Hubbell, 2009). Nevertheless, neutral and positive role of conspecific adult neighbours on seedling emergence and survival has also been documented (Crawford & Whitney, 2010; Lara-Romero *et al.*, 2014).

Despite the importance of spatial structure of plants to plant population dynamics (Theodore & Bowman, 1997; Jones & Comita, 2008; Reid & Lortie, 2010) we still do not know up to what extent aggregation or segregation in plant populations can be attributed to the existence of positive or negative interactions among its individuals. In the case that a direct relationship between plant structure and plant-plant ecological interactions exist if such relationship is conserved across life–stages and populations is something far from understood. This implies that the mechanism and ecological processes fostering the observed spatial patterns remain in many cases unrevealed. To contribute to filling this gap of knowledge, here we studied the spatial pattern of the Mediterranean high-mountain specialist *Silene ciliata* Poiret (Caryophyllaceae) at five populations located along an altitudinal gradient. We aimed to depict the spatial structure of the species and its relation to plant performance across different populations assessing the contribution of dispersal, habitat heterogeneity and conspecific plant interactions to the observed patterns. *S. ciliata* recruitment is mainly constrained by summer drought at the seedling emergence and survival stages (Gimenez-Benavides *et al.*, 2007; Gimenez-

Benavides *et al.*, 2011). Furthermore, it has been suggested that adults facilitate emergence and survival of conspecific seedlings leading to a reduction of primary seed dispersal distance (Lara-Romero *et al.*, 2014). Consequently, we expected a marked spatial aggregation between seedling recruits (seedlings surviving the first summer season) and adults of *S. ciliata*. Moreover, we hypothesized this spatial aggregation to be conserved along subsequent life stages, as the main limiting recruitment processes of the species (*i.e.*, seedling emergence and survival) have already been overcome (Gimenez-Benavides *et al.*, 2007; Gimenez-Benavides *et al.*, 2011). Finally, we expected a strong effect of the environmental heterogeneity on the spatial pattern within and between populations (Pescador *et al.*, 2014; Chapter 2).

We used state-of-the-art techniques of spatial point pattern analysis (Wiegand & Moloney, 2004; Law *et al.*, 2009) to decompose the realized spatial patterns into biotic and abiotic underlying processes applying, at a wide array of scales, specific a priori hypotheses linked to spatial pattern creation (McIntire & Fajardo, 2009; Brown *et al.*, 2011). We first described the spatial pattern of both recruits and adult plants and assessed the contribution of environmental heterogeneity to the pattern by contrasting the plant spatial arrangement of each population to four null models, namely homogeneous and heterogeneous Poisson and Poisson cluster processes (Diggle, 2003; Waagepetersen, 2007; Wiegand *et al.*, 2013). Secondly, we used a case-control design using a null model of bivariate random-labelling to contrast the patterns of both recruits and adults (De La Cruz *et al.*, 2008; Getzin *et al.*, 2008; Chacón-Labella *et al.*, 2014). This allowed us to evaluate whether the spatial pattern induced by seed dispersal and subsequent seedling recruitment persists in the adult class and also to analyse the spatial relationships between recruits and adults (*e.g.*, attraction vs segregation). Finally, we analysed the effects of conspecific neighbour density on individual reproductive performance to assess whether plant fitness is density dependent. The three analyses, specific hypotheses and related tables and figures are summarized in Table 1.

Table 1. Research hypotheses, statistical analyses and related figures of the study

Hypotheses	Statistical analyses/Null models	Related figures and tables
(1) The spatial distribution of recruits and adults of <i>Silene ciliata</i> within and between populations is affected by environmental heterogeneity.	<i>g</i> -functions of adults and recruits contrasted to homogeneous and heterogeneous Poisson (<i>i.e.</i> , CSR) and Poisson cluster processes.	Fig 1, Table 2
(2) The spatial distribution of both stage classes is spatially structured even after having statistically corrected for heterogeneity		
(3) The pattern induced by seed dispersal and subsequent seedling recruitment persist in the adult class.	Evaluation of <i>g</i> -functions under a case-control design using a null model of bivariate random labelling: <i>a</i> = control = adults, <i>s</i> = cases = recruits.	Fig 2
(4) Adults facilitate seedling recruitment, and thus, both stage classes are spatially correlated.		
(5) The density of conspecifics around each adult increases individual reproductive performance	Linear mixed model (LMM) for number of flowerheads, with plant size and density of conspecific neighbours around each adult plant as predictor continuous variable and population as random factor.	Table 3

METHODS

Study site and species

Silene ciliata Poiret (Caryophyllaceae) is a chamaephytic perennial cushion plant that occurs in the Mediterranean mountain ranges of southern Europe. It is self-compatible, although autogamy is restricted by pronounced protandry (Garcia-Fernandez *et al.*, 2012). It blooms in late summer, with a peak in early August (Gimenez-Benavides *et al.*, 2007). Flowering stems reach 15 cm in height and have 1–5 flowers. *S. ciliata* is pollinated at night by *Hadena consparcatoides* Schawerda (Lepidoptera, Noctuidae) but can also be pollinated by diurnal insects (Gimenez-Benavides *et al.*, 2007). Fruit capsules, dehiscing by teeth, contain up to 100 seeds that are dispersed in August-September. Because its seeds lack specialized structures, seed dispersal is essentially barochorous and most seed are dispersed at

short distance (*i.e.*, 50 cm) (Lara-Romero *et al.*, 2014). The study was carried out in Sierra de Guadarrama, a mountain range located in central Spain, where *S. ciliata* reaches one of its southernmost distribution limits. It grows locally at altitudes from 1900 to 2430 m in orophyllous cryophilic pastures dominated by *Festuca curvifolia* Lag ex Lange and other perennial plants that are interspersed in a shrub matrix characterized by *Cytisus oromediterraneus* Rivas Mart. & al. and *Juniperus communis* subsp. *alpina* (Suter) Čelak.

Field data

In August and September 2010, we established a plot of approx. 100m² in five populations of *S. ciliata* distributed along an altitudinal gradient at the study site (Table S1). The five populations spanned gradients in plant density and climatic conditions (Table S1). We mapped every *S. ciliata* adult plant and seedling recruit (hereafter recruit) in each study plot (Figure S1) using two high-resolution Differential Global Navigation Satellite System (DGNSS) receivers (Viva GS15, Leica, Switzerland) with an accuracy of 5 cm for *x* and *y* coordinates. We also measured the number of inflorescences of each adult plant. This variable is significantly related to total seed production (Gimenez-Benavides *et al.*, 2007), thus we used it as a plant fitness correlate. Our definition of recruits includes up to one-year individuals (*i.e.*, plants that had germinated in fall or spring and had survived their first full summer), because the seedlings that survive a second growth period grow into the reproductive stage (Gimenez-Benavides *et al.*, 2011). Within this framework, we included all critical stages of effective dispersal: seed dispersal, seed germination and seedling recruitment.

Summary statistics: the g-functions

We employed spatial point pattern analyses to characterize the spatial structure of recruits and adult plants at each of the five study populations. Specifically, we used test statistics based on the pair-correlation functions $g(r)$, which are a derivative of the Ripley's K function (Stoyan & Stoyan, 1994; Wiegand & Moloney, 2004). The function $\lambda K(r)$ is the expected number of points (*i.e.*, plants) within a distance *r* of an

arbitrary point (Ripley, 1976), where λ is the intensity of the pattern in the study area (*i.e.*, plant density) and $K(r)$ is the univariate K function. The univariate pair correlation function $g(r)$ is related to Ripley's $K(r)$ in the form $g(r) = K'(r)/(2\pi r)$ (Stoyan & Stoyan, 1994; Wiegand & Moloney, 2004). It gives the expected number of points in a ring (not a circle, as in the K -function) of radius r and width w centred at an arbitrary point, divided by λ . Bivariate extension of the pair correlation function $g_{as}(r)$ and $g_{sa}(r)$, comprising type “*a*” and type “*s*” points (where type “*a*” represents adult plants and type “*s*” recruits) are defined as $\lambda_s K_{as}(r)/(2\pi r)$ and $\lambda_a K_{sa}(r)/(2\pi r)$, respectively (Wiegand & Moloney, 2004).

We calculated univariate and bivariate g -functions considering concentric rings of constant width $w = 0.1$ m and increasing (1 cm lag) radius r centred at each plant, and plotted $g(r)$ as a function of the distance r to the plant. To avoid estimation bias, g -functions were estimated between $r = 0.10$ and 2.5 m, *i.e.* up to a quarter of the plot width (Baddeley & Turner, 2005). We contrasted the empirical g -functions to those expected under specific null models (see below) which represent point processes with known structure, thus allowing the comparison of observed patterns to those expected under biological hypotheses (Wiegand & Moloney, 2004).

Spatial pattern of recruits and adult plants

To select the null model that appropriately describes the spatial dispersion of recruits and adults in each population and to assess the existence of large-scale heterogeneity (Hypotheses 1 and 2, Table 1), we assumed the pattern reconstruction strategy proposed by Wiegand *et al.*, (2013). We first fitted four commonly employed spatial (null) models for each class: a) a homogeneous Poisson process (*i.e.*, complete spatial randomness: CSR) with constant intensity λ equal to the density of the observed pattern; b) a homogeneous Poisson cluster process with constant intensity λ and parameters σ and ρ fitted by minimum contrast (Diggle, 2003), where ρ is the density of clusters and σ^2 denotes the variance of the normal distribution $h(r, \sigma)$ describing the locations of points relative to the cluster centre (Jacquemyn *et al.*, 2007; Wiegand *et al.*, 2007); c) a heterogeneous Poisson process

with $\lambda(x,y)$ estimated with a Gaussian kernel (Wiegand *et al.*, 2007; Pescador *et al.*, 2014) with a bandwidth of 3.7 m, d) a heterogeneous Poisson cluster processes (Waagepetersen, 2007) with parameters σ and ρ fitted by minimum contrast and intensity functions estimated as in c). Heterogeneous models account for spatial variation of λ related to habitat heterogeneity (*i.e.*, “first-order effects” *sensu* Wiegand & Moloney, 2004). We specified a bandwidth of 3.7m on the basis that at this distance the cumulative probability of seedling recruitment is over 95% for all study population (Lara-Romero *et al.*, 2014), and therefore, it should be a sufficient distance to remove all potential heterogeneities in the point pattern without affecting plant-plant interactions and the pattern of small-scale clustering (*i.e.*, “second-order effects” *sensu* Wiegand & Moloney, 2004) that tend to occur at scales lower than 2 m in the study species (Pescador *et al.*, 2014). When evaluating heterogeneous models, we substituted the homogeneous- $g(r)$ by the inhomogeneous- $g(r)$ (Baddeley & Turner, 2005).

To evaluate whether a given null model is able to describe and quantify the empirical spatial structures of recruits and adults in each population, 95% simulation envelopes associated to a given null model were generated by calculating for each distance r the 10th-lowest and 10th-highest $g(r)$ value from 399 Monte Carlo simulations of each null model. The simulation envelopes cannot be interpreted as confidence intervals because the null hypothesis is tested simultaneously at many scales, which may inflate Type-I error (Loosmore & Ford, 2006). We therefore evaluated the overall fit of four alternative spatial models with a goodness-of-fit (GOF) test in the range $r = 0.10$ and 2.5 using the u statistic (Diggle, 2003; Loosmore & Ford, 2006),

$$u = \int_{r=0.1}^{r=2.5} \{ \hat{g}(r) - g(r) \}^2 dr$$

where $\hat{g}(r)$ is the empirical estimation of the pair correlation function, $g(r)$ the mean of the pair correlation functions of 399 patterns simulated according to the

evaluated null model. Therefore, u represents the total squared deviation between the empirical pattern and the theoretical result across the distances of interest. If u of the observed pattern is larger than those of 380 simulations (of the 399 total simulations), the null hypothesis is rejected with at the 5% significance level. When two spatial models were not rejected for a spatial pattern, we selected the model which produced the smallest u value as the "appropriate" model for describing spatial structure of recruits and adults in each population (Pescador *et al.*, 2014).

Difference in the strength of clustering and spatial relationship between recruits and adult plants

We used the null model of random labelling within a case-control design to test whether the strength of clustering varies between recruits and adults and to investigate the spatial relationship between both stage classes (Hypotheses 3 and 4, Table 1). Case-control designs are commonly used in epidemiology and have recently been applied to point pattern analysis (De La Cruz *et al.*, 2008; Getzin *et al.*, 2008), where the control pattern is used to account for environmental heterogeneity in the population of interest (*i.e.*, cases). The random-labelling null model assumes that the pattern of cases is identical from that of controls and therefore cases are a random subset of the population group composed of cases and controls (De La Cruz *et al.*, 2008; Getzin *et al.*, 2008). We used recruits as “cases” and compared them to the pattern of adults (*i.e.*, controls). Random labelling implies that K and g -functions are invariant, and therefore, we expected that $g_s(r) = g_a(r) = g_{as}(r)$. Departure from random labelling indicates that the process assigning labels to the points does not act randomly.

Strength of clustering of recruits and adult plants

To evaluate if the strength of clustering varies across stage classes (Hypotheses 3, Table 1), we applied the null-model, $g_a(r) - g_s(r)$, which has an expected value of 0 for all r distances. Significant negative values at short scales of r indicate that recruits are more clustered than adults, suggesting the existence of a thinning

process. Positive deviations at short scales would indicate increased aggregation with ontogeny (Getzin *et al.*, 2008; Chacón-Labella *et al.*, 2014).

Recruit-adult relationship

We used the null-model $g_{as}(r)$ (Goreaud & Pélissier, 2003) to test if the pattern of recruits shows a significant association with the pattern of adults (Hypotheses 4, Table 1). Positive departures from the null model would indicate attraction of recruits to adults (*i.e.*, clumps of recruits located near adults), which suggest some facilitative effects of adults on germination and/or establishment. Conversely, spatial segregation (*i.e.*, recruits spatially segregated from adults) would be indicated by negative departures from the null model.

We also employed the differences $g_{aa}(r) - g_{as}(r)$, and $g_{ss}(r) - g_{sa}(r)$, which have also an expected value of 0 under random labelling. These functions evaluate whether one type of point tends to be more frequently surrounded by points of the other type than by points of its own type. Hence, if $g_{aa}(r) - g_{as}(r) = 0$, the relative concentration of recruits near adults is similar than that of adults, indicating that adult and seedlings exploit the available habitat in the same way. Negative deviations at short scales would indicate increased regeneration in the proximity of adults, and thus, the existence of some facilitative effects of adults on seed germination or seedling establishment, while positive deviations would indicate segregation between adults and seedlings (Getzin *et al.*, 2008; Chacón-Labella *et al.*, 2014). A value of $g_{ss}(r) - g_{sa}(r) \gg 0$, on other hand, would indicate the existence of an additional clustering mechanism within recruits, irrespective from the spatial pattern of the adults, and therefore, from the “first-order” environmental heterogeneity (Getzin *et al.*, 2008).

Significant departure from random labelling of the observed difference-functions at each distance r was assessed by computing 95% simulation envelopes from 399 Monte Carlo simulations of each random labelling null model. The overall departure from random labelling was evaluated computing GOF test in the range $r = 0.10$ and 2.5m.

Density-dependent effects on reproductive performance

We used a linear mixed model (LMM) to test whether the strength of spatial clustering affected individual reproductive performance. The model included number of flowerheads of each adult plant as dependent variable, the clustering of conspecific around each adult and plant size as predictor continuous variable and population as random factor to control for potential spatial non-independence. The clustering around each adult plant (hereafter conspecific neighbour density) was the intensity of the point pattern in its neighbourhood λ_i , which was estimated with a Gaussian kernel with a bandwidth equal to the maximum distance to a nearest-neighbour in each study population (Diggle, 2003). Response variables were ln-transformed to reach normality. We included plant size in the LMM because it is positively related with production of flowerheads in the study species (Gimenez-Benavides *et al.*, 2007). This variable was entered first in the models to control its potential effects before calculating the effects of conspecific neighbour density on number of flowerheads.

We performed all statistical and spatial analyses in the open source software R. v. 3.0.3 (R Core Team, 2014). Most of the spatial analyses were conducted with the R package “spatstat” (Baddeley & Turner, 2005), while differences between g -functions were calculated with the R package “ecespa” (De la Cruz, 2008). LMM model was fitted with the R package “lmer” (Pinheiro *et al.*, 2014).

RESULTS

We mapped a total of 832 recruits and 3643 adults across all populations of the species (Figure S1 and Table S1 in Supporting Information). Adult and recruit densities varied between populations (Table S1). The average (\pm SD) of adult individuals per population was 728 ± 509 individuals (ranging from 1330 in Peñalara to 149 in Laguna) meanwhile the seedling average was 166 ± 172 (ranging from 469 in Peñalara to 10 in Laguna).

Spatial pattern of recruits and adult plants

When we contrasted the spatial pattern of recruits and adults in the five study populations to the four null models we found that the pattern of *S. ciliata* recruits and adults were consistent with a heterogeneous Poisson cluster (HPC) process in all five populations (GOF tests: all $p > 0.05$). In all cases, the HPC model provided the best fits across all populations (*i.e.*, GOF test with the lowest u value), while the empirical in-homogeneous $g(r)$ functions lied completely within the simulation envelopes of this null model (Fig 1). This indicates that the spatial patterns of recruits and adults exhibit a pronounced small-scale aggregation and large-scale heterogeneity. On the contrary, null models based on Poisson processes were rejected (all GOF test: $p < 0.001$).




Fig 1. Patterns of adults (top) and recruits (bottom) of *Silene ciliata* contrasted to the null model of Poisson cluster process, using the inhomogeneous pair correlation function $g(r)$. The $g(r)$ is contrasted to 95% simulation envelopes (grey area) constructed using 399 simulations of the Poisson cluster processes. The solid and dashed lines denote the observed pattern and the expectation of the null model, respectively.

The parameters ρ and σ of the HPC model measure basic characteristics of clustering of the empirical patterns, and allow detecting possible differences between the evaluated patterns. The pattern is more clustered when there are fewer clusters or when cluster size, defined as $r_C \approx 2\sigma$, is smaller (Jacquemyn *et al.*, 2007). For the fitted HPC processes the number of clusters ranged between $A\rho = 30$ and $A\rho = 200$ for adult plants and between $A\rho = 10$ and $A\rho = 133$ for recruits (Table 2). Cluster size ranged between $r_C = 0.36$ and $r_C = 0.88$ m and between $r_C = 0.16$ and $r_C = 0.72$ m for adults and recruits, respectively (Table 2). According to these parameters, the process was more clustered for recruits than for adult plants in all the study populations. Similarly, differences were also found among populations within stage classes (Table 2).

Table 2. Estimated parameters for the heterogeneous Poisson cluster (HPC) model fitted for recruits and adult plants of *S. ciliata*.

Population					
Peñalara	Cabezas de Hierro	Nevero	Najarra	Laguna	
HPC fitted for adults					
σ^2	0.096	0.13	0.096	0.194	0.032
ρ	0.94	2	0.28	1.2	0.46
r_c	0.62	0.72	0.62	0.88	0.36
$A\rho$	99	202	30	125	45
HPC fitted for recruits					
σ^2	0.053	0.13	0.068	0.014	0.006
ρ	0.75	0.54	0.21	1.27	0.1
r_c	0.46	0.72	0.52	0.24	0.16
$A\rho$	79	55	22	133	10

σ^2 , variance; ρ , number of cluster/m²; $A\rho$, number of clusters (A is the area of the study plot); r_c cluster radius (m).

Difference in the strength of clustering and spatial relationship between recruits and adult plants*Strength of clustering of recruits and adult plants*

In all populations, the difference $g_a(r) - g_s(r)$ showed significant negative departures from the null model of random labelling at short scales (GOF test: all $p < 0.05$; Fig. 2), indicating that the strength of clustering declined systematically from recruits to adults and therefore the existence of a thinning process due to seedling mortality in their transition to adults.

Recruit-adult relationship

With the function g_{as} empirical spatial patterns did not depart significantly from the null model of random labelling (GOF test: all $p > 0.05$; Fig. 2) in all populations except in Najarra, which showed a slight spatial segregation of recruits from adults pointed by negative departures from the null model (GOF test: $p = 0.01$; Fig. 2).

The empirical $g_{aa}(r) - g_{as}(r)$ difference did not depart significantly from the random labelling null model at scales between 0.10 and 2.5 m (GOF test: all $p > 0.05$; Fig. 2), indicating a spatially homogeneous mixing in the distribution of recruits and adults. However, Najarra was once again the exception to the overall pattern due to the spatial segregation between adult and recruits indicated by significant positive departure from the null model (GOF test: all $p = 0.01$; Fig. 2). Finally, the difference $g_{ss}(r) - g_{sa}(r)$ revealed strong departures at small scales from random labelling in all populations (GOF test: all $p > 0.05$; Fig. 2), which indicates that an additional aggregation mechanism of recruits occurs, independent of the adult pattern.




Fig 2. Differences of pair correlation functions $g(r)$ using a null model of random labelling, where subscript “ a ” and “ s ” indicate adult and recruits of *Silene ciliata*, respectively. Approximately 95% simulation envelopes (grey area) were constructed using 399 simulations of the random labelling model. The solid and dashed lines denote the observed pattern and the expectation of the null model, respectively.

Density-dependent effects on reproductive performance

Reproductive plant performance was negatively affected by conspecific neighbour once corrected by plant size, despite a substantial random effect of population (Table 3). Thus, big plants in low-dense conspecific neighbourhoods yielded the highest reproductive performance.

Table 3. LMM results for testing whether plant size and neighbourhood density affects reproductive performance.

Effect	Solution for effects				Deviance change	
	β	SE	t	p	$F_{1, 3636}$	p
<i>Fixed effects</i>						
Intercept	0.53	0.27	1.97	0.048	36.12	<0.001
Plant size	0.016	0.001	45.09	<0.001	2047	<0.001
Neighbours density	-0.010	0.002	-5.39	<0.001	29.08	<0.001
<i>Random effects (Population)</i>						
Intercept					SD	
					0.6	

SD, standard deviation.

DISCUSSION

Despite the strong variation in plant demography and climatic conditions across the studied populations the spatial pattern of *S. Ciliata* was consistently explained by the Heterogeneous Poisson cluster process (HPC) and both stage classes exhibited a pronounced small-scale aggregation and large-scale heterogeneity in all studied populations. Nevertheless, the pattern induced by seed dispersal and subsequent seedling recruitment was thinned. Thus adults showed a decrease in the strength of clustering compared to recruits, likely due to the impact of mortality in the transition from recruit to adult. Contrary to our expectations the spatial pattern of recruits after seed germination and establishment was independent of the spatial pattern of adults. Even though the aggregated structure of adult plants might suggest the presence of facilitative interactions among them, a negative density-dependent effect on

reproductive output was found. We suggest that the aggregated pattern observed is the result of environmental heterogeneity and short distance dispersal of the species together with the availability of safe-sites for establishment, with a reduced effect of plant-plant interactions. Thus, our results provide empirical evidence for the lack of a simple direct correlation between the presence of spatial structure and the sign of intraspecific interactions (aggregation and segregation meaning facilitation and competition, respectively) across all populations and life history stages.

Habitat heterogeneity

The heterogeneity in the spatial pattern of *S. ciliata* adults indicates the existence of relevant differentiation in environmentally driven habitat quality (*i.e.*, a non-random distribution of resources), as adult individuals are expected to exploit all favourable sites for the species (Wiegand *et al.*, 2007; Getzin *et al.*, 2008). This heterogeneity may be related to a patchy distribution of limiting resources (*e.g.*, water or soil nutrients) or to periglacial process involving snow cover and duration which would affect the density and distribution of the species (Pescador *et al.*, 2014). Environmental heterogeneity can mask the dispersal and demographic behaviour of a given species (North & Ovaskainen, 2007; Getzin *et al.*, 2008; Chacón-Labella *et al.*, 2014), and can introduce biases in the detection of the pattern of small-scale clustering (*i.e.*, “second-order effects”, *sensu* Wiegand & Moloney, 2004). However, by fitting the HPC model we accounted for the effect of heterogeneity and revealed an additional small-scale aggregation on both stage classes in all populations.

Small-scale aggregation of recruits and adults

The limited dispersal capacity of *S. ciliata* (Lara-Romero *et al.*, 2014) may explain the spatial aggregation of recruits by generating clusters of seeds around mother plants (Howe, 1989; Seidler & Plotkin, 2006). However, seed recruitment not only depends on primary seed dispersal, but is also affected by multiple biotic and abiotic factors controlling seed germination and further establishment of seedlings (Schupp & Fuentes, 1995; Shimono & Kudo, 2003; Pías *et al.*, 2014

2014). We expected *S. ciliata* adults to facilitate germination and early establishment of recruits generating an aggregated pattern among both stages (Lara-Romero *et al.*, 2014). However we found no spatial relation between adults and recruits in most of our populations suggesting that facilitation of conspecific adult neighbours on seedling emergence and survival did not contribute substantially to the spatial clustering of recruits in our study species. In contrast we found an additional aggregation of recruits independently of adults, matching a positive density dependent germination pattern (Antonovics, 1980 #429}. Previous works have related positive density-dependent germination to the scarcity and irregular distribution of suitable sites for germination and establishment (*i.e.*, safe-sites). This leads to high-density clumps of seedlings, but at the same time trigger high mortality due to competition among recruits, thus resulting in a thinning process (Antonovics & Levin, 1980; Stoll & Prati, 2001; Turnbull *et al.*, 2004). Our overall results match this pattern: the clumping patterns generated by seed dispersal seemed to be further reinforced by the heterogeneous distribution of safe-sites (*i.e.*, small gaps of bare soil with suitable conditions for seedling establishment and survival). However, they were followed by a subsequent process of seedling mortality in agreement with the decline in clustering from seedlings to adults. Seedling recruitment into the adult class was probably restricted by competitive exclusion as established seedlings are resource-limited (Gimenez-Benavides *et al.*, 2007; Gimenez-Benavides *et al.*, 2011) and may therefore exhibit lower survival rates in areas of high conspecific density (Antonovics & Levin, 1980; Comita & Hubbell, 2009). This is a common feature in Mediterranean mountains where plant recruitment is extremely limited by high seedling mortality during summer drought (Castro *et al.*, 2005; Gimenez-Benavides *et al.*, 2007).

This general pattern was not fully met in Najarra as our results showed the existence of spatial segregation between adults and recruits. This is in agreement with our previous estimates of greater effective seed dispersal in this population (Lara-Romero *et al.*, 2014). We suspect that this over-dispersed spatial distribution of recruits in Najarra may be due to the greater herbivore-mammal associated

perturbation detected in this population (*e.g.*, herbivory, soil nutrients modifications and trample) (Lara-Romero & Iriondo personal obs.). We speculate that this perturbation may modify the main biotic and abiotic factors controlling recruit-adult relationships which in all the other four populations studied were species-specific dispersal traits, the occupancy of heterogeneously distributed safe-sites and further thinning due to mortality at the recruit-adult transition.

Density dependence effects of the conspecific neighbourhood

Despite the thinning process, the aggregated spatial structure induced by seedling recruitment persisted into the adult class, as the HPC model characterized the empirical pattern of adults with high fidelity in all populations. Previous works assume that positive spatial associations detected between plants at a certain scale are the result of facilitative interactions (McIntire & Fajardo, 2009; Soliveres & Maestre, 2014). In our study, adult reproductive performance measured as the number of flowerheads produced by each adult plant was negatively related to conspecific neighbours density. This result provides empirical evidence for negative density-dependence despite plant aggregation, thereby pointing-out the lack of a simple direct relation between the presence of spatial structure and the sign of plant-plant interactions (*i.e.*, aggregation and segregation meaning facilitation and competition, respectively) (Schupp & Fuentes, 1995; Plotkin *et al.*, 2000; Turnbull *et al.*, 2004). Indeed, taken together, our findings show that facilitative intraspecific (adult-recruit) interactions did not contribute substantially to the tendency for *S. ciliata* to aggregate. Similarly, the presence of competitive intraspecific (recruit-recruit) interactions was not able to generate a segregated spatial pattern. Thus, this suggests that plant-plant interactions, comparatively to other ecological process such as dispersal, safe-site occupation and seedling mortality, played a less important role in determining the spatial pattern of *S. ciliata*.

Conclusions

The spatial distribution of *S. ciliata* was aggregated at both, the adult and recruit stages. We successfully described such distribution with an HPC process for the five

populations studied that occupy a wide range of environmental variability for *S. ciliata* in the Mediterranean region. Thus, our findings highlight the potential of spatial point pattern analyses to the study of plant population dynamics, especially when combined with additional sources of information such as demographic data. Plant-plant interactions played a minor role in explaining the overall pattern, being other biotic and abiotic factors such as seed dispersal, and the non homogeneous distributions of safe sites more determinant. We found a negative relationship between conspecific neighbour density and plant reproductive performance providing empirical evidence for the lack of a simple and direct relationship between plant population structure and the sign of plant-plant interactions (*i.e.*, competition and facilitation). Therefore, it is recommended to avoid overly simplistic inferences on the information contained in plant spatial patterns and that all relevant ecological factors involved in determining such patterns are considered. This way, the establishment of spurious relationships will be prevented and underlying mechanisms of plant spatial patterns will be more easily understood.

ACKNOWLEDGMENTS

The staff of Parque Nacional de Guadarrama gave their permission to work in the protected natural area. This work was supported by the Spanish Ministry of Science and Innovation (LIMITES: CGL2009-07229). CLR was supported by a F.P.I fellowship (Spanish Ministry of Science and Innovation).

REFERENCES

- Antonovics J, Levin DA (1980) The ecological and genetic consequences of density-dependent regulation in plants. *Annual Review of Ecology and Systematics*, 411-452.
- Baddeley A, Turner R (2005) Spatstat: an R package for analyzing spatial point patterns. *Journal of statistical software* **12**, 1-42.
- Bertness MD, Callaway R (1994) Positive interactions in communities. *Trends in Ecology & Evolution* **9**, 191-193.
- Brown C, Law R, Illian JB, Burslem DFRP (2011) Linking ecological processes with spatial and non-spatial patterns in plant communities. *Journal of Ecology* **99**, 1402-1414.

- Callaway RM, Brooker RW, Choler P, *et al.* (2002) Positive interactions among alpine plants increase with stress. *Nature* **417**, 844-848.
- Castro J, Zamora R, Hodar JA, Gomez JM (2005) Alleviation of summer drought boosts establishment success of *Pinus sylvestris* in a Mediterranean mountain: an experimental approach. *Plant Ecology* **181**, 191-202.
- Comita LS, Hubbell SP (2009) Local neighborhood and species' shade tolerance influence survival in a diverse seedling bank. *Ecology* **90**, 328-334.
- Crawford KM, Whitney KD (2010) Population genetic diversity influences colonization success. *Molecular Ecology* **19**, 1253-1263.
- Chacón-Labella J, De la Cruz M, Vicuña R, Tapia K, Escudero A (2014) Negative density dependence and environmental heterogeneity effects on tree ferns across succession in a tropical montane forest. *Perspectives in Plant Ecology, Evolution and Systematics* **16**, 52-63.
- De la Cruz M (2008) Metodos para analizar datos puntuales. In: *Introducción al Análisis Espacial de Datos en Ecología y Ciencias Ambientales: Métodos y Aplicaciones*. (eds. Maestre FT, Escudero A, Bonet A), pp. 76-127. Universidad Rey Juan Carlos and Caja de Ahorros del Mediterráneo, Mostoles.
- De La Cruz M, Romao RL, Escudero A, Maestre FT (2008) Where do seedlings go? A spatio-temporal analysis of seedling mortality in a semi-arid gypsophyte. *Ecography* **31**, 720-730.
- Diggle P (2003) *Statistical analysis of spatial point patterns*, 2nd edn. Arnold, London.
- Dvorský M, Dolezal J, Kopecký M, *et al.* (2013) Testing the Stress-Gradient Hypothesis at the roof of the world: Effects of the cushion plant *Thylacospermum caespitosum* on species assemblages. *PloS one* **8**, e53514.
- Garcia-Fernandez A, Iriondo JM, Escudero A (2012) Inbreeding at the edge: does inbreeding depression increase under more stressful conditions? *Oikos* **121**, 1435-1445.
- Getzin S, Wiegand T, Wiegand K, He F (2008) Heterogeneity influences spatial patterns and demographics in forest stands. *Journal of Ecology* **96**, 807-820.
- Gimenez-Benavides L, Albert MJ, Iriondo JM, Escudero A (2011) Demographic processes of upward range contraction in a long-lived Mediterranean high mountain plant. *Ecography* **34**, 85-93.
- Gimenez-Benavides L, Escudero A, Iriondo JM (2007) Reproductive limits of a late-flowering high-mountain Mediterranean plant along an elevational climate gradient. *New Phytologist* **173**, 367-382.
- Goreaud F, Pélassier R (2003) Avoiding misinterpretation of biotic interactions with the intertype K_{12} -function: population independence vs. random labelling hypotheses. *Journal of Vegetation Science* **14**, 681-692.
- Howe HF (1989) Scatter-and clump-dispersal and seedling demography: hypothesis and implications. *Oecologia* **79**, 417-426.
- Jacquemyn H, Brys R, Vandepitte K, *et al.* (2007) A spatially explicit analysis of seedling recruitment in the terrestrial orchid *Orchis purpurea*. *New Phytologist* **176**, 448-459.

- Jones FA, Comita LS (2008) Neighbourhood density and genetic relatedness interact to determine fruit set and abortion rates in a continuous tropical tree population. *Proceedings of the Royal Society B: Biological Sciences* **275**, 2759-2767.
- Lara-Romero C, Robledo-Arnuncio JJ, García-Fernández A, Iriondo JM (2014) Assessing Intraspecific Variation in Effective Dispersal Along an Altitudinal Gradient: A Test in Two Mediterranean High-Mountain Plants. *PLOS ONE* **9**, e87189.
- Law R, Illian J, Burslem DFRP, *et al.* (2009) Ecological information from spatial patterns of plants: insights from point process theory. *Journal of Ecology* **97**, 616-628.
- Loosmore NB, Ford ED (2006) Statistical inference using the *G* or *K* point pattern spatial statistics. *Ecology* **87**, 1925-1931.
- Maestre FT, Callaway RM, Valladares F, Lortie CJ (2009) Refining the stress-gradient hypothesis for competition and facilitation in plant communities. *Journal of Ecology* **97**, 199-205.
- Maestre FT, Valladares F, Reynolds JF (2005) Is the change of plant-plant interactions with abiotic stress predictable? A meta-analysis of field results in arid environments. *Journal of Ecology* **93**, 748-757.
- McIntire EJB, Fajardo A (2009) Beyond description: the active and effective way to infer processes from spatial patterns. *Ecology* **90**, 46-56.
- North A, Ovaskainen O (2007) Interactions between dispersal, competition, and landscape heterogeneity. *Oikos* **116**, 1106-1119.
- Pescador DS, Chacón-Labolla J, Cruz M, Escudero A (2014) Maintaining distances with the engineer: patterns of coexistence in plant communities beyond the patch-bare dichotomy. *New Phytologist*, 140-148.
- Pías B, Escribano-Avila G, Virgós E, *et al.* (2014) The colonization of abandoned land by Spanish juniper: Linking biotic and abiotic factors at different spatial scales. *Forest Ecology and Management* **329**, 186-194.
- Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team (2014) nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1. R Foundation for Statistical Computing, Viena.
- Plotkin JB, Potts MD, Leslie N, *et al.* (2000) Species-area curves, spatial aggregation, and habitat specialization in tropical forests. *Journal of Theoretical Biology* **207**, 81-99.
- R Core Team (2014) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria
- Reid AM, Lortie CJ (2010) Cushion plants are foundation species with positive effects extending to higher trophic levels. *Ecosphere* **3**, art96.
- Ripley BD (1976) The second-order analysis of stationary point processes. *Journal of applied probability*, 255-266.
- Schupp EW, Fuentes M (1995) Spatial patterns of seed dispersal and the unification of plant population ecology. *Ecoscience* **2**, 267-275.
- Seidler TG, Plotkin JB (2006) Seed dispersal and spatial pattern in tropical trees. *PLoS biology* **4**, e344.
- Shimono Y, Kudo G (2003) Intraspecific variations in seedling emergence and survival of *Potentilla matsumurae* (Rosaceae) between alpine fellfield and snowbed habitats. *Annals of Botany* **91**, 21-29.

- Soliveres S, Maestre FT (2014) Plant-plant interactions, environmental gradients and plant diversity: A global synthesis of community-level studies. *Perspectives in Plant Ecology, Evolution and Systematics* **16**, 154–163.
- Stoll P, Prati D (2001) Intraspecific aggregation alters competitive interactions in experimental plant communities. *Ecology* **82**, 319-327.
- Stoyan D, Stoyan H (1994) *Fractals, random shapes, and point fields: methods of geometrical statistics*, 1st edn. Wiley, Chichester.
- Theodore T, Bowman W (1997) The influence of interspecific competition on the distribution of an alpine graminoid: evidence for the importance of plant competition in an extreme environment. *OIKOS* **79**, 101-114.
- Turnbull LA, Coomes D, Hector A, Rees M (2004) Seed mass and the competition/colonization trade-off: competitive interactions and spatial patterns in a guild of annual plants. *Journal of Ecology* **92**, 97-109.
- Waagepetersen RP (2007) An estimating function approach to inference for inhomogeneous Neyman-Scott processes. *Biometrics* **63**, 252-258.
- Wiegand T, Moloney K (2004) Rings, circles, and null-models for point pattern analysis in ecology. *Oikos* **104**, 209-229.
- Wiegand T, Gunatilleke S, Gunatilleke N, Okuda T (2007) Analyzing the spatial structure of a Sri Lankan tree species with multiple scales of clustering. *Ecology* **88**, 3088-3102.
- Wiegand T, He F, Hubbell SP (2013) A systematic comparison of summary characteristics for quantifying point patterns in ecology. *Ecography* **36**, 092-103.

Figure S1. Spatial distribution of adults (circles) and seedlings (crosses) in each study plot of *S. ciliata*.




Figure S1. (A) Spatial distribution of adults (circles) and seedlings (crosses) in the Laguna population of *Silene ciliata*. Axes show spatial position in meters.




Figure S1. (B) Spatial distribution of adults (circles) and seedlings (crosses) in the Najarra population of *Silene ciliata*. Axes show spatial position in meters.




Figure S1. (C) Spatial distribution of adults (circles) and seedlings (crosses) in the Nevero population of *Silene ciliata*. Axes show spatial position in meters.




Figure S1. (D) Spatial distribution of adults (circles) and seedlings (crosses) in the Cabezas de Hierro population of *Silene ciliata*. Axes show spatial position in meters.




Figure S1. (E) Spatial distribution of adults (circles) and seedlings (crosses) in the Peñalara population of *Silene ciliata*. Axes show spatial position in meters.

Table S1. Description of *S. ciliata* sampling sites.

Population	UTM Coordinates (x,y)		Altitude m	Pm mm	Tm °C	Ws m/s	Seedlings	Adults
	m		m	mm	°C			
Peñalara	419427.3	4522814.15	2405	82.5 (32, 149)	8.77 (0.2, 14.1)	10.62	469	1330
Cabezas de Hierro	421169.5	4516859.61	2305	85.17 (34, 151)	8.37 (0.2, 13.7)	10.87	243	1303
Nevero	428863.74	4537167.57	2190	71 (30, 121)	9.67 (2.1, 14.6)	8.97	21	215
Najarra	430196.59	4518886.29	2080	72.17 (27, 112)	10.45 (2.8, 15.7)	9	89	646
Laguna	419931.16	4521082.09	1946	77 (31, 134)	10.27 (2.8, 15.2)	7.17	10	149

Geographical coordinates, altitude, climatic variables (mean and range) for the growing season (April-September) and seedling and adult plant density (individuals per 10x10 m plot) in study populations. Pm: monthly precipitation, Tm: monthly temperature, Ws: wind speed at 80 m height.. The estimation of climatic variables is detailed in Lara-Romero *et al.* (2014).

CAPÍTULO 4/CHAPTER 4

Genetic variation in flowering phenology and reproductive performance in a Mediterranean high-mountain specialist.



Carlos Lara-Romero, Raúl García-Camacho, Adrian Escudero, Jose M. Iriondo

Manuscrito aceptado en *Botanical Journal of the Linnean Society*

ABSTRACT

Adaptive responses to past climate change may play an important role in the persistence of high-mountain plants, which are vulnerable to global warming. *Armeria caespitosa* is a high-mountain plant, endemic to the Iberian Central Range. Differences in abiotic environment along the elevational gradient impose the species to two opposing stress gradients concerning water stress and duration of the growth season. Furthermore, the species is found in two interspersed, contrasting microhabitats (rocky outcrops and dry cryophilic grasslands) that have different effects on plants depending of the altitude. As a result of this, the species shows great among-population variation in many reproductive and vegetative traits. We used a common garden approach to assess whether this phenotypic variation has a genetic basis or is the result of plastic responses shaped by heterogeneous environmental conditions. Plants from the high-elevation edge and dry cryophilic grasslands flowered earlier and produced more viable fruits but were smaller. These results confirm that among-population variation in flowering phenology and reproductive performance traits in *A. caespitosa* is partially genetically based. Results also showed that the stronger selection response in favor of early-flowering individuals in the low-edge populations did not correspond with greater proportion of early-flowering individuals. Genetic variability associated with flowering onset may be relevant in coping with the impacts of ongoing global warming.

Keywords: common garden, flowering phenology, reproductive performance, mountain plants, local adaptation, altitudinal gradients.

INTRODUCTION

High-mountain plant species are among the organisms thought to be especially vulnerable to global warming (Thuiller *et al.*, 2005; Nogues-Bravo *et al.*, 2007). This becomes a critical issue considering the exceptionally high warming rates projected for the 21st century (Nogues-Bravo *et al.*, 2007). The main response of high-mountain plants to this temperature increase seems to be upward range shifts tracking their current climatic niche (Gottfried, Pauli & Futschik, 2012; Pauli, Gottfried & Dullinger, 2012). However, this may not be feasible for some species because upslope habitat may not be available due to interspecific competition, unfavorable soil conditions or because other species already inhabit summit areas (Pauli *et al.*, 2003; Körner *et al.*, 2007; Larsen, Brehm & Navarrete, 2011). Furthermore, the seed dispersal capacity of some high-mountain species may not be large enough to allow them to track their climatic niche under rapid climate warming (Lara-Romero *et al.*, 2014). In these cases, plants can prevent local extinction by varying their phenotype by means of phenotypic plasticity and/or genetic change through an adaptive evolutionary process (Davis, Shaw & Etterson, 2005; Jump & Peñuelas, 2005; Miner *et al.*, 2005; Gienapp *et al.*, 2008; Chuine *et al.*, 2010; Lloret *et al.*, 2012).

Phenotypic plasticity allows for a more rapid and flexible adjustment to environmental change than adaptation by natural selection (Sultan, 2000; Miner *et al.*, 2005). However, in the long run, plastic response to extreme environmental changes has its limits (Bradshaw, 1965; Bradshaw & McNeilly, 1991; DeWitt *et al.*, 1998; Jump & Peñuelas, 2005; de Jong, 2005; Gienapp *et al.*, 2008). Adaptive evolution can greatly influence the patterns and rates of response to new climatic conditions (Jump & Peñuelas, 2005; Davis and Shaw, 2001; Davis *et al.*, 2005; Parmesan, 2006; Gienapp *et al.*, 2008) and complement other responses, including plasticity or migration (Davis *et al.*, 2005, Parmesan, 2006).

Environmental-change driven phenotypic responses in mountain plant populations has been reported in several studies (Inouye, Morales & Dodge, 2002;

Dune *et al.*, 2003; Inouye, 2008; Kudo & Hiro, 2006; Giménez-Benavides *et al.*, 2011). Nevertheless knowledge on the evolutionary consequences of such phenotypic shifts is still scarce (Gienapp, 2008; Chuine *et al.*, 2010). A change in a species' environment can impose new or intensified selection regimes on traits that may lead to genetically based evolutionary shifts (Hoffmann & Hercus, 2000; Davis *et al.*, 2005; Parmesan, 2006; Gienapp *et al.*, 2008; Munguía-Rosas *et al.*, 2011). However, a heritable trait under natural selection may be prevented from an adaptive response due to evolutionary constraints such as correlations among traits, lack of sufficient genetic variability, fluctuating selection or deleterious mutations (Linhart & Grant, 1996; García-Ramos and Kirkpatrick, 1997; Hoffmann & Hercus, 2000; Jump & Peñuelas, 2005; Parmesan, 2006; Gienap *et al.*, 2008). Actually, many phenotypic differences observed in natural populations may simply be the result of non-evolutionary variation (Miner *et al.*, 2005; Levin, 2009). Consequently, in order to determine a species' potential to adapt to ongoing climate warming, it is critical to know if phenotypic shifts along environmental gradients have a genetic basis.

Armeria caespitosa (Ortega) Boiss. *in* DC. (Plumbaginaceae) is a high-mountain specialist plant, endemic to the Iberian Central Range. Differences in abiotic environment along the elevational gradient impose the species to two opposing stress gradients concerning water stress and duration of the growth season. Hence, plants at higher altitudes experience more favourable conditions in terms of temperature and water availability but have less time to grow, flower and disperse (Gimenez-Benavides *et al.*, 2007; García-Camacho & Escudero 2009; Gimenez-Benavides *et al.*, 2011, Lara-Romero *et al.*, 2014). Furthermore, the species is found in two interspersed, contrasting microhabitats (rocky outcrops and dry cryophilic high-mountain grasslands) that have different effects on plants depending of the elevation. Overall, nutrient and water content in soil is likely to be higher in grassland than in rocks (Körner 2003). Nevertheless, rocky outcrops allow for higher number of orientations in less space than grasslands enabling plants to reach favorable exposure conditions depending on the altitude: orientations with low insolation in summer at lower altitudes and orientations with greater exposure to

prevailing wind at higher altitudes. The latter allows earlier snowmelt at rocky outcrops than at grasslands (Palacios, de Andrés & Luengo, 2003), which may increase the length of growing season and reproductive success of the species (García-Camacho & Escudero 2009; Giménez-Benavides *et al.*, 2011). Earlier studies have shown that *A. caespitosa* is likely to withstand environmental conditions along these complex gradients varying some ecologically important traits along its distribution range. Hence, differences in reproductive performance and demographic vital rates have been found in populations at each microhabitat and edges of the species elevation range (García-Camacho & Escudero, 2009; García-Camacho, Iriondo & Escudero, 2010; García-Camacho *et al.*, 2012). This reproductive and demographic variation suggests that *A. caespitosa* is able to persist at the low edge of its elevation range through “demographic compensation” (García-Camacho *et al.*, 2012); a mechanism by which a decrease in certain vital rates (*i.e.*, life span and fecundity rates) imposed by stressful environmental conditions is buffered by compensatory changes in other demographic vital rates (*i.e.*, growth rate) due to phenotypic plasticity and/or local adaptation (Doak & Morris, 2010). This species seems to have the potential to respond to early snowmelt in warm years by means of shifts in reproductive phenology (García-Camacho & Escudero, 2009; Giménez-Benavides *et al.*, 2011). This ensures an adequate match between flowering period and optimal environmental conditions, which is essential for reproductive success since the growing season is usually very short due to water shortage derived from high temperatures and lack of precipitation as the summer progresses (Giménez-Benavides, Escudero & Iriondo, 2007; Giménez-Benavides *et al.*, 2011). Moreover, flowering onset in this species has been found to be under selection: plants that bloomed earlier flowered for longer periods and produced more fruits (Giménez-Benavides *et al.*, 2011). However, the stronger selection responses for earlier flowering at the lower elevations (Giménez-Benavides *et al.*, 2011) indicates that there is spatial variation across the different elevations and microhabitats. Nevertheless, it remains unknown whether the selective advantage of early-flowering individuals is great enough to cause the predominance of early-flowering phenotypes in nature (Giménez-Benavides *et al.*, 2011).

In this context, we aimed to assess whether the observed variation in a suite of reproductive and vegetative traits in natural populations has a genetic basis or is the result of plastic responses shaped by heterogeneous environmental conditions. We used a common garden approach to study the variation of such traits in four populations of *A. caespitosa*. Common garden experiments allowed us to control the effect of environmental factors on the expression of the traits of interest, ensuring that any differences found among the considered populations would have a genetic origin (Turesson, 1922; Clausen, Keck & Hiesey, 1940; Kawecki & Ebert, 2004). We hypothesized that part of the phenotypic variability observed in natural environments has a genetic basis and therefore, we would detect significant differences among populations in traits related to demographic and reproductive performance and flowering phenology. Concerning flowering phenology, we asked whether the stronger selection responses for earlier flowering observed at lower elevations are promoting the predominance of early-flowering phenotypes (Giménez-Benavides *et al.*, 2011).

MATERIAL AND METHODS

Study species and source populations

Armeria caespitosa (Ortega) Boiss. *in DC.* is a high-mountain dwarf chamaephytic cushion plant, endemic to the mountains of central Spain (Sierra de Guadarrama, Ayllón and East Gredos). It grows at elevations from 1600 to 2430 m in two interspersed but contrasting microhabitats: crevasses and ledges of granite and gneiss outcrops (hereafter rock microhabitat), and in dry cryophilic *Festuca curvifolia* Lag. *ex Lange* fell fields (hereafter pasture microhabitat). *A. caespitosa* is self-incompatible and pollinated by bees, bumblebees and syrphid flies (García-Camacho, Mendez & Escudero, 2009). Its flowers are grouped in short-scaped flowerheads of 19 ± 8 (mean \pm SD) flowers. Further information on the natural history of *A. caespitosa* is available in García-Camacho and Escudero (2009) and García-Camacho *et al.*, (2009, 2010, 2012).

The study site was located in the Sierra de Guadarrama mountain range (Madrid, Spain). Vegetation composition of these high-mountain systems is detailed in Escudero *et al.*, (2004). We replicated the design in García-Camacho and Escudero (2009) to allow the comparison of previous results on the phenology and reproductive output of the species in its natural environment. Hence, we selected four *A. caespitosa* populations at the high and low edges of the species distribution range (hereafter, high-edge and low-edge populations, respectively), two in the rock microhabitat and two in the pasture microhabitat (Table 1). We avoided isolated or small populations to minimize bias due to confounding factors. Thus, we selected genuine functional populations with more than 300 individuals under similar environmental conditions, such as orientation and shrub cover. These populations could not be replicated because they constitute the edge populations of *A. caespitosa* in the Sierra de Guadarrama. On the Western distribution edge in Sierra de Gredos, *A. caespitosa* hybridizes with a close congener, *Armeria bigerrensis* subsp. *bigerrensis*, whereas on the Eastern distribution edge in Sierra de Ayllón *A. caespitosa* populations are small and scarce. More information on the selected populations can be found in García-Camacho & Escudero (2009).

Seed collection and cultivation of plants

In July 2006, we collected fruits from 100 healthy individuals haphazardly chosen at each site and microhabitat (fruits were collected from only 60 plants on the cliffs of Site 1 due to inaccessibility). Fruits were dissected in the laboratory and all the seeds collected per population were pooled. Plants were grown in the CULTIVE Facility (<http://servicat.escet.urjc.es:8080>) situated at 690 m.a.s.l. in Universidad Rey Juan Carlos, Móstoles (Spain). Mean annual precipitation and temperature measured at the experimental field of the CULTIVE Facility are 494 mm (13 – 51 mm) and 14.4 °C (5.8 - 24.9°C), respectively. In spring 2007, we sowed 200 seeds from each population in rectangular plastic pots (5 cm³ volume approximately) with a commercial substrate enriched with NPK and distributed them haphazardly in a greenhouse bench. We allowed plants to develop without stress caused by temperature, nutrient or water limitations. In April 2010, we

transplanted the plants that had grown into the reproductive stage (188 plants in total) in 1800 cm³ rectangular plastic pots with a commercial substrate enriched with NPK. We randomly distributed the pots in the experimental field of the CULTIVE Facility to allow natural pollination. Once again, plants were cultivated without stress caused by temperature, nutrient or water limitations. Because maternal effects are usually only pronounced at the seedling stage and decrease over time (Mousseau & Fox, 1998), we only collected data after the plants had grown under homogeneous conditions for five years, minimizing bias in results due to maternal effects. Voucher specimens were deposited in the Universidad Rey Juan Carlos Germplasm bank (Supporting Information, Table S1).

Sampling methods

We waited until the flowering season of 2011 to monitor phenology and reproductive performance. At that time we also measured the maximum diameter of each plant both as an estimation of plant size at flowering time and growth rate (as all plants were sown at the same time). We counted the total number of flowers per plant every 7 days throughout the flowering period for a total of 15 censuses. All phenological variables were measured with reference to 15th of February. When fruits were ripe at the end of the flowering period, we collected and counted all the flowerheads from each plant. We dissected ten flowerheads per individual, counted the total number of fruits and viable fruits per flowerhead (fruits with one fully matured viable seed) and obtained the mean values for each variable. Following García-Camacho & Escudero (2009) and Nieto-Feliner, Izuzquiza & Lansac (1996), we considered fruits to be viable when they had enlarged and hardened enclosing calyces with wide intercostal spaces, patent hairs and wide-open scarious involucres, which were resistant to pressure from forceps.

We also calculated four individual-based phenological variables: flowering onset, duration, moment and synchrony. Flowering onset was defined as the number of days elapsed between the 15th February and the first flower on the plant. Duration was estimated as the number of days elapsed between the onset and end of blooming

and moment as the number of days elapsed between the blooming of the first flower in all populations studied and the flowering peak. Finally, synchrony was calculated considering the following function (Albert, Escudero & Iriondo, 2001):

$$S_i = \frac{1}{n-1} \sum_{j=1}^n \frac{a_{ij}}{b_{ij}}$$

where n is the number of plants, a_{ij} is the number of days during which individuals i and j are in flower simultaneously, and b_{ij} is the number of days during which at least one of them is in flower.

Table 1. Ecogeographic characterization of *A. caespitosa* sampling sites.

Sampling site	Elevational range	Habitat	Geographic coordinates	Altitude (m)	Pm (mm)	Tm (°C)
Porrones	Low	Rock	40° 45'N, 3° 56'W	1750	1133 (25 – 130)	8.2 (5.2 – 18.3)
Loma de Cabezas	Low	Pasture	40° 46'N, 3° 56'W	1950	1351 (30 – 153)	5.7 (-0.7 – 15.1)
Cabezas de Hierro	High	Rock	40° 48'N, 3° 55'W	2350	1594 (34 – 193)	4.2 (-2.8 – 13.7)
Cabezas de Hierro	High	Pasture	40° 47'N, 3° 56'W	2300	1594 (34 – 193)	4.2 (-2.8 – 13.7)

Pm: mean annual precipitation, *Tm*: mean annual temperature. Minimum and maximum monthly precipitation and temperatures are provided in brackets. Climatic data obtained from Gonzalo-Jiménez (2011).

Statistical analysis

We used General Linear Models (hereafter GLMs) to test for genetic differences in reproductive phenology. Hence, we fitted models for onset, duration, moment and synchrony. All these models included position in the elevational range (hereafter elevational range) and microhabitat as categorical predictors. As in García-Camacho & Escudero (2009), we considered range edge instead of elevation, because we were especially interested in how plants behave at the local low and high edges of their ranges. Similarly, rock and pasture were included as the two levels of microhabitat. We also included interaction effects between elevational range and microhabitat. In the case of the models fitted for duration, synchrony and moment, we included plant size and onset as predictor continuous variables.

We also used GLMs to assess differences in reproductive output. We fitted models for number of flowerheads, fruits per flowerhead and viable fruits per flowerhead. All these models included elevational range, microhabitat and their interaction effects as predictors. In the case of the model fitted for number of flowerheads, we included plant size and onset as continuous predictor variables, while in models fitted for fruits per flowerhead and viable fruits per flowerhead, the continuous predictor variables were plant size and the two phenological variables onset and synchrony. In addition, we fitted GLMs with elevational range, microhabitat and their interaction to model plant size. We did not include moment and duration in the models as predictor continuous variables because both phenological traits were significantly related to onset (see below).

We fitted all models applying the principle of parsimony by means of a stepwise model simplification process or deletion test (Crawley, 2007). We achieved this by fitting a maximal model (*i.e.*, the model containing all the predictor variables mentioned above) and then removing sequentially non-significant interaction terms and explanatory variables from the maximal model to obtain a final model containing nothing but significant and non-redundant predictor variables (*i.e.*,

minimal adequate model, Crawley, 2007). The continuous predictor variables were entered first in the models to minimize potential effects of plant size and/or phenological variables before calculating categorical predictor's effects. We assumed Poisson error and log link functions for all the response variables except fruits per flowerhead and viable fruits per flowerhead, for which we included a quasi-Poisson error and a log-link function.

To examine the effect of elevational range and microhabitat on the long-term survival of adult plants, we tested for differences in mean survival times using the log-rank χ^2 - test of homogeneity among groups of the Kaplan-Meier product-limit survival analysis (Kleinbaum & Klein, 2005). Data for the survival analysis were collected in May 2010, September 2010, May 2011 and September 2011. All statistical analyses were performed in the open source software package R v 3.0.2 (R Core Team, 2014).

RESULTS

Means, standard deviations and sample size of phenology and reproductive variables for each population are presented in Supporting Information, Table S2.

Models fitted for onset and duration found a significant effect of elevational range and microhabitat (Table 2a, b). Plants from the highest elevation and from the pasture microhabitat flowered earlier and had shorter flower duration, once the negative effect of flower onset on flower duration had been taken into account (Figure 1; Table 2a, b; Table S2). No significant elevational range x microhabitat interactions was found. The remaining phenological variables were not significantly affected by elevational range and microhabitat. However, onset significantly affected flowering moment and duration (Table 2b, c). Hence, plants that flowered earlier had a longer flowering duration and a shorter flowering moment. Plant size did not affect any of the studied phenological variables.

Table 2. Minimal adequate model for reproductive phenology of *Armeria caespitosa* plants grown in the common garden experiment.

Effect	Parameter value	Solution for effects			Deviance Change	
		SE	z value	Pr(> z)	χ^2	Pr(> χ^2)
a) Flowering onset						
Intercept	3.957	0.023	171.81	<0.001	-	-
Range	-	-	-	-	5.02	0.02
Low	0.062	0.026	2.41	<0.01	-	-
Microhabitat	-	-	-	-	15.19	<0.001
Rock	0.099	0.025	3.04	<0.01	-	-
b) Flowering duration						
Intercept	4.712	0.056	83.94	<0.001	-	-
Onset	-0.021	0.001	-20.03	<0.001	67.60	<0.001
Range	-	-	-	-	6.31	0.01
Low	0.084	0.032	2.67	<0.01	-	-
Microhabitat	-	-	-	-	5.60	0.02
Rock	0.075	0.031	2.38	0.02	-	-
c) Flowering moment						
Intercept	4.226	0.047	89.84	<0.001	-	-
Onset	0.002	0.0008	2.75	<0.01	7.61	<0.01

SE: Standard Error. Missing factors were removed during model simplification process based on a deletion test. Missing levels of factors (range: high; microhabitat: pasture) are included in the intercept.

Elevational range and microhabitat had a significant effect on the number of flowerheads (Table 3). Plants from low-edge populations and the pasture microhabitat produced more flowerheads (Figure 2a, Table 3). There was also a significant elevational range x microhabitat interaction due to a greater variation between microhabitats at the low-edge populations (Figure 2a, Table 3). Plant size and onset also had a significant positive and negative effect on number of flowerheads, respectively (Table 3).




Figure 1. Variation in flowering phenology between microhabitats and between elevation edges. Horizontal bars indicate flowering onset, duration and end per elevation edge and microhabitat. Dashed lines represent the range of flowering duration in each study population; solid lines delimited by small vertical bars represent the range of mean individual flowering duration. Circles represent the mean flowering moment.

Table 3. Minimal adequate model for number of flowerheads of *Armeria caespitosa* plants grown in the common garden experiment.

Effect	Solution for effects				Deviance change	
	Parameter value	SE	z value	Pr(> z)	χ^2	Pr(> χ^2)
Intercept	2.84	0.11	25.96	<0.001	-	-
Plant size	0.012	0.001	12.79	<0.001	67.41	<0.001
Onset	-0.007	0.001	-7.09	<0.001	38.35	<0.001
Range	-	-	-	-	67.41	<0.001
Low	0.575	0.036	15.94	<0.001	-	-
Microhabitat					57.94	<0.001
Rock	-0.099	0.046	-2.15	0.03	-	-
Range:Microhabitat	-	-	-	-	10.08	<0.01
Low:Rock	-0.177	0.056	-3.18	<0.01	-	-

SE: Standard Error. Missing factors were removed during model simplification process based on a deletion test. Missing levels of factors (range: high; microhabitat: pasture) are included in the intercept.

Number of flowerheads was positively related to total fruits per flowerhead, whereas non-significant effects of elevational range and microhabitat were found (Figure 2b, Table 4a). On the other hand, elevational range had a significant effect on the number of viable fruits per flowerhead, although a significant elevational range x microhabitat interaction was also found (Table 4b). Consequently, the number of viable fruits per flowerhead was greater at the high-edge populations in the pasture microhabitat than at the low-edge populations but was lower in the rock microhabitat (Figure 2c, Table 4b). Plant size and the two phenological variables onset and synchrony were not significantly related with fruit performance of plants grown in the common garden experiment.




Figure 2. Mean values and standard deviations for the components of reproductive success and plant size of *Armeria caespitosa* in the common garden experiment. Black bars: rock microhabitat; grey bars: pasture microhabitat.

Table 4. Minimal adequate model fitted for fruit performance of *Armeria caespitosa* plants grown in the common garden experiment.

Effect	Parameter value	Solution for effects				Deviance change	
		SE	t value	Pr(> t)	χ^2	Pr(> χ^2)	
a) Total fruits per flowerhead							
Intercept	2.661	0.05	9.80	<0.001	-	-	-
Flowerheads	0.002	0.0007	2.10	0.04	4.31	0.04	
b) Viable fruits per flowerhead							
Intercept	2.040	0.076	26.74	<0.001	-	-	-
Range	-	-	-	-	5.32	0.02	
Low	-0.432	0.110	-3.93	<0.001	-	-	-
Microhabitat	-	-	-	-	0.11	0.74	
Rock	-0.262	0.121	-2.16	0.03	-	-	-
Range:Microhabitat	-	-	-	-	11.09	<0.001	
Low:Rock	0.553	0.167	3.32	<0.01	--	--	--

SE: Standard Error. Missing factors were removed during model simplification process based on a deletion test. Missing levels of factors (range: high; microhabitat: pasture) are included on the intercept.

A significant effect of elevational range and microhabitat on plant size was found (Table 5). Plants were found to be smaller in the high-edge populations and in the pasture microhabitat (Figure 2d, Table 5). Kaplan-Meier product-limit survival analysis did not find significant differences in the shape of the survival curves across elevations or microhabitats ($\log\text{-rank } \chi^2 = 5.6$, $df = 3$, $P = 0.13$).

Table 5. Minimal adequate model fitted for plant size of *Armeria caespitosa* plants grown in the common garden experiment.

Effect	Parameter value	Solution for effects			Deviance change	
		SE	z value	Pr(> z)	χ^2	Pr(> χ^2)
Intercept	4.571	0.017	270.56	<0.001	-	-
Range	-	-	-	-	12.63	<0.001
Low	0.072	0.019	3.82	<0.001	-	-
Microhabitat	-	-	-	-	36.53	<0.001
Rock	0.112	0.019	6.06	<0.001	-	-

SE: Standard Error. Missing factors were removed during model simplification process based on a deletion test. Missing levels of factors (range: high; microhabitat: pasture) are included on the intercept.

DISCUSSION

This study shows that in *A. caespitosa*, among-population variation in ecologically relevant traits related to reproductive performance and flowering phenology is genetically based. The inherent genetic variation observed in flowering onset was not consistent with the selective pressures previously identified in natural populations.

Flowering onset

The common garden was established at lower elevation than the natural populations. As a result, plants flowered several weeks earlier in our common garden (early March) than in natural populations (early May: García-Camacho & Escudero, 2009). This is congruent with previous studies that have shown that *A. caespitosa* comes into bloom soon after snowmelt. These results further support the idea that this species is photoperiodically neutral (García-Camacho & Escudero, 2009; Giménez-Benavides *et al.*, 2011) and that a temperature threshold triggers flowering onset. In natural conditions, *A. caespitosa* flowers earlier at the low-

elevation edge, because the critical threshold temperature is reached earlier. However, under common garden conditions, flower onset was earlier in plants from the high-elevation edge. This suggests that the threshold temperature for flowering is lower for individuals from the high-elevation edge.

The genetic variation in flowering onset detected in the common garden experiment between the low and high-elevation edges does not correspond with the selective pressures previously identified in natural populations, where the selection response for earlier flowering was found to be stronger in the low-elevation edge and in the rock microhabitat, but practically negligible in all the other studied populations (Giménez-Benavides *et al.*, 2011). The results reported by Giménez-Benavides *et al.* (2011) are based on selection differentials and gradients. These analyses reveal the strength and direction of the selection acting on each trait just during the selective episodes studied (Lande & Arnold 1983, Arnold & Wade 1984), while the genetically based variations found in a common garden experiment are the outcome of both recent and more ancient processes (Kawecki & Ebert 2004). Consequently, selection responses within a population and genetic differentiation between populations for flowering onset do not need to be causally connected. No selection pressures are expected when the populations have reached their optimal because phenotypes are already differentiated; whereas selection pressures may be strong and different when populations are yet undifferentiated but growing in different environments (Lande & Arnold 1983, Arnold & Wade 1984, Kawecki & Ebert 2004).

Flowering onset at lower temperatures at the high-elevation edge may be linked to the shorter effective growth period on the mountain summits in terms of temperature and the positive correlation between flowering duration and seed production found in natural populations (Giménez-Benavides *et al.* 2011). In this context, earlier flowering at lower temperatures would increase seed production. On the contrary, the threshold temperature for flowering is reached at the low-elevation edge earlier in the year when sudden spring freezing events are more likely. These

episodic spring freezing events may eliminate the advantage of flowering at an earlier date. Giménez-Benavides *et al.* (2011) found directional selection for early flowering genotypes at the lower-elevation edge for two consecutive years, one which was climatically benign and one which was extremely warm and dry. However, episodic late spring freezing events may stabilize selection at a higher threshold temperature when averaged throughout time.

Similarly, the earlier flowering onset found in the pasture microhabitat compared to the rock microhabitat may be related to earlier snowmelt in the latter due to its lower specific heat (Körner, 2003) and greater exposure to prevailing winds (Palacios, de Andrés & Luengo, 2003). In this way, plant populations in the pasture microhabitat would have shorter effective growth periods and need to lower the threshold temperature for flowering onset to maintain the length of the reproductive period and maximize seed production. As a result, the growing season is shorter at the high-elevation edge and in the pasture microhabitat (García-Camacho & Escudero 2009, Giménez-Benavides *et al.*, 2011, Escudero *et al.*, 2012), which may have led to selective pressures towards earlier flowering onset. This is supported by the lower flowering duration showed by plants from these populations in relation to plants from low-elevation edge and rock microhabitat under common garden conditions. In any case, substantial within-population variation in flowering onset was also detected in the common garden experiment (coefficients of variation 0.2 - 0.4). This suggests that relevant genetic variation in flowering onset also exists at the population level.

Flowerheads, viable fruits per flowerhead and plant size

Previous studies on the reproductive success of *A. caespitosa* in the same natural populations used for this study found no significant differences between populations in terms of number of flowerheads (García-Camacho & Escudero, 2008). In contrast, the number of viable fruits per flowerhead was higher at the higher-elevation edge and at the pasture microhabitat in the climatically benign year

2004, but no significant differences were found in the warmer and drier year of 2005 (García-Camacho & Escudero, 2009). Therefore, it is clear that differences in reproductive success greatly depend on environmental conditions. These previous studies also showed that higher-elevation edge and pasture microhabitat tend to harbor smaller plants than lower-elevation edge and rock microhabitat (García-Camacho & Escudero 2008). The sign of the differences found in the natural populations in benign conditions is in accordance with those found in the common garden experiment, suggesting that differences in these traits are also partially due to inherent genetic variation between populations. This variation may be related to the way plants allocate resources to reproduction versus further vegetative growth and survival (Cousens, Dytham & Law, 2008) and may explain the process of “demographic compensation” detected in this species (García-Camacho *et al.*, 2012). These findings agree with those of previous studies that have reported genetic differences in reproductive traits among high-mountain plant populations (*e.g.* Pluess & Stöcklin, 2005; Stöcklin, Kuss & Pluess, 2009; Frei *et al.*, 2011). Once again, relevant variation in these traits was also found at the within-population level in the common garden experiment, suggesting substantial genetic variation within populations (coefficients of variation: 0.4 - 0.7 and 0.4 - 0.5 for number of flowerheads and number of viable fruits per flowerhead, respectively).

Genetic variation in the context of global warming

The inherited capacity to shift flowering onset may be of adaptive value for natural populations under ongoing global warming. In mountains, temperature and precipitation modifications due to climate change seem to primarily affect snow and snowmelt timing (Beniston *et al.*, 1997; Dye, 2002; Laternser & Schneebeli, 2003; García-Romero *et al.*, 2010) and, consequently, the onset and length of the growing season (Inouye *et al.*, 2002; Dune *et al.*, 2003; Inouye, 2008; Kudo & Hiro, 2006; Giménez-Benavides *et al.*, 2011). This effect is particularly critical in Mediterranean mountains because summer precipitation is scarce and soil moisture mainly depends on the supply of snowmelt water, which decreases progressively as the season

advances (Giménez-Benavides *et al.*, 2007). In this context, the genetic variation found within *A. caespitosa* populations may allow them to advance their flowering period in response to warmer conditions, increasing flowering duration and reproductive performance (Giménez-Benavides *et al.*, 2011). This would represent an adaptive response that may enhance persistence under ongoing global warming (Linhart & Grant 1996; Davis, Shaw & Etterson, 2005; Gienapp *et al.*, 2008; Chuine *et al.*, 2010).

Limitations of the common garden experiment

It should be noted that the observed genetic differences among populations may not necessarily be adaptive. In theory, they could also be caused by neutral genetic drift resulting from historical processes irrelevant for the contemporary persistence of populations (Bradshaw, 1965; Linhart & Grant, 1996; Kawecki & Ebert, 2004; Gienapp *et al.*, 2008), or even by cryptic genetic variation (Rutherford, 2000; Ghalambor *et al.*, 2007). Reciprocal transplants or tests of population responses to specific experimental conditions are needed to explore the adaptive significance of the variation in phenotypes among populations (Kawecki & Ebert, 2004).

The environmental conditions experienced in the common garden differed from those in the natural populations because the experimental site was located at a lower elevation than the populations from the low and high-elevation edges. Therefore, the phenotypes expressed in this experiment may not be the same as those that would have been expressed in the natural populations. In any case, the climatic conditions in the common garden are expected to have been equally advantageous for all the studied populations, as they were grown without significant limitations due to temperature conditions or water and mineral nutrient availability.

Conclusions

This study has established the presence of genetic variation in a suite of female reproductive traits of *Armeria caespitosa*. Trait differences between plants of the populations studied found in the common garden experiment are only partially in accordance with those found in the natural populations, suggesting that differences in these traits are partially due to genetic variation between populations. Furthermore, the later flowering onset found in the low-edge populations did not correspond with the stronger selection response in favor of early-flowering individuals previously reported in the low-edge populations. The inherited capacity of this species to shift its flowering phenology coupled with the observed genetic variation within populations indicates that there is adaptive potential to respond to ongoing global warming. Future research should assess the adaptive significance of the documented variation in phenology and life history traits by means of reciprocal transplants in the field.

ACKNOWLEDGEMENTS

The authors thank José Margalet, Carlos Díaz and Javier Morente for help with the field work and Lori De Hond for linguistic assistance. The staff of Parque Natural de las Cumbres, Circo y Lagunas de Peñalara and Parque Regional de la Cuenca Alta del Manzanares gave their permission to work in the protected natural areas. An early draft of the manuscript was written during a short stay of CLR in Jon Ågren's research group (Department of Plant Ecology and Evolution, Evolutionary Biology Centre, Uppsala University, Sweden). This work was supported by the Spanish Ministry of Science and Innovation (MOUNTAINS CGL-2012-38427 and LIMITES: CGL2009-07229), the Spanish Ministry of Economy and Competitiveness (AdAptA: CGL2012-33528), and the Government of Autonomous Region of Madrid (REMEDINAL2). CLR was supported by a F.P.I fellowship (Spanish Ministry of Science and Innovation).

REFERENCES

- Albert MJ, Escudero A, Iriondo JM. 2001. Female reproductive success of narrow endemic *Erodium paularense* in contrasting microhabitats. *Ecology* 82: 1734-1747.
- Arnold SJ, Wade, MJ. 1984. On the measurement of natural and sexual selection: applications. *Evolution* 38: 720-734.
- Beniston M, Stephenson DB, Christensen OB *et al.* 2007. Future extreme events in European climate: an exploration of regional climate model projections. *Climatic Change* 81: 71-95.
- Bradshaw AD. 1965. Evolutionary Significance of Phenotypic Plasticity in Plants. *Advance in Genetics* 13: 115-155.
- Bradshaw AD, McNeilly T. 1991. Evolutionary response to global climatic change. *Annals of Botany* 67: 5-14.
- Clausen J, Keck DD, Hiesey WM. 1948. *Environmental responses of climatic races of Achillea. Experimental studies on the nature of species*. Washington D.C: Carnegie Institute of Washington.
- Chuine I. 2010. Why does phenology drive species distributions? *Philosophical transactions of The Royal Society B* 365: 3149-3160.
- Crawley MJ. 2007. *The R book*. Chichester: John wiley & Sons.
- Cousens R, Dytham C, Law R. 2008. *Dispersal in plants. A population perspective*. Oxford: Oxford University Press.
- Davis MB, Shaw RG. 2001. Range shifts and adaptive responses to Quaternary climate change. *Science* 292: 673-679.
- Davis MB, Shaw RG, Etterson, J.R. 2005. Evolutionary responses to changing climate. *Ecology* 86: 1704-1714.
- DeWitt TJ, Sih A, Wilson DS. 1998. Costs and limits of phenotypic plasticity. *Trends in Ecology & Evolution* 13: 77-81.
- Doak DF, Morris WF. 2010. Demographic compensation and tipping points in climate-induced range shifts. *Nature* 467: 959-962.
- Dunne JA, Harte J, Taylor KJ. 2003. Subalpine meadow flowering phenology responses to climate change: Integrating experimental and gradient methods. *Ecological Monographs* 73: 69-86.
- Dye DG. 2002. Variability and trends in the annual snow cover cycle in Northern Hemisphere land areas, 1972-2000. *Hydrological Processes* 16: 3065-3077.
- Escudero A, García-Camacho R, García Fernández A *et al.* 2012. Vulnerability to global change in Mediterranean high mountains plants - Vulnerabilidad al cambio global en la alta montaña mediterránea. *Ecosistemas* 21: 63-72.
- Escudero A, Giménez-Benavides L, Iriondo JM, Rubio A. 2004. Patch dynamics and islands of fertility in a high mountain Mediterranean community. *Arctic Antarctic and Alpine Research* 36: 518-527.
- Frei ES, Scheepens JF, Armbruster GFJ, Stöcklin J. 2011. Phenotypic differentiation in a common garden reflects the phylogeography of a widespread Alpine plant. *Journal of Ecology* 100: 297-308.

- Garcia-Camacho R, Albert MJ, Escudero A. 2012. Small-scale demographic compensation in a high-mountain endemic: the low edge stands still. *Plant Ecology & Diversity* 25: 37-44.
- Garcia-Camacho R, Escudero A. 2009. Reproduction of an early-flowering Mediterranean mountain narrow endemic (*Armeria caespitosa*) in a contracting mountain island. *Plant Biology* 11: 515-524.
- Garcia-Camacho R, Iriondo JM, Escudero A. 2010. Seedling Dynamics at Elevation Limits: Complex Interactions Beyond Seed and Microsite Limitations. *American Journal of Botany* 97: 1791-1797.
- Garcia-Camacho R, Mendez M, Escudero A. 2009. Pollination Context Effects in the High-Mountain Dimorphic *Armeria Caespitosa* (Plumbaginaceae): Neighborhood Is Something More Than Density. *American Journal of Botany* 96: 1620-1626.
- Garcia-Ramos G, Kirkpatrick M. 1997. Genetic models of adaptation and gene flow in peripheral populations. *Evolution* 51: 21-28.
- Garcia-Romero A, Munoz J, Andres N, Palacios D. 2010. Relationship between climate change and vegetation distribution in the Mediterranean mountains: Manzanares Head Valley, Sierra de Guadarrama (Central Spain). *Climatic Change* 100: 645-666.
- Gienapp P, Teplitsky J, Alho S, Mills A, Merila J. 2008. Climate change and evolution: disentangling environmental and genetic responses. *Molecular Ecology* 17: 167-178.
- Ghalambor CK, McKay JK, Carroll SP, Reznick DN. 2007. Adaptive versus non adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Functional Ecology* 21: 394-407.
- Giménez-Benavides L, Escudero A, Iriondo JM. 2007. Local adaptation enhances seedling recruitment along an elevational gradient in a high mountain Mediterranean plant. *Annals of Botany* 99: 723-734.
- Giménez-Benavides L, Garcia-Camacho R, Iriondo JM, Escudero A. 2011. Selection on flowering time in Mediterranean high-mountain plants under global warming. *Evolutionary Ecology* 25: 777-794.
- Gonzalo-Jiménez J 2011. *Diagnosis fitoclimática de la España peninsular*. Madrid: Organismo Autónomo de Parques Nacionales.
- Gottfried M, Pauli H, Futschik A et al. 2012. Continent-wide response of mountain vegetation to climate change. *Nature Climate Change* 2: 111-115.
- Hoffmann A, Hercus M. 2000. Environmental stress as an evolutionary force. *BioScience* 50: 217-226.
- Inouye DW. 2008. Effects of climate change on phenology, frost damage, and floral abundance of montane wildflowers. *Ecology* 89: 353-362.
- Inouye DW, Morales MA, Dodge GJ. 2002. Variation in timing and abundance of flowering by *Delphinium barbeyi* Huth (Ranunculaceae): the roles of snowpack, frost, and La Niña, in the context of climate change. *Oecologia* 130: 543-550.
- de Jong G. 2005. Evolution of phenotypic plasticity: patterns of plasticity and the emergence of ecotypes. *New Phytologist* 166: 101-118.

- Jump AS, Penuelas J. 2005. Running to stand still: adaptation and the response of plants to rapid climate change. *Ecology Letters* 8: 1010-1020.
- Kawecki TJ, Ebert D. 2004. Conceptual issues in local adaptation. *Ecology Letters* 7: 1225-1241.
- Kleinbaum DG, Klein M. 2005. Kaplan-Meier survival curves and the log-rank test. *Survival analysis*: 55-96.
- Körner C. 2003. *Alpine plant life: functional plant ecology of high mountain ecosystems*. Heidelberg: Springer.
- Körner C. 2007. The use of 'altitude' in ecological research. *Trends in Ecology & Evolution* 22: 569-574.
- Kudo G, Hirao AS. 2006. Habitat-specific responses in the flowering phenology and seed set of alpine plants to climate variation: implications for global-change impacts. *Population Ecology* 48: 49-58.
- Lande R, Arnold SJ. 1983. The Measurement of Selection on Correlated Characters. *Evolution* 37: 1210-1226.
- Lara-Romero C, Robledo-Arnuncio JJ, García-Fernández A, Iriondo JM. 2014. Assessing intraspecific variation in effective dispersal along an altitudinal gradient: a test in two Mediterranean high-mountain plants. *PLoS ONE* 9(1): e87189.
- Larsen TH, Brehm G, Navarrete H et al. 2011. Range shifts and extinctions driven by climate change in the Tropical Andes: Synthesis and directions. In: Herzog SK, Martínez R, Jørgensen PM, Tiessen H, eds. *Climate change and biodiversity in the Tropical Andes*. São José dos Campos: Inter-American Institute for Global Change Research and Scientific Committee on Problems of the Environment, 47-67.
- Laternser M, Schneebeli M. 2003. Long term snow climate trends of the Swiss Alps (1931-1999). *International Journal of Climatology* 23: 733-750.
- Levin DA. 2009. Flowering-time plasticity facilitates niche shifts in adjacent populations. *New Phytologist* 183: 661-666.
- Linhart YB, Grant MC. 1996. Evolutionary significance of local genetic differentiation in plants. *Annual Review of Ecology Evolution and Systematics* 27: 237-277.
- Lloret F, Escudero A, Iriondo JM, Martínez-Vilalta J, Valladares F. 2012. Extreme Climatic Events and Vegetation: The Role of Stabilizing Processes. *Global Change Biology* 18: 797-805.
- Miner BG, Sultan SE, Morgan SG, Padilla DK, Relyea RA. 2005. Ecological consequences of phenotypic plasticity. *Trends in Ecology & Evolution* 20: 687-692.
- Mousseau TA, Fox CW. 1998. The adaptive significance of maternal effects. *Trends in Ecology & Evolution* 13: 403-407.
- Munguía-Rosas MA, Ollerton J, Parra-Tabla V, De-Nova JA. 2011. Meta-analysis of phenotypic selection on flowering phenology suggests that early flowering plants are favoured. *Ecology Letters* 14: 511-521.
- Nieto-Feliner GN, Izuzquiza A, Lansac AR. 1996. Natural and experimental hybridization in Armeria (Plumbaginaceae): *Armeria villosa* subsp *carratricensis*. *Plant Systematics and Evolution* 201: 163-177.

- Nogues-Bravo D, Araujo MB, Errea MP, Martinez-Rica JP. 2007. Exposure of global mountain systems to climate warming during the 21st Century. *Global Environmental Change-Human and Policy Dimensions* 17: 420-428.
- Palacios D, de Andrés, N, Luengo, E. 2003. Distribution and effectiveness of nivation in Mediterranean mountains: Peñalara (Spain). *Geomorphology* 54: 157-178.
- Parmesan C. 2006. Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology Evolution and Systematics*. 637-669.
- Pauli H, Gottfried M, Dirnbock T, Dullinger S, Grabherr G. 2003. Assessing the long term dynamics of endemic plants at summit areas. In: Nagy L, Grabherr G, Körner C, Thompson DBA, eds. *Alpine Biodiversity in Europe*. Berlin: Springer-Verlag, 243-248.
- Pauli H, Gottfried M, Dullinger S *et al.* 2012. Recent plant diversity changes on Europe's mountain summits. *Science* 336: 353-355.
- Pluess AR, Stöcklin J. 2005. The importance of population origin and environment on clonal and sexual reproduction in the alpine plant *Geum reptans*. *Functional Ecology* 19: 228-237.
- R Core Team (2014). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rutherford SL. 2000. From genotype to phenotype: buffering mechanisms and the storage of genetic information. *BioEssays* 22: 1095-1105.
- Sandring S, Riihimäki MA, Savolainen O, Agren J. 2007. Selection on flowering time and floral display in an alpine and a lowland population of *Arabidopsis lyrata*. *Journal of Evolutionary Biology* 20: 558-567.
- Stöcklin J, kuss P, Pluess AR. 2009. Genetic diversity, phenotypic variation and local adaptation in the alpine landscape: case studies with alpine plant species. *Botanica Helvetica* 119: 123-133.
- Sultan SE. 2000. Phenotypic plasticity for plant development, function and life history. *Trends in plant science* 5: 537-542.
- Thuiller W, Lavorel S, Araujo MB, Sykes MT, Prentice IC. 2005. Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences of the United States of America* 102: 8245-8250.
- Turesson G. 1922. The genotypical response of the plant species to the habitat. *Hereditas* 3: 211-350.

SUPPORTING INFORMATION**Table S1.** Voucher information for populations used in the study.

Collection Locality^a	Microhabitat^b	Voucher specimen accesión no.^c	Geographic coordinates
Cuerda de los Porrones, Madrid	Rock	E0265	40° 45'N, 3° 56'W
Loma de Cabezas, Madrid	Pasture	E0268	40° 46'N, 3° 56'W
Cabezas de Hierro Menor, Madrid	Rock	E0270	40° 48'N, 3° 55'W
Cabezas de Hierro Menor, Madrid	Pasture	E0270	40° 47'N, 3° 56'W

^aLocality and Spanish province.

^b Microhabitat in which the specimen was collected: rock (rocky outcrops), pasture (dry cryophilic high-mountain grasslands).

^c Vouchers deposited at Universidad Rey Juan Carlos, Departamento de Biología y Geología, Germplasm bank. (<http://www.gbif.org/dataset/59bf2c83-1e3c-40c8-9437-39ce3d3d462c>).

Table S2. Reproductive output, size, and phenological traits, of *Armeria caespitosa* in the common garden experiment. All the variables are expressed as mean \pm sd.

Variable	Low		High	
	Rock (n=26)	Pasture (n= 37)	Rock (n= 22)	Pasture (n=26)
Size (mm)	118.58 \pm 13.91	102.19 \pm 14.26	105.36 \pm 13.23	99.08 \pm 12.15
Onset (days)	49 \pm 10	44 \pm 13	47 \pm 15	40 \pm 14
Duration (days)	38 \pm 9	38 \pm 12	37 \pm 15	40 \pm 14
Moment (days)	69 \pm 3	65 \pm 5	66 \pm 4	63 \pm 5
Synchrony	0.60 \pm 0.10	0.60 \pm 0.08	0.58 \pm 0.10	0.58 \pm 0.09
Flowerheads	68.42 \pm 30.33	75.76 \pm 31.6	39.5 \pm 25.1	42.31 \pm 30.76
Total fruits per flowerhead	15.16 \pm 3.20	15.69 \pm 3.84	15.76 \pm 4.11	16.10 \pm 5.18
Viable fruits per flowerhead	6.68 \pm 2.43	4.99 \pm 2.38	5.92 \pm 2.14	7.69 \pm 3.73

CAPÍTULO 5/CHAPTER 5

Direct and indirect effects of shrub encroachment on alpine grasslands by altering the assemblage of pollinators



Carlos Lara-Romero, Cristina García, Javier Morente-López, Jose M. Iriondo

Manuscrito inédito

ABSTRACT

Mutualistic interactions structure ecological communities and they are strongly influenced by the combined effect of different drivers of global change. Land-use changes and global warming have elicited a fast shrub encroachment in alpine grasslands during the latest decades, with potential detrimental outcomes for native alpine forbs. Based on the Niche Variation hypothesis (NVH) and the Functional Biodiversity hypothesis (FBH), we asked whether the increase in nutritional resources, as a consequence of the shrub expansion, modifies pollinator's niche breadth and species richness and whether these changes affect plant-plant interactions. For this purpose, we compared quantitative plant-pollinator visitation networks assemblages at replicated plots located at two types of habitats in dry cryophilic grasslands of Sierra de Guadarrama (Spain): *i*) encroached pastures (EP); and *ii*) pastures dominated by forb species where shrub species are absent (PA). As predicted by the FBH, pollinator richness increased at EP, but their niche breadth did not vary, contrary to the NVH's expectation. From the perspective of plant species, shrub expansion reduced the number of visits received per flowering plant in forb species. Additionally, shrubs had more interactions with pollinators and received more visits per flowering plant than forbs at EP, which was in agreement with their significantly higher linkage and strength. This study reveals that moderate levels of shrub encroachment affect the pollination patterns of forb species in alpine grasslands as the diversity of pollinators increased (direct effect) and the plant-plant competition for shared pollinators strengthened (indirect effects). These findings point out the need of an integrative approach to study the cascading effects of global change drivers on species interactions and their impact on the structure and functioning of threatened ecological communities.

Key words: competition, functional-biodiversity hypothesis, specialization, shared pollinators, land-use change, plant-pollinator interactions, niche variation hypothesis.

INTRODUCTION

Global change drivers cause species extinction, modify species distributions and promote the emergence of novel communities, ultimately affecting the state of ecosystems and their associated services (Tylianakis *et al.*, 2008; Algar *et al.*, 2009). High-mountain ecosystems are particularly responsive to the action of global change drivers (Wookey *et al.*, 2009; Pauli *et al.*, 2012). Climate warming and extensification of traditional land management have shifted the distribution ranges of plant species and the composition of plant communities worldwide (Pauli *et al.*, 2012; Brandt *et al.*, 2013). In many cases, these alterations have caused an increase in density, cover and biomass of native woody plants, to the detriment of herbaceous flowering plants (Sanz-Elorza *et al.*, 2003; Wookey *et al.*, 2009; Brandt *et al.*, 2013). Numerous studies document the effect of shrub encroachment on the abiotic environment (*e.g.*, soil erosion and nutrient cycling), plant-plant interactions and vegetation composition (Reviewed in Wookey *et al.*, 2009). By contrast, we lack empirical data showing how shrub encroachment modifies plant-animal interactions (*e.g.*, herbivory, parasitism, pollination: Tylianakis *et al.*, 2008; Ferreira *et al.*, 2013). This is indeed a relevant issue that needs to be addressed as species interactions are among the most important forces structuring ecological communities (Jordano *et al.*, 2003; Tylianakis *et al.*, 2010; Svenning *et al.*, 2014).

Harsh environmental conditions of high-mountain ecosystems limit pollinator richness and pollination services for plant reproduction (Inouye & Pyke, 1988; Kearns & Inouye, 1994; Dupont *et al.*, 2003). In this context of limiting resources, shrub encroachment implies an augment of the nutritional resources for pollinators (Ebeling *et al.*, 2008; Blüthgen & Klein, 2011). According to the biodiversity → (functional) niche hypothesis (Blüthgen & Klein, 2011) (FBH), higher diversity of functionally complementary plants (the producer trophic level) contributes to better sustain pollinators' requirements (the consumer trophic level). This implies a positive feedback of biodiversity at both trophic levels in which more flowering plant species would tend to harbour higher diversity of pollinator species at encroached areas, which in turn might increase pollination success for local plants

(Ebeling *et al.*, 2008; Blüthgen & Klein, 2011). Furthermore, a wider availability of trophic resources results in a reduction of specific niche's breadths of the consumers, according to the “Niche Variation hypothesis” (NVH) (Van Valen, 1965). Consequently, the specific niche breadths of each pollinator would narrow at the same time that species richness augments causing the expansion of the overall niche breadth of the pollinator's community (Bolnick *et al.*, 2007; Ebeling *et al.*, 2011). Additionally, indirect plant-plant competitive interactions for shared pollinators might arise between shrubs and co-flowering forb species. Empirical studies addressing the consequences of shifts in plant communities show mixed results. On one hand, higher pollinator diversity in encroached pastures may increase visitation rates to forbs increasing their reproductive output (Johnson *et al.*, 2003; Ghazoul, 2006). On the other hand, pollinators may shift their preference from forbs to expanding shrubs in encroached areas, thus leading to decreased visitation to forb plants that may reduce their seed set (Revised in Morales & Traveset, 2009). The strength of the plant-plant competition for shared pollinators depends on their flowering phenology and phenotypic similarity in floral resources (Jordano *et al.*, 2003; Morales & Traveset, 2009). Shrub species tend to have less accessible flowers, and therefore, competition for pollinators between pasture and shrub flowering plants tend to involve a subset of pollinators that have access to shrub flowers, such as bees and bumblebees (Fenster *et al.*, 2004; Fontaine *et al.*, 2006; Weiner *et al.*, 2014).

Although previous experimental research has shown that diet breadth is a flexible trait of pollinators affected partially by behavioural adaptation to resource availability (Fontaine *et al.*, 2006; Fontaine *et al.*, 2008), variation in diet breadth and its effects at a larger community scale remains largely unexplored (Fontaine *et al.*, 2008). The above-described community-wide patterns in plant-pollinator and plant-plant interactions can be explored through the application of a species interaction network approach (Jordano, 1987). Interaction networks summarize the diversity of interactions among species in a community and they are widely used to explore the effect of changes in species composition and interactions on ecological

communities (*e.g.*, Lopezaraiza-Mikel *et al.*, 2007; Aizen *et al.*, 2008; Vilá *et al.*, 2009). Topological properties of these networks inform on community organization and they are instrumental to predict community dynamics in response to ecosystem disturbance (Jordano *et al.*, 2003; Blüthgen, 2010; Weiner *et al.*, 2014).

In this study, we tested two well-established hypotheses on the relationship between ecological heterogeneity and niche breadth to investigate the cascading effects of shrub encroachment on alpine plant-pollinator networks. The study was carried out in orophyllous cryophilic pastures of Sierra de Guadarrama, a mountain range located in central Spain. In these mountains, the abandonment of traditional pastoral activities coupled with an increase in winter temperatures has caused an expansion of native woody plants in cryophilic grasslands to the detriment of herbaceous flowering plants (Sanz-Elorza *et al.*, 2003; Garcia-Romero *et al.*, 2010). Two dominant leguminous species of the shrub belt have showy floral displays and co-flower with the pasture community (Castroviejo *et al.*, 1990; Escudero *et al.*, 2004). Based on NVH and FBH, we expected that shrub encroachment would increase pollinator's species richness, and by doing so, pollinator's niche breadth would narrow, ultimately affecting plant-plant interactions through shared pollinators. Specifically, we asked:

- (1) Does higher diversity at the producer trophic level (plants) result in an increased diversity at the consumers (pollinators) trophic level, as predicted by FBH?
- (2) Does shrub encroachment reduce the specific niche breadth of the pollinators, according to the NVH?
- (3) Do shared-pollinators mediate facilitative or competitive indirect interactions between shrubs and forbs?

MATERIAL AND METHODS

Study system

The present study was carried out in the orophyllous cryophilic pastures of Sierra de Guadarrama, a mountain range located in the Central System in Spain ($40^{\circ} 50'N$, $3^{\circ} 57'W$; Figure 1). Dry cryophilic pastures occur in the higher summits above the tree line between 2000 and 2430 m.a.s.l, characterized by a Mediterranean-Alpine-type climate (Escudero *et al.*, 2004). The dominant vegetation is an extremely short pasture (rarely exceeding 3 cm height) dominated by graminoid *Festuca curvifolia* Lag. ex Lange with several creeping chamaephytes and caespitose grasses (hereafter forb species). Timberline is located between 1900 and 2100 m and is dominated by stunted pines (*Pinus sylvestris*), which appear interspersed in a shrub matrix characterized by *Cytisus oromediterraneus* (Fabaceae) Rivas Mart. *et al.*, *Adenocarpus hispanicus* (Fabaceae) (Lam. DC.) and *Juniperus communis* subsp. *alpina* (Cupressaceae) (Suter) Čelak. The first two shrubs (hereafter shrub species) co-flower with forb species. These species produce showy resource-rich flowers with higher reward levels for pollinators compared to the small and resource-poor flowers produced by forb species (Castroviejo *et al.*, 1990).

Sampling design and field survey

Study sites

We replicated the study in two localities separated by a distance of about 17 kilometres (Peñalara and Nevero peaks, Figure 1). In each locality, we selected two sampling sites; a pasture site dominated by forb species where shrub species are absent (hereafter, PA) and one located in an encroached pasture (hereafter, EP) with ~25% of their area covered by shrub species. We selected EP sites with moderate level of shrub encroachment to avoid comparing communities in which forb species have already been displaced. Sampling sites were 700 and 900 m apart from each other in Peñalara and Nevero, respectively. Previous studies show that this distance allows spatially independent pollinator communities (e.g., Aizen *et al.*, 2008;

Kaiser-Bunbury *et al.*, 2009). Differences in altitude of sampling sites within localities and between habitat types (*i.e.*, EP vs. PA) were less than 100 m (Figure 1). All sampling sites had an NW orientation and similar plant communities (74% of forb species occurred in both habitat types, Table S1).




Figure 1. Location of the sampling sites in Sierra de Guadarrama (Spain). Darker-shaded areas show higher altitude. Circles denote pasture habitat (PA) whereas triangles designate encroached pastures (PE). Upper left map indicates the location of Sierra de Guadarrama in the Iberian Peninsula. Photographs on the right side show PA and PE sites in Nevero to illustrate the differences between habitat types. Sampling plots were located at 2264 m (Peñalara-PA), 2214 m (Peñalara-PE), 2202 m (Nevero-PA) and 2126 m (Nevero-PE) above sea level.

Monitoring of plant-pollinator interactions

We collected data throughout the flowering period (Figure S1: 13 June to 28 July 2011) for a total of 10 observation days for each sampling site at Nevero and 11 census days for each sampling site at Peñalara. We added a census day to Peñalara in order to balance the number of hours surveyed in both sites (approximately 160 hours per site). Weather determined the number of census days, as warm, dry and light wind conditions are unusual in high-mountain environments (Dupont *et al.*, 2003). Observations were made on sunny days without wind, from 10:00 to 18:00 h when the environmental conditions at these altitudes allow pollination activity (Dupont *et al.*, 2003). To collect visitation data, we established two 60 x 100m sampling plots in each locality (one at PA and another at EP) where we set 10 linear transects across the width of the plots (60m long and 5m wide). We made successive walks through the line transects and we noted the identity and number of contacts between plant and insect species. Using this methodology the complete area of each sampling plot was observed. Transect methods are very effective for monitoring plant-pollinator interactions in environments where time available for observations is limited (Gibson *et al.*, 2011), as in the case of high-mountain environments. We defined a plant-pollinator interaction as the contact of an insect with the reproductive organs of a flower for more than 1 second (Gibson *et al.*, 2011). Thus, flowering visiting insects were regarded as pollinators regardless of the efficacy of the visit. The two sampling sites in each locality were simultaneously surveyed by two work teams. Insects were captured and determined to the lowest taxonomic category possible with the help of experts (see acknowledgements). Voucher specimens were deposited at Rey Juan Carlos University. We grouped the pollinators into functional groups to facilitate the detection of general patterns. We define ‘functional group’ as a set of pollinators that tend to interact with flowers in a similar way (Fenster *et al.*, 2004). Following the methodology employed in Gómez *et al.*, (2008) we used criteria of similarity in size, proboscis length, foraging behaviour and feeding habits rather than taxonomic relationship. We established eleven functional groups (Appendix 1, Table S1). Plant species were determined in the field. For each census

day, the number of flowering plants of each species was recorded in 10 transects (60m long and 2m wide) per sampling site. The sum of all transects was used as an estimation of the total number of flowering plants (hereafter flowering plant abundance).

Data analysis

Effect of shrub encroachment on species assemblages and visitation patterns

We used linear mixed models (LMMs) to test whether shrub encroachment affected flowering plant abundance, the number of plant-pollinator interactions, total number of visits and the number of visits per flowering plant. All models included habitat type (PA and EP) as a fixed factor, and locality and census day nested within locality as random factors to control for potential temporal and spatial non-independence (Zuur *et al.*, 2009). In the case of the models fitted for the number of interactions, total number of visits and the number of visits per flowering plant we included also flowering plant abundance as a continuous predictor variable because it affects visitation patterns (Vázquez, 2005; Blüthgen *et al.*, 2006).

After evaluating the overall effects of shrub encroachment on visitation patterns and flowering plant abundance, we performed more detailed analysis to assess if the presence of shrub species affected pollinators and forbs in a similar way, or whether, on the contrary, it had different effects for certain plant species or groups of pollinators.

From the pollinators' perspective we tested FBH (*i.e.*, whether the total number of pollinator species increased at EP) by using rarefaction curves. Rarefaction methods allow for comparison of datasets standardizing by sampling effort (Gotelli & Entsminger, 2004). We used ECOSIM ver 7.71 (Gotelli & Entsminger, 2004) to compute sample-based rarefaction curves and their 95% confidence interval curves for each habitat type pooling data from all census days and sampling sites. Curves were rescaled by the number of observations (*i.e.*, number of visits) to allow for species richness comparison (Gotelli & Entsminger,

2004). Then, we applied LMM to test whether shrub encroachment affected the number of interactions and total number of visits recorded for each functional group of pollinators. All models included habitat type, functional group, and their interaction terms as fixed factors, and locality and census day nested within locality as random factors, in addition to flowering plant abundance as continuous predictor.

From the plant species point of view, we tested whether shrub encroachment affected the number of interactions, total number of visits, and the number of visits per flowering plant by applying LMMs. These models allowed detecting shifts in pollinator preferences and indirect plant-plant interactions mediated by shared pollinators. We compared whether visitation patterns differed among plant species according to their life form (fixed factor: shrub or forb) in EP. Note that both life forms only co-occur at EP sites. Once we assessed the effect of the life form in shaping pollinators' preference, we estimated the effect of habitat type (fixed factor) by comparing visitation patterns among forb species that are found at PA and EP sampling sites. The model also included life form and the interaction term with habitat type as fixed factors. All models fitted for plant species included locality and census day nested in locality as random factors and flowering plant abundance of each species as continuous predictor.

We assumed Gaussian error for all the LMM models. Response variables were ln-transformed to reach normality. Residuals did not meet the homocedasticity requirement, and therefore, we modelled the variance of the residuals as σ^2 multiplied with the power of the absolute value of the flowering plant abundance (*i.e.*, “power of the covariate” variance structure, Zuur, 2009). We performed all statistical analyses in the open source software R v 3.0.3 using the package “lmer” (Pinheiro *et al.*, 2014).

Topological network metrics

We built quantitative bipartite networks for full-season data for each habitat type (Jordano, 1987). We used visitation frequencies as a surrogate for interaction strength (Vázquez *et al.*, 2005). We defined visitation frequency between a

pollinator i and a plant species j as the total number of visits of i to j multiplied by relative specific flowering plant abundance of j , *i.e.* the total number of visits was weighted by flowering plant abundance. To address the questions posed in this study, we calculated eight niche-based topological metrics, which describe specialization, niche breadth and niche overlap and their implications with interspecific competition within trophic levels (Jordano *et al.*, 2003; Blüthgen *et al.*, 2006; Blüthgen, 2010):

- 1) number of plant species (P);
- 2) number of pollinator species (A);
- 3) total number of interactions (I);
- 4) total number of visits (V);
- 5) Connectance ($C=I/AP$), *i.e.* the fraction of realised interactions in the network (Jordano, 1987);
- 6, 7) Weighted linkage for plants (lw_p) and pollinators (lw_a) (also called weighted generality and vulnerability) as the effective mean number of partners per plant (lw_p), and per pollinator (lw_a) (Bersier *et al.*, 2002). Both indices are, therefore, a measure of the niche breadth;
- 8) Index of network specialization ($H'2$), that quantifies the degree of niche divergence of elements within an entire bipartite network (Blüthgen *et al.*, 2006). It ranges from 0 (low specialisation, high niche overlap) to 1 (high specialisation, low niche overlap); and

We also estimated three species-level metrics for each plant (p) and pollinator (a) species:

- 1) Linkage level (L ; the number of links or partners that each species had in the network);

2) Index of species specialization (d'), expresses levels of specialization of each species based on its discrimination from random selection of partners (Blüthgen *et al.*, 2006; Dormann *et al.*, 2008). It ranges from 0 (no specialisation, or species that interact with their partners proportionally to their availability) to 1 (perfect specialist, or species that disproportionately interact with rare partners); and

3) Species strength (S) is the sum of dependencies of the partners relying on each species (Dormann *et al.*, 2008). It quantifies the importance of each species from the perspective of their partners. This metric is a quantitative extension of the linkage level.

All these metrics were calculated using the R package ‘*bipartite*’ v. 2.0.4 (Dormann *et al.*, 2008). We used unpaired two-sample t -test to check for differences on species-level metrics between habitats, trophic level (*i.e.*, plant and pollinators) and functional group of pollinators and one-sample t -tests to test for differences between plant life forms within EP.

We finally estimated the modularity (M) of the networks which measures the extent to which species interactions are organized into modules (Olesen *et al.*, 2007). Network modules are groups of species more strongly linked to each other than to species in other modules (Olesen *et al.*, 2007; Tylianakis *et al.*, 2010). We used the software NETCARTO to estimate M of the networks employing the algorithm SA (Guimera & Amaral 2005). We assessed the significance level of M running an SA-analysis of 100 random networks with the same species degree distribution as the empirical one (Guimera & Amaral, 2005; Olesen *et al.*, 2007). M increases with increasing link density within modules and decreasing connectedness between different modules (*i.e.*, as increase specialization) (Olesen *et al.*, 2007; Albrecht *et al.*, 2014). Thus, under NVH, we expected a decrease in the connectivity among modules, and therefore a considerable reduction in M at EP in comparison with AP.

RESULTS

Effect of shrub encroachment on species assemblages and visitation patterns

Overall, the number of flowering plants, number of interactions and the total number of visits recorded per census was higher in encroached pastures (EP) than in pastures (PA), while the number of visits per flowering plant was lower (Fig. 2). Flowering plant abundance only influenced the number of visits per flowering plant ($F_{1, 19} = 74.57, p < 0.001; \beta = -0.0012; t = -8.64, p < 0.001$), and was more stable along the flowering season in EP (weekly mean \pm SD: $0.9 \text{ plant/m}^2 \pm 0.5$) than in PA ($0.6 \text{ plant/m}^2 \pm 0.7$).




Figure 2. Number of flowering plants, number of interactions, total number of visits and number of visits per flowering plant measured in pastures (PA) and encroached pastures (EP). In boxplots, thick lines represent the median, boxes the lower and upper quartile, and whiskers above and below the box indicate the 90th and 10th percentiles. LMM results for testing the effects of shrub encroachment are displayed above each graph.

Pollinator richness was higher in EP than in PA in all functional groups (Table S1). As a result, sample-based rarefied estimates of pollinator richness were significantly higher in EP than in PA (Curves rescaled to 2000 visits: 108 species, 95% CI: 102–113; 91 species, 95% CI: 87–94, respectively). The main functional groups of pollinators did not vary between habitat types: butterflies were the predominant pollinators [28 species (23.5%) at EP, 17 species (18.1%) at PA]; followed by hoverflies [13 species at EP (10.9%), 14 species at PA (14.9%)]; bees [13 species (10.9%) at EP, 11 species (11.7%) at PA]; and beetles [12 species (10.1%) at EP, 11 species (11.7%) at PA]. The greater number of species in EP compared to PA resulted in an increased number of interactions and visits recorded for each functional group (Table 1, Table S2: $\beta = 0.42$, $t = 2.39$, $p = 0.02$ for number of interactions; $\beta = 0.78$, $t = 2.45$, $p = 0.01$ for number of visits). There was also a significant effect of the functional group on the visitation patterns observed (Table 1) due to the higher number of interactions and visits by butterflies, hoverflies, small bees and beetles in both habitat types (Table S2, Tukey's Post hoc tests: Table S3). Furthermore, the interaction habitat type x functional group was not significant (Table 1), suggesting that insects from all eleven functional groups responded to the shrub encroachment in a similar manner. Flowering plant abundance did not affect visitation patterns (Table 1).

Life form (*i.e.*, shrub and forb species) affected the number of interactions with pollinators and the number of visits per flowering plant observed in EP (Table 2a). Hence, shrub species had more interactions with pollinators (Table S4: $\beta = 0.53$, $t = 2.43$, $p = 0.02$) and received more visits per flowering plant (Table S4: $\beta = 0.81$, $t = 2.35$, $p = 0.02$) than forb species. Models fitted for forb species that occurred at both habitat types showed that the number of interactions with pollinators and total number of visits recorded for forb species did not differ between PA and EP (Table 2b). There was a significant effect of plant species on these two variables (Table 2b, S5), mainly because *Jasione crispa*, *Jurinea humilis* and *Senecio pyrenaicus* had significantly greater number of interactions and visits in comparison with most of the forb plants as indicated by Tukey's Post-hoc tests

(Table S6). The interaction habitat type x plant species was not significant (Table 2b). In contrast, the number of visits received per flowering plant was significantly lower in EP (Table 2b, Table S5: $\beta = -1.61$, $t = -3.04$, $p < 0.01$). There was also a significant habitat type x plant species interaction for this variable (Table 2b). Although all forb species received less visits per flowering plant in encroached pastures, two of them showed greater negative effect of shrub encroachment, namely *Biscutella intermedia* ($\beta = -2.30$, $t = -2.39$, $p = 0.02$) and *Solidago virgaurea* ($\beta = -1.63$, $t = -1.98$, $p = 0.04$) (Table S4). Flowering plant abundance had a significant effect on all models fitted for plant species (Table 2a,b), which suggests that visitation patterns to plant species in both habitat types were partially explained by this variable. Hence, plant species with greater flowering plant abundance had more interactions with pollinators (Table S4, S5), and therefore, received more visits (Tables S4, S5). However, flowering plant abundance was inversely related to the number of visits per flowering plant (Tables S4, S5).

Table 1. LMM results for testing the effects of habitat type, functional group and flowering plants abundance on the number of interactions and total number of visits recorded for each functional group of pollinators.

	Num. <i>Df</i>	Den. <i>Df</i>	Number of interactions		Number of visits	
			<i>F</i> value	<i>Pr(> F)</i>	<i>F</i> value	<i>Pr(> F)</i>
Intercept	1	419	1109.51	< 0.001	773.74	< 0.001
Habitat type	1	419	20.75	< 0.001	14.16	< 0.001
Functional group	10	419	34.96	< 0.001	25.40	< 0.001
Plants abundance	1	419	0.07	0.78	0.10	0.76
Habitat type: F. group	10	419	1.42	0.17	1.34	0.21

Num. *Df*, numerator degrees of freedom of fixed effects; Den. *Df*, denominator degrees of freedom for fixed effects. Parameter estimates fitted for fixed effects and standard deviations fitted for random effects are provided as Supplementary data (Table S2).

Table 2. LMM results for testing the effects of habitat type, plant species, life form and flowering plant abundance on number of interactions, total number of visits and number of visits per flowering plant recorded for each plant species.

	Num. <i>Df</i>	Den. <i>Df</i>	Number of interactions		Number of visits		Visits per flowering plant	
			<i>F</i> value	<i>Pr(> F)</i>	<i>F</i> value	<i>Pr(> F)</i>	<i>F</i> value	<i>Pr(> F)</i>
a) Species that occurred at encroached pastures								
Intercept	1	115	356.87	< 0.001	423.69	< 0.001	140.69	< 0.001
Life form	1	115	3.91	0.05	1.12	0.29	12.26	< 0.001
Plants abundance	1	115	10.88	< 0.01	19.46	< 0.001	53.28	< 0.001
b) Species that occurred at both habitat types								
Intercept	1	155	580.23	< 0.001	645.22	< 0.001	359.16	< 0.001
Habitat type	1	155	0.019	0.89	0.12	0.73	25.40	< 0.001
Plant species	10	155	4.64	< 0.001	4.70	< 0.001	22.00	< 0.001
Plants abundance	1	155	26.98	< 0.001	37.84	< 0.001	47.71	< 0.001
H. Type:P. Species	10	155	0.52	0.87	0.45	0.92	2.36	0.01

Num. *Df*, numerator degrees of freedom of fixed effects; Den. *Df*, denominator degrees of freedom for fixed effects. Parameter estimates fitted for fixed effects and standard deviations fitted for random effects are provided as Supplementary Data (Table S4, S5).

Topological network metrics

Topological network metrics confirmed that the total number interactions and visits recorded were higher in EP than in PA (Table 3). Four functional groups of pollinators (butterflies, hoverflies, small bees and beetles) accounted for the majority of interactions and visits in both habitat types. They summed up to 63% of interactions and 74% of visits in PA and up to 60% of interactions and 76% of visits in EP. The forb species *Jasione crispa*, *Jurinea humilis* and *Senecio pyrenaicus* were involved in 40% of all links and received 59 % of all visits at PA, while these species together with the shrub species *Cytisus oromediterraneus* and *Adenocarpus hispanicus* were involved in 54% of all links and had 62% of all visits at EP. Nevertheless, most plant species were highly connected, with 10 species (85%) at PA and 12 species (80%) at EP that had at least 10 interactions and 37 visits or more. For further details, the number of interactions and total number of visits recorded for each plant species are shown in Table S1. Regarding network-level metrics, connectance (C) values were similar between habitat types, while the index of network specialization ($H'2$) was slightly lower in EP than in PA (Table 3). Weighted linkage for plants (lw_p) was four times lower than for pollinators (lw_a) showing that plants interact with a high number of pollinator species. Regarding habitat types, lw_p and lw_a in EP increased by 25% and 17%, respectively, compared to PA (Table 3). The plant-pollinator networks of both habitat types were not significantly modular: $M = 0.34$ (mean \pm SD, of 100 randomizations: $M= 0.35 \pm 0.007$; $p > 0.05$) for PA and $M = 0.32$ (0.33 ± 0.006 ; $p > 0.05$) for EP.

Means of all species-level metrics, namely number of links (L), species specialization (d') and species strength (S) estimated for both plant (p) and pollinators (a) were similar between habitat types (t -test: all $p > 0.05$; Table 3). This pattern was constant for all pollinator functional groups (t -test: all $p > 0.05$), but not for plant life forms: linkage ($L = 31.5 \pm 7.5$) and strength ($S = 11.08 \pm 5.7$) of shrub species were higher than those of forb species in EP ($L = 20.86 \pm 4.0$, $t = 2.66$, $p < 0.02$; $S = 6.53 \pm 2.03$, $t = 2.23$, $p < 0.02$). Furthermore, mean L and mean S were

significantly higher in plants than in pollinators at both PA and EP (*t*- test: $p < 0.001$), but non-significant differences were found for mean d' (*t*- test: $p > 0.05$).

Table 3. Topological network metrics for a bipartite plant-pollinator network obtained for pastures (PA) and encroached pastures (EP).

Metrics	PA	EP
Network-level metrics		
Number of plants (P)	15	17
Number of animals (A)	94	119
Number of interactions (I)	268	367
Number of visits (V)	2423	3116
Connectance (C)	0.19	0.18
Weighted plant linkage (lw_p)	8.76	11.57
Weighted pollinator linkage (lw_a)	2.36	2.86
Network specialization ($H'2$)	0.49	0.42
Species level metrics (Mean \pm SE)		
Plant linkage (L_p)	17.87 ± 3.07	22.12 ± 3.68
Pollinator linkage (L_a)	2.82 ± 0.27	3.13 ± 0.26
Plant species specialization (d'_p)	0.33 ± 0.04	0.31 ± 0.04
Pollinator species specialization (d'_a)	0.25 ± 0.02	0.24 ± 0.02
Plant species strength (S_p)	6.33 ± 1.70	7.06 ± 1.94
Pollinator species strength (S_a)	0.16 ± 0.04	0.14 ± 0.03

SE, standard error. Non-significant differences between PA and EP were found in all tested species level metrics ($p > 0.05$)

DISCUSSION

Pollination patterns shifted in encroached pastures (EP) compared to pastures (PA) and our results show distinctive effects of shrub encroachment on plant and pollinator species. Overall, the number of flowering plants, number of interactions and the total number of visits recorded increased in EP. As predicted by the Functional Biodiversity hypothesis (FBH), the integration of shrub species into the network increased richness and visitation rates of all functional groups of pollinators, but their niche breadth did not vary, contrary to the expectation of the Niche Variation hypothesis (NVH). From the perspective of plant species, shrub expansion reduced the number of visits received per flowering plant in forb species. Shrubs had more interactions with pollinators and received more visits per flowering plant than forb species, which was in agreement with their significantly higher linkage and strength. These latest results point to a strengthening of the plant-plant competition for shared pollinators.

Effects of shrub encroachment on pollinator diversity and visitation patterns

The increase in the number of pollinator species (*i.e.*, species richness) in all functional groups is congruent with our expected results based on the FBH that predicts higher pollinator's biodiversity in rich-plant communities (Ebeling *et al.*, 2008; Blüthgen & Klein, 2011). This implies a positive feedback between species richness of both trophic levels: more flowering plant species in EP derives into higher pollinator species richness. Consequently, the overall niche breadth of the pollinator community expanded. Furthermore, based on the NHV we would have expected that the specific niche breadth of pollinators would decrease suggesting an increased trophic specialization. Yet, network metrics showed that shrub encroachment did not change pollinator niche breadth (lw_a, d'_a); rather, the index of network specialization ($H'2$) decreased suggesting a higher generalisation (higher niche overlap) with shrub encroachment. Previous empirical work found no changes in the overall network specialization with increasing flowering plant species richness, but did detect specialization of particular functional groups of pollinators

(Ebeling et al 2011). Yet, means of all species-level metrics (lw_a , d'_a , s_a) estimated for each functional groups of pollinator were similar between habitat types, which reveals that no group of pollinators narrowed their niche's breadths in response to shrub encroachment. Our findings might be partly explained by a temporal partitioning in the flowering period of plant species throughout the growing season. The flowering period of the study species rarely overlapped (Fig S1), and they show short blooming periods (mean \pm SD: 19 days \pm 8.2 at PA; 20 days \pm 9.5 at EP). Contrarily, pollinators tended to be active across the growing season exceeding the blooming period of each plant species (Morente, Lara-Romero & Iriondo, Unpublished results). From the perspective of pollinators, this means that different plant species provided resources at different time lags, and thus several plant species with non-overlapping flowering peaks are required to fulfil their overall foraging demands in alpine grasslands. This phenological complementarity (*sensu* Blüthgen & Klein, 2011) increases the stability of pollinators' community, but it may not entail trophic specialization towards particular plant species, as shown by our results (Benadi *et al.*, 2014).

Plant-plant interactions mediated by shared pollinators

Generalization can improve the stability of a community, but also enhance interspecific competition when resources are limited (Blüthgen & Klein, 2011), as may be the case in our study system. Shrub encroachment increased plant linkage (*i.e.*, L_p and lw_p) while reducing the number of visits per flowering plant of forb species without affecting their specialization (d'_p). These findings suggest the existence of pollinator-mediated competition among plant species mediated by pollinator sharing, which can potentially result in reduced reproductive success of some plant species (Morales & Traveset, 2009).

Previous studies have shown that generalists have the largest potential to compete with other plants because their resources are accessible to a wide range of pollinators for lengthy time lags (Lopezaraiza-Mikel *et al.*, 2007; Morales & Traveset, 2009; Seifan *et al.*, 2014). In EP results show a low degree of

specialization (d'), but high linkage (L) and pollinator visitation, for shrub species, who would act as super-generalists. Further research is needed to evaluate the effective outcome of the integration of shrubs into the network (increased stability *vs.* increased competition). For instance, the evaluation of the presence of shrub pollen grains in the pollen transport network (Lopezaraiza-Mikel *et al.*, 2007) and the joint estimation of the “quantity” (*i.e.*, visit number) and “quality” (*i.e.*, visit efficiency) component of the interaction and the subsequent effect on plant fitness (Fenster *et al.*, 2004; Seifan *et al.*, 2014) would provide very valuable information.

Effects of shrub encroachment on the topology of plant–pollinator network

In spite of the central role of shrub species, the structure of the networks was very robust to the effects of shrub encroachment (*e.g.*, connectance C , modulariry M and $H'2$). This result is congruent with previous findings in the literature where strong shifts on network properties were found only in highly invaded habitats, when invader species had already displaced native ones (*e.g.*, Aizen *et al.*, 2008). The relative invariance of network structure implies that plant-pollinator interaction networks might be less prone to detrimental effects of disturbance than previously thought (Petanidou *et al.*, 2008; Vilá *et al.*, 2009), as may be the case of the plant-pollinator network studied here. Moreover, a strong dependence of pollinators on a narrow set of plant species represents a risk that renders species more vulnerable to co-extinction (Tylianakis *et al.*, 2010; Svenning *et al.*, 2014). In this context, the integration of generalist shrub species would be advantageous because it may increase network resilience after disruptions and the capacity to respond to environmental changes (Elmqvist *et al.*, 2003; Blüthgen & Klein, 2011). Indeed, recent simulated species extinctions models have shown that the integration into the network of species with higher generalization levels but without displacement of native species makes networks more robust against random secondary species extinctions (Albrecht *et al.*, 2014). Notwithstanding the foregoing, this study focused on mountain sites with moderate level of shrub encroachment. The integration of shrub species into the network may represent a risk if shrub cover reaches a threshold value that leads to significant decline in forbs richness (Aizen *et*

al., 2008; Seifan *et al.*, 2014). In the face of increasing shrub encroachment rates in mountain environments worldwide (Wookey *et al.*, 2009), we urgently need further empirical studies to measure the amount of shrub expansion that would disrupt the original properties of plant-pollination networks in alpine grasslands.

Conclusions

This study reveals contrasting effects of moderate levels of shrub encroachment on plant and pollinator species because it increased species richness and visitation rates of pollinators, but it also increased indirect plant-plant interactions mediated by shared pollinators, which can result in reduced reproductive success of key native forbs. In spite of the central role of shrub species, the topological structure of the network was robust to the integration of shrub species, although it slightly increased network generalisation. Results did not support NVH, while FBH explained the direct outcome of shrub encroachment for forb species as the diversity of pollinators increased. These findings advance our understanding of the role of shrub encroachment in high-mountain ecosystems and point out the need of an integrative approach to study the effect of global change drivers on species interactions and associated ecosystem services.

ACKNOWLEDGEMENTS

We are grateful to Silvia Santamaría, Gema Escribano-Ávila and Pedro Quintana-Ascencio for the helpful recommendations and discussions along the work. The taxonomic identification was made by Concepción Ornosa, Miguel Carles-Torla, Xavier Espadaler, José Luis Yela and Margarita Bravo. The staff of Parque Nacional de Guadarrama gave their permission to work in the protected natural areas. We thank Pedro Quintana-Ascencio, Silvia Santamaría, Gema Escribano, Alfredo García, Samuel Prieto, Alberto Teixido, Marta Rodriguez, María José Albert and Miguel Cantero for the field work. This work was supported by the Spanish Ministry of Science and Innovation (LIMITES: CGL2009-07229). CLR and JML were supported by a F.P.I fellowship (Spanish Ministry of Science and Innovation). CG

was supported by the Investigador (IF) FCT program funded by Fundação para a Ciência e a Tecnologia (FCT) (PTDC/BIA-ECS/116521/2010) and co-funded by the European Program COMPETE: FCOMP-01-0124-FEDER-019772.

REFERENCES

- Aizen MA, Morales CL, Morales JM (2008) Invasive mutualists erode native pollination webs. *PLoS biology* **6**, e31.
- Albrecht M, Padrón B, Bartomeus I, Traveset A (2014) Consequences of plant invasions on compartmentalization and species's roles in plant-pollinator networks. *Proceedings of the Royal Society B: Biological Sciences* **281**, 20140773.
- Algar AC, Kharouba HM, Young ER, Kerr JT (2009) Predicting the future of species diversity: macroecological theory, climate change, and direct tests of alternative forecasting methods. *Ecography* **32**, 22-33.
- Benadi G, Hovestadt T, Poethke H-J, Blüthgen N (2014) Specialization and phenological synchrony of plant-pollinator interactions along an altitudinal gradient. *Journal of Animal Ecology* **83**, 639-650.
- Bersier L-F, Banasek-Richter C, Cattin M-F (2002) Quantitative descriptors of food-web matrices. *Ecology* **83**, 2394-2407.
- Blüthgen N (2010) Why network analysis is often disconnected from community ecology: a critique and an ecologist's guide. *Basic and Applied Ecology* **11**, 185-195.
- Blüthgen N, Klein A-M (2011) Functional complementarity and specialisation: the role of biodiversity in plantâ-pollinator interactions. *Basic and Applied Ecology* **12**, 282-291.
- Blüthgen N, Menzel F, Blüthgen N (2006) Measuring specialization in species interaction networks. *BMC ecology* **6**, 9.
- Bolnick DI, Svanbäck R, Araújo M, Persson L (2007) Comparative support for the niche variation hypothesis that more generalized populations also are more heterogeneous. *Proceedings of the National Academy of Sciences* **104**, 10075-10079.
- Brandt JS, Haynes MA, Kuemmerle T, Waller DM, Radeloff VC (2013) Regime shift on the roof of the world: Alpine meadows converting to shrublands in the southern Himalayas. *Biological Conservation* **158**, 116-127.

- Castroviejo S, Laínz M, López González G, et al. (1990) *Flora iberica. Plantas vasculares de la Península Ibérica e Islas Baleares. Vol. II* Real Jardín Botánico. CSIC, Madrid.
- Dormann CF, B. G, Fruend J (2008) Introducing the bipartite Package: Analysing Ecological Networks. *R news* **8**, 8-11.
- Dupont YL, Hansen DM, Olesen JM (2003) Structure of a plant-flower-visitor network in the high-altitude sub-alpine desert of Tenerife, Canary Islands. *Ecography* **26**, 301-310.
- Ebeling A, Klein A-M, Schumacher J, Weisser WW, Tscharntke T (2008) How does plant richness affect pollinator richness and temporal stability of flower visits? *Oikos* **117**, 1808-1815.
- Ebeling A, Klein A-M, Tscharntke T (2011) Plant-flower visitor interaction webs: temporal stability and pollinator specialization increases along an experimental plant diversity gradient. *Basic and Applied Ecology* **12**, 300-309.
- Elmqvist T, Folke C, Nyström M, et al. (2003) Response diversity, ecosystem change, and resilience. *Frontiers in Ecology and the Environment* **1**, 488-494.
- Escudero A, Gimenez-Benavides L, Iriondo JM, Rubio A (2004) Patch dynamics and islands of fertility in a high mountain Mediterranean community. *Arctic Antarctic and Alpine Research* **36**, 518-527.
- Fenster CB, Armbruster WS, Wilson P, Dudash MR, Thomson JD (2004) Pollination syndromes and floral specialization. *Annual Review of Ecology, Evolution, and Systematics* **35**, 375-403.
- Ferreira PA, Boscolo D, Viana BF (2013) What do we know about the effects of landscape changes on plant-pollinator interaction networks? *Ecological Indicators* **31**, 35-40.
- Fontaine C, Collin CL, Dajoz I (2008) Generalist foraging of pollinators: diet expansion at high density. *Journal of Ecology* **96**, 1002-1010.
- Fontaine C, Dajoz I, Meriguet J, Loreau M (2006) Functional diversity of plant-pollinator interaction webs enhances the persistence of plant communities. *PLoS biology* **4**, e1.
- Garcia-Romero A, Munoz J, Andres N, Palacios D (2010) Relationship between climate change and vegetation distribution in the Mediterranean mountains: Manzanares Head Valley, Sierra de Guadarrama (Central Spain). *Climatic Change* **100**, 645-666.
- Ghazoul J (2006) Floral diversity and the facilitation of pollination. *Journal of ecology* **94**, 295-304.

- Gibson RH, Knott B, Eberlein T, Memmott J (2011) Sampling method influences the structure of plant-pollinator networks. *Oikos* **120**, 822-831.
- Gomez JM, Bosch J, Perfectti F, *et al.* (2008) Spatial variation in selection on corolla shape in a generalist plant is promoted by the preference patterns of its local pollinators. *Proceedings of the Royal Society B: Biological Sciences* **275**, 2241-2249.
- Gotelli NJ, Entsminger G (2004) EcoSim: Null models software for ecology. Acquired Intelligence & Kesey-Bear, Jericho.
- Guimera R, Amaral LAN (2005) Functional cartography of complex metabolic networks. *Nature* **433**, 895-900.
- Inouye DW, Pyke GH (1988) Pollination biology in the Snowy Mountains of Australia: comparisons with montane Colorado, USA. *Australian Journal of Ecology* **13**, 191-205.
- Johnson SD, Peter CI, Nilsson LA, Agren J (2003) Pollination success in a deceptive orchid is enhanced by co-occurring rewarding magnet plants. *Ecology* **84**, 2919-2927.
- Jordano P (1987) Patterns of mutualistic interactions in pollination and seed dispersal: connectance, dependence asymmetries, and coevolution. *American naturalist*, 657-677.
- Jordano P, Bascompte J, Olesen JM (2003) Invariant properties in coevolutionary networks of plant-animal interactions. *Ecology letters* **6**, 69-81.
- Kaiser-Bunbury CN, Memmott J, Müller CB (2009) Community structure of pollination webs of Mauritian heathland habitats. *Perspectives in Plant Ecology, Evolution and Systematics* **11**, 241-254.
- Kearns CA, Inouye DW (1994) Fly pollination of *Linum lewisii* (Linaceae). *American journal of botany* **81**, 1091–1095.
- Lopezaraiza-Mikel ME, Hayes RB, Whalley MR, Memmott J (2007) The impact of an alien plant on a native plant-pollinator network: an experimental approach. *Ecology Letters* **10**, 539-550.
- Morales CL, Traveset A (2009) A meta-analysis of impacts of alien vs. native plants on pollinator visitation and reproductive success of co-flowering native plants. *Ecology Letters* **12**, 716-728.
- Olesen JM, Bascompte J, Dupont YL, Jordano P (2007) The modularity of pollination networks. *Proceedings of the National Academy of Sciences* **104**, 19891-19896.
- Pauli H, Gottfried M, Dullinger S, *et al.* (2012) Recent plant diversity changes on Europe's mountain summits. *Science* **336**, 353-355.

- Petanidou T, Kallimanis AS, Tzanopoulos J, Sgardelis SP, Pantis JD (2008) Long-term observation of a pollination network: fluctuation in species and interactions, relative invariance of network structure and implications for estimates of specialization. *Ecology Letters* **11**, 564-575.
- Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team (2014) nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1. R Foundation for Statistical Computing, Viena.
- Sanz-Elorza M, Dana ED, Gonzalez A, Sobrino E (2003) Changes in the High-mountain Vegetation of the Central Iberian Peninsula as a Probable Sign of Global Warming. *Annals of Botany* **92**, 273-280.
- Seifan M, Hoch E-M, Hanoteaux S, Tielbörger K (2014) The outcome of shared pollination services is affected by the density and spatial pattern of an attractive neighbour. *Journal of Ecology* **102**, 953-962.
- Svenning J-C, Gravel D, Holt RD, *et al.* (2014) The influence of interspecific interactions on species range expansion rates. *Ecography*.
- Tylianakis JM, Didham RK, Bascompte J, Wardle DA (2008) Global change and species interactions in terrestrial ecosystems. *Ecology letters* **11**, 1351-1363.
- Tylianakis JM, Laliberté E, Nielsen A, Bascompte J (2010) Conservation of species interaction networks. *Biological conservation* **143**, 2270-2279.
- Van VAlen L (1965) Morphological variation and width of ecological niche. *American Naturalistk*, **99**, 377-399)
- Vázquez DP (2005) Degree distribution in plant-animal mutualistic networks: forbidden links or random interactions? *Oikos* **108**, 421-426.
- Vázquez DP, Morris WF, Jordano P (2005) Interaction frequency as a surrogate for the total effect of animal mutualists on plants. *Ecology Letters* **8**, 1088-1094.
- Vilá M, Bartomeus I, Dietzsch AC, *et al.* (2009) Invasive plant integration into native plant-pollinator networks across Europe. *Proceedings of the Royal Society B: Biological Sciences* **276**, 3887-3893.
- Weiner CN, Werner M, Linsenmair KE, Blüthgen N (2014) Land-use impacts on plant-pollinator networks: interaction strength and specialization predict pollinator declines. *Ecology* **95**, 466-474.
- Wookey PA, Aerts R, Bardgett RD, *et al.* (2009) Ecosystem feedbacks and cascade processes: understanding their role in the responses of Arctic and alpine ecosystems to environmental change. *Global Change Biology* **15**, 1153-1172.

Zuur A, Ieno EN, Walker N, Saveliev AA, Smith GM (2009) *Mixed effects models and extensions in ecology with R* Springer.

APPENDIX 1.

Functional group of pollinators defined following the criteria of similarity in size, proboscis length, foraging behaviour and feeding habits.

- (i) Bumblebees, pollen-and nectar-collecting members of the bee genus *Bombus*;
- (ii) Bees, mostly pollen- and nectar-collecting females measuring 7 mm in body length or larger;
- (iii) Small bees, mostly pollen- and/or nectar-collecting females smaller than 7 mm;
- (iv) Wasps, including aculeate wasps, large parasitic wasps collecting only nectar;
- (v) Bee flies, long-tongued nectar-collecting Bombyliidae;
- (vi) hoverflies, nectar- and pollen-collecting Syrphidae and short-tongued Bombyliidae;
- (vii) Flies, including nectar- and pollen- feeding species mainly belonging to the families tachinidae and muscidae;
- (viii) Small flies, including feeding nectar- and/or pollen-feeding species smaller than 5 mm;
- (ix) Beetles, including species collecting nectar and/or pollen;
- (x) Butterflies (including diurnal moths), all nectar collectors;
- (xi) Others, including nectar-collecting ants, small parasitic wasps, Microlepidoptera (micromoths), bugs and other occasional flower visitors.

SUPPORTING INFORMATION




Fig. S1. Flowering phenology of the study species at pastures (PA) and encroached pastures (EP) from 13 June to 28 July 2011. Lines indicate flowering period and circles represent the flowering peak. Table S1 contains acronyms used for plant species.

Table S1. List of species studied in each habitat type. Acronym, scientific name, family, functional group and number of interactions and total number of visits recorded are indicated for each species.

Acronym	Scientific name	Family	F. Group	Encroached Pastures (EP)		Pastures (PA)	
				Interactions	Visits	Interactions	Visits
adehis	<i>Adenocarpus hispanicus</i> (Lam.) DC.	Fabaceae	Plant	24	255	n.p.	n.p.
armcae	<i>Armeria caespitosa</i> (Gómez Ortega) Boiss.	Plumbaginaceae	Plant	16	33	21	100
bisint	<i>Biscutella intermedia</i> (Boiss. & Reut.) Malag.	Cruciferae	Plant	4	12	8	36
cytoro	<i>Cytisus oromediterraneus</i> Rivas Mart. & al.	Fabaceae	Plant	39	205	n.p.	n.p.
eupwil	<i>Euphrasia willkommii</i> Freyn.	Scrophulariaceae	Plant	2	3	4	23
gagnev	<i>Gagea nevaensis</i> (Boiss.) O. Bolòs	Liliaceae	Plant	n.p.	n.p.	5	8
genlut	<i>Gentiana lutea</i> L.	Gentianaceae	Plant	9	37	n.p.	n.p.
hiease	<i>Pilosella vahlii</i> (Froel.) F.W. Sch. & Sch. Bip. <i>Pilosella argylocoma</i> (Fr.) F.W. Sch. & Sch.	Compositae	Plant	10	31	n.p.	n.p.
hiebla	Bip.	Compositae	Plant	6	14	n.p.	n.p.
hiesp	<i>Lentodon</i> sp. L.	Compositae	Plant	23	139	24	303
jascri	<i>Jasione crispa</i> (Pourr.) Samp.	Campanulaceae	Plant	44	570	31	414
jurhum	<i>Jurinea humilis</i> (Desf.) DC.	Compositae	Plant	47	723	40	691
leualp	<i>Leucanthemopsis alpina</i> (L.) Heywood	Compositae	Plant	28	176	28	189
ransp	<i>Ranunculus</i> sp. L.	Ranunculaceae	Plant	n.p.	n.p.	4	7
sedbre	<i>Sedum brevifolium</i> DC.	Crassulaceae	Plant	24	187	17	155
sedcan	<i>Sedum candollei</i> Raym.	Crassulaceae	Plant	n.p.	n.p.	18	58
senpyr	<i>Senecio pyrenaicus</i> L.	Compositae	Plant	48	377	36	316
silcil	<i>Silene ciliata</i> Pourr.	Caryophyllaceae	Plant	8	88	4	19
solvir	<i>Solidago virgaurea</i> L.	Compositae	Plant	22	155	18	47

Table S1 (Cont.).

Acronim	Scientific name	Family	F. Group	Encroached Pastures (EP)		Pastures (PA)	
				Interactions	Visits	Interactions	Visits
thypra	<i>Thymus praecox</i> Opiz.	Labiatae	Plant	23	108	17	57
bombus3	<i>Bombus</i> sp. (Latrelle, 1802)	Apidae	Bombus	1	2	2	8
bommax	<i>Bombus maxillosus</i> (Klug, 1817)	Apidae	Bombus	3	8	4	167
bomter	<i>Bombus terrestris</i> (Linnaeus, 1758)	Apidae	Bombus	10	168	n.p.	n.p.
andfla	<i>Andrena flavipes</i> (Panzer 1799)	Andrenidae	Bees	6	10	1	4
andlab	<i>Andrena labialis</i> (Kirby, 1802)	Andrenidae	Bees	n.p.	n.p.	1	1
andnit	<i>Andrena nitida</i> (Müller 1776)	Andrenidae	Bees	1	6	1	2
andova	<i>Andrena ovatula</i> (Kirby 1802)	Andrenidae	Bees	5	54	1	14
andpil	<i>Andrena pilipes</i> (Fabricius, 1781)	Andrenidae	Bees	2	5	n.p.	n.p.
andsue	<i>Andrena suerinensis</i> (Friese 1884)	Andrenidae	Bees	3	7	n.p.	n.p.
antret	<i>Anthophora retusa</i> (Linnaeus, 1758)	Anthophoridae	Bees	n.p.	n.p.	1	1
apimel	<i>Apis mellifera</i> (Linnaeus, 1758)	Apidae	Bees	5	17	1	4
dasmor	<i>Dasypoda morotei</i> (Quilis, 1928)	Melittidae	Bees	n.p.	n.p.	2	3
halcon	<i>Halictus confusus</i> (Smith, 1853)	Halictidae	Bees	9	56	6	17
halqua	<i>Halictus quadricinctus</i> (Fabricius, 1776)	Halictidae	Bees	3	8	2	8
halrub	<i>Halictus rubicundus</i> (Christ, 1791)	Halictidae	Bees	3	5	n.p.	n.p.
halsca	<i>Halictus scabiosae</i> (Rossi 1790)	Halictidae	Bees	1	1	2	7
haltet	<i>Halictus tetrazonius</i> (Klug, 1817)	Halictidae	Bees	1	1	n.p.	n.p.
hopmit	<i>Hoplitis mitis</i> (Nylander, 1852)	Megachilidae	Bees	4	18	3	3
hylpra	<i>Hylaeus praenotatus</i> (Förster, 1871)	Colletidae	Bees	1	1	n.p.	n.p.
cercuc	<i>Ceratina cucurbitina</i> (Rossi, 1792)	Anthophoridae	Small Bees	2	2	n.p.	n.p.
lasalb	<i>Lasioglossum albipes</i> (Fabricius, 1781)	Halictidae	Small Bees	4	4	1	1
lascos	<i>Lasioglossum costulatum</i> (Kriechbaumer, 1873)	Halictidae	Small Bees	4	10	3	5

Table S1 (Cont.).

Acronym	Scientific name	Family	F. Group	Encroached Pastures (EP)		Pastures (PA)	
				Interactions	Visits	Interactions	Visits
lasdus	<i>Lasioglossum dusmeti</i> (Blügheten, 1924)	Halictidae	Small Bees	1	1	n.p.	n.p.
lasiog	<i>Lasioglossum leucoprys y cupromicans</i> (Kirby 1802 & Pérez, 1903)	Halictidae	Small Bees	14	378	14	339
lasleu	<i>Lasioglossum leucozonium</i> (Schrank, 1781)	Halictidae	Small Bees	n.p.	n.p.	1	6
lassub	<i>Lasioglossum subfasciatum</i> (Imhoff, 1832)	Halictidae	Small Bees	4	60	1	3
lasxan	<i>Lasioglossum xanthopus</i> (Kirby, 1802)	Halictidae	Small Bees	1	1	n.p.	n.p.
sphsca	<i>Sphecodes scabricollis</i> (Wesmael, 1865)	Halictidae	Small Bees	4	12	3	4
panban	<i>Panurgus banksianus</i> (Panzer 1806)	Panurginae	Small Bees	8	22	2	13
pancal	<i>Panurgus calcaratus</i> (Panzer 1807)	Panurginae	Small Bees	4	15	2	17
bracon	<i>Braconidae</i> (Benoit, 1949)	Braconidae	Wasps	n.p.	n.p.	1	1
cerare	<i>Cerceris Arenaria</i> (Linnaeus, 1758)	Crabronidae	Wasps	4	46	5	24
cerchy	<i>Cerceris rhybyensis</i> (Linnaeus, 1771)	Crabronidae	Wasps	1	1	n.p.	n.p.
ichneu	<i>Ichneumonidae</i>	Ichneumonidae	Wasps	1	1	n.p.	n.p.
odyner	<i>Odynerus</i> sp. (Latreille, 1802)	Eumenidae	Wasps	3	3	n.p.	n.p.
ophyon	<i>Ophyon</i> (Fabricius, 1798)	Ichneumonidae	Wasps	2	2	1	1
philan	<i>Philantus</i> sp. (Fabricius, 1790)	Sphecidae	Wasps	2	8	n.p.	n.p.
podalo	<i>Podalonia tydei</i> & <i>Podalonia</i> sp. (Le Guillou, 1842 & Fernald, 1927)	Sphecidae	Wasps	2	17	n.p.	n.p.
sphex	<i>Sphex</i> sp. (Linnaeus, 1758)	Sphecidae	Wasps	6	43	8	78
tenmes	<i>Tenthredo mesomela</i> (Linnaeus, 1758)	Sphecidae	Wasps	1	1	n.p.	n.p.
chfes	<i>Chrysotoxum festivum</i> (Linnaeus, 1758)	Syrphidae	Hoverflies	3	7	2	3
chinter	<i>Chrysotoxum intermedium</i> (Linnaeus, 1758)	Syrphidae	Hoverflies	2	4	n.p.	n.p.
chsp	<i>Chrysotoxum</i> sp. (Linnaeus, 1758)	Syrphidae	Hoverflies	n.p.	n.p.	3	4

Table S1 (Cont.).

Acronim	Scientific name	Family	F. Group	Encroached Pastures (EP)		Pastures (PA)	
				Interactions	Visits	Interactions	Visits
didint	<i>Didea intermedia</i> (Loew, 1854)	Syrphidae	Hoverflies	2	4	1	2
erissi	<i>Eristalis similis</i> (Fallen, 1817)	Syrphidae	Hoverflies	11	248	9	217
eriste	<i>Eristalis tenax</i> (Linnaeus, 1758)	Syrphidae	Hoverflies	9	185	9	174
eupco	<i>Eupeodes corollae</i> (Fabricius, 1794)	Syrphidae	Hoverflies	12	32	7	57
mercry	<i>Merodon crypticus</i> (Marcos-García, Vujic & Mengual, 2007)	Syrphidae	Hoverflies	1	1	1	1
	<i>Paragus</i> sp. (Latreille, 1804)					n.p.	n.p.
scaeva	<i>Scaeva albomaculata/pyrastri</i> (Macquart, 1842 & Linnaeus, 1758)	Syrphidae	Hoverflies	11	104	10	119
sersil	<i>Sericomyia silentis</i> (Harris, 1776)	Syrphidae	Hoverflies	n.p.	n.p.	1	1
sphscr	<i>Sphaeroforia scripta</i> (Linnaeus, 1757)	Syrphidae	Hoverflies	10	46	3	4
sphsp	<i>Spaherophoria</i> sp. (Linnaeus, 1758)	Syrphidae	Hoverflies	9	18	6	11
syrib	<i>Syrphus ribesii</i> (Linnaeus, 1758)	Syrphidae	Hoverflies	1	3	1	3
syrtor	<i>Syrphus torvus</i> (Osten Sacken, 1875)	Syrphidae	Hoverflies	n.p.	n.p.	1	1
volele	<i>Volucella elegans</i> (Loew, 1862)	Syrphidae	Hoverflies	n.p.	n.p.	1	2
volpel	<i>Volucella pellucens</i> (Loew, 1862)	Syrphidae	Hoverflies	1	2	n.p.	n.p.
convir	<i>Conophorus virescens</i> (Fabricius, 1787)	Bombyliidae	Bee Flies	1	18	2	63
hemham	<i>Hemipenthes hamifera</i> (Loew, 1854)	Bombyliidae	Bee Flies	5	31	1	12
hemmor	<i>Hemipenthes morio</i> (Loew, 1854)	Bombyliidae	Bee Flies	1	2	1	1
phthi	<i>Phthiria</i> sp. (Evenhuis, 1985)	Bombyliidae	Bee Flies	6	13	6	14
sysgra	<i>Systoechus gradatus</i> (Wiedemann in Meigen, 1820)	Bombyliidae	Bee Flies	5	37	4	29
vilhot	<i>Villa hottentotta</i> (Linnaeus, 1758)	Bombyliidae	Bee Flies	3	23	5	11
calli	<i>Calliphoridae</i>	Calliphoridae	Bigflies	n.p.	n.p.	1	1

Table S1 (Cont.).

Acronim	Scientific name	Family	F. Group	Encroached Pastures (EP)		Pastures (PA)	
				Interactions	Visits	Interactions	Visits
dilmin	<i>Dilophus minor</i> (Strobl, 1900)	Bibionidae	Bigflies	n.p.	n.p.	1	1
dysma	<i>Dysmachus</i> sp. (Loew, 1860)	Asilidae	Bigflies	1	3	1	1
musci	Muscidae	Muscidae	Bigflies	3	5	2	4
neocor	<i>Neomyia cornicina</i> (Fabricius, 1781)	Muscidae	Bigflies	1	1	n.p.	n.p.
tachi1	Tachinidae1	Tachinidae	Bigflies	2	2	n.p.	n.p.
tachi2	Tachinidae2	Tachinidae	Bigflies	4	6	2	4
tachi3	Tachinidae3	Tachinidae	Bigflies	3	9	n.p.	n.p.
tachi4	Tachinidae4	Tachinidae	Bigflies	1	2	n.p.	n.p.
tephri	Tephritidae	Tachinidae	Bigflies	1	1	n.p.	n.p.
antho1	<i>Anthomyiidae1</i>	Anthomyiidae	Smallflies	1	1	n.p.	n.p.
antho2	<i>Anthomyiidae2</i>	Anthomyiidae	Smallflies	1	1	1	2
antho3	<i>Anthomyiidae3</i>	Anthomyiidae	Smallflies	n.p.	n.p.	1	9
chloro	Chloropidae	Chloropidae	Smallflies	3	4	n.p.	n.p.
		Phoridae					
dypt1	<i>Megaselia & Meoneura</i> (Rondani, 1856)	&Carnidae	Smallflies	5	9	6	13
myotes	<i>Myopa testacea</i> (Linneaus 1767)	Conopidae	Smallflies	1	2	n.p.	n.p.
phecin	<i>Pherbellia cinerella</i> (Fallén, 1820)	Sciomyzidae	Smallflies	3	7	1	1
salspho	<i>Saltella sphondyliae</i> (Schrank, 1803)	Sepsidae	Smallflies	7	52	6	52
scaster	<i>Scathophaga stercoraria</i> (Linnaeus, 1758)	Scathophagidae	Smallflies	1	3	n.p.	n.p.
sepneo	<i>Sepsis neocynipsea</i> (Melander & Spuler, 1917)	Sepsidae	Smallflies	1	5	1	1
adsger	<i>Adscita geryon</i> (Hübner 1813)	Zygaenidae	Butterflies	n.p.	n.p.	1	2
aglurt	<i>Aglais urticae</i> (Linneo, 1758)	Nymphalidae	Butterflies	5	74	6	39
argagl	<i>Argynnis aglaja</i> (Linnaeus, 1758)	Nymphalidae	Butterflies	1	5	1	2

Table S1 (Cont.).

Acronym	Scientific name	Family	F. Group	Encroached Pastures (EP)		Pastures (PA)	
				Interactions	Visits	Interactions	Visits
argsp	<i>Argynnis</i> sp. (Rafinesque 1815)	Nymphalidae	Butterflies	1	1	1	2
colalf	<i>Colias alfacariensis</i> (Berger, 1948)	Pieridae	Butterflies	1	2	n.p.	n.p.
colcro	<i>Colias crocea</i> (Geoffroy, 1785)	Pieridae	Butterflies	4	30	5	20
crambi	Crambidae (Meyrick, 1890)	Crambidae	Butterflies	n.p.	n.p.	1	2
eremeo	<i>Erebia meolans</i> (Prunner, 1798)	Nymphalidae	Butterflies	5	48	1	1
eretri	<i>Erebia triaria</i> (Prunner, 1799)	Nymphalidae	Butterflies	1	2	n.p.	n.p.
helvir	<i>Heliothis viriplaca</i> (Ochsenheimer, 1816)	Noctuidae	Butterflies	2	6	n.p.	n.p.
	<i>Hipparchia alcyone</i> (Denis & Schiffermüller, 1775)	Nymphalidae	Butterflies	n.p.	n.p.	n.p.	n.p.
hipalc	<i>Hipparchia semele</i> (Linnaeus, 1758)	Nymphalidae	Butterflies	2	3	n.p.	n.p.
inaio	<i>Inachis io</i> (Linnaeus, 1759)	Nymphalidae	Butterflies	1	1	1	1
isslat	<i>Issoria lathonia</i> (Linnaeus, 1758)	Nymphalidae	Butterflies	11	551	9	159
lamboe	<i>Lampides boeticus</i> (Linnaeus, 1767)	Lycaenidae	Butterflies	1	4	2	2
lycalc	<i>Lycaena alciphron</i> (Rottemburg, 1775)	Lycaenidae	Butterflies	2	2	1	2
lycphl	<i>Lycaena phlaeas</i> (Linnaeus, 1761)	Lycaenidae	Butterflies	3	3	n.p.	n.p.
macste	<i>Macroglossum stellatarum</i> (Linnaeus, 1758)	Sphingidae	Butterflies	5	25	5	54
manjur	<i>Maniola jurtina</i> (Linnaeus, 1758)	Nymphalidae	Butterflies	1	1	n.p.	n.p.
meldid	<i>Melitaea didyma</i> (Esper, 1779)	Nymphalidae	Butterflies	2	2	n.p.	n.p.
	<i>Pandoriana pandora</i> (Denis & Schiffermüller, 1775)	Nymphalidae	Butterflies	3	32	4	207
parapo	<i>Parnassius apollo</i> (Linnaeus, 1758)	Papilionidae	Butterflies	3	8	n.p.	n.p.
piebra	<i>Pieris brassicae</i> (Linnaeus, 1758)	Pieridae	Butterflies	2	3	2	2
pienap	<i>Pieris napi</i> (Linnaeus, 1758)	Pieridae	Butterflies	1	2	n.p.	n.p.

Table S1 (Cont.).

Acronym	Scientific name	Family	F. Group	Encroached Pastures (EP)		Pastures (PA)	
				Interactions	Visits	Interactions	Visits
pierap	<i>Pieris rapae</i> (Linnaeus, 1758)	Pieridae	Butterflies	1	1	n.p.	n.p.
pondap	<i>Pontia daplidice</i> (Linnaeus, 1758)	Pieridae	Butterflies	1	1	n.p.	n.p.
pyrali1	Pyralidae1	Pyralidae	Butterflies	1	2	n.p.	n.p.
pyrali2	Pyralidae2	Pyralidae	Butterflies	2	6	3	10
synant	<i>Synanthedon</i> sp. (Hübner 1819)	Sesiidae	Butterflies	2	2	1	1
vanata	<i>Vanessa atalanta</i> (Linnaeus, 1758)	Nymphalidae	Butterflies	1	1	3	47
vancar	<i>Vanessa cardui</i> (Linnaeus, 1758) <i>Acmaeoderella coarctata coarctata</i> (Lucas, 1846)	Nymphalidae	Butterflies	2	46	n.p.	n.p.
acmcoa		Buprestidae	Beetles	3	11	4	6
chryso	Chrysomelidae (Latreille, 1802)	Chrysomelidae	Beetles	1	1	n.p.	n.p.
cocsep	<i>Coccinella septepunctata</i> (Linnaeus, 1758)	Coccinellidae	Beetles	4	15	6	26
curcul1	Curculionidae1	Curculionidae	Beetles	1	3	3	7
curcul2	Curculionidae2	Curculionidae	Beetles	n.p.	n.p.	1	1
enicop	<i>Enicopus</i> sp. (Scopoli, 1763)	Dasytidae	Beetles	7	108	10	162
enipil	<i>Enicopus pilosus</i> (Scopoli, 1763)	Dasytidae	Beetles	n.p.	n.p.	1	1
helper	<i>Heliopathes perrouidi</i> (Mulsan & Rey, 1854)	Tenebrionidae	Beetles	1	1	2	4
lablon	<i>Labidostomis longimana</i> (Linnaeus, 1760)	Chrysomelidae	Beetles	n.p.	n.p.	n.p.	n.p.
lytves	<i>Lytta vesicatoria</i> (Linnaeus, 1758)	Meloidae	Beetles	2	30	n.p.	n.p.
mylvar	<i>Mylabris variabilis</i> (Pallas, 1781)	Meloidae	Beetles	7	105	2	3
nitidu	Nitidulidae (Latreille, 1802)	Nitidulidae	Beetles	3	10	5	14
phalac1	Phalacridae1	Phalacridae	Beetles	2	18	2	7
phalac2	Phalacridae2	Phalacridae	Beetles	3	11	2	2
phyvul	<i>Phyllodecta vulgatissima</i> (Linnaeus, 1758)	Chrysomelidae	Beetles	1	2	n.p.	n.p.

Table S1 (Cont.).

Acronim	Scientific name	Family	F. Group	Encroached Pastures (EP)		Pastures (PA)	
				Interactions	Visits	Interactions	Visits
camlat	<i>Camptopus lateralis</i> (German, 1817)	Alydidae	Others	1	1	n.p.	n.p.
forfus	<i>Formica fusca</i> (Linnaeus 1758)	Formicidae	Others	2	4	2	3
lygaei2	Lygaeidae (Schilling, 1829)	Lygeidae	Others	n.p.	n.p.	2	5
lygeid	Lygeidae (Schilling, 1829)	Lygeidae	Others	n.p.	n.p.	1	1
micbla	Microlepidoptero 1	-	Others	1	2	1	2
micbri	Microlepidoptero 2	-	Others	1	1	n.p.	n.p.
miccam	Microlepidoptero 3	-	Others	1	1	1	1
micman	Microlepidoptero 4	-	Others	n.p.	n.p.	2	2
micpar	Microlepidoptero 5	-	Others	6	32	10	65
micral	Microlepidoptero 6	-	Others	2	5	n.p.	n.p.
mirida	Miridae (Hahn, 1831)	Miridae	Others	n.p.	n.p.	1	1
scioco	<i>Sciocoris sp.</i> (Fallén, 1829)	Pentatomidae	Others	n.p.	n.p.	1	2
spisax	<i>Spilostethus saxatilis</i> (Scopoli, 1763)	Lygeidae	Others	1	1	n.p.	n.p.
tetimp	<i>Tetramorium impurum</i> (Fürster 1850)	Formicidae	Others	n.p.	n.p.	2	4

Table S2. Parameter estimates and standard deviation fitted for fixed and random effects by LMM performed for testing the effects of habitat type, functional group and flowering plants abundance on number of interactions and total number of visits recorded for each functional group.

		Number of interactions			Number of visits		
		Parameter Value	t value	Pr(> t)	Parameter Value	t value	Pr(> t)
Fixed effects							
	Intercept	0.40	3.05	< 0.001	0.85	3.62	< 0.001
Habitat type	Encroached	0.42	2.39	0.02	0.78	2.45	0.01
Functional Group	Big flies	0.04	0.22	0.83	-0.59	-1.87	0.06
	Bees	0.34	1.93	0.06	0.10	0.31	0.76
	Beetles	1.04	5.86	< 0.01	1.20	3.78	< 0.001
	Bee flies	0.08	0.44	0.66	0.04	0.11	0.91
	Butterflies	1.20	6.78	< 0.001	1.50	4.75	< 0.001
	Others	0.47	1.68	0.10	0.36	1.15	0.25
	Smallbees	0.99	5.60	< 0.001	1.53	4.83	< 0.001
	Smallflies	0.49	2.76	0.01	0.37	1.15	0.25
	Hoverflies	1.37	7.72	< 0.001	1.87	5.90	< 0.001
	Wasps	0.28	1.61	0.11	0.31	0.97	0.33
Plant abundance H. Type:F. Group		0.0001	0.27	0.78	0.0001	0.31	0.75
	Encroached:Flies	-0.19	-0.76	0.45	-0.20	-0.44	0.66
	Encroached:Bees	0.15	0.61	0.54	0.03	0.08	0.94
	Encroached:Beetles	-0.25	-1.02	0.31	-0.31	-0.70	0.48
	Encroached:Bee flies	-0.26	-1.06	0.29	-0.54	-1.20	0.23
	Encroached:Butterflies	-0.01	-0.05	0.96	-0.10	-0.21	0.83
	Encroached:Others	-0.62	-2.50	0.01	-1.21	-2.69	0.01
	Encroached:Small bees	-0.15	-0.59	0.56	-0.49	-1.08	0.28
	Encroached:Small flies	-0.34	-1.39	0.17	-0.71	-1.59	0.11
	Encroached:Hoverflies	-0.03	-0.13	0.90	-0.47	-1.04	0.30
	Encroached:Wasps	-0.28	-1.15	0.25	-0.70	-1.56	0.12
Random effects							
Intercept				SD		SD	
	Population			0.18		0.31	
	Residual			0.18		0.31	
				0.64		0.96	

SD, standard deviation. Significant effects ($p < 0.05$) are indicated in bold. Missing levels of factors (Habitat type: Pasture; Functional group: Bumblebees) are included on the intercept. Bumblebees was the functional group with the lowest number of interactions and visits measured, which facilitates the comparison among functional groups.

Table S3. Z ratio values for Tukey's post-hoc tests used to detect differences between functional groups of pollinators in the number of interactions and number of visits recorded. *significant differences after Tukey's adjustment for a family of 11 means.

	Bombus	Bees	Beetles	Big flies	Bee flies	Butterflies	Others	Small bees	Small flies	Hoverflies	Wasps
Number of interactions recorded for each functional group of pollinators											
Bombus	-										
Bees	-3.37*	-									
Beetles	-7.33*	-3.96*	-								
Big flies	0.45	3.82*	7.78*	-							
Bee flies	0.43	3.80*	7.76*	-0.02	-						
Butterflies	-9.61*	-6.25*	-2.29	-10.06*	-10.04*	-					
Others	-1.32	2.05	6.01*	-1.77	-1.75	8.29*	-				
Small bees	-7.38*	-4.02*	-0.06	-7.83*	-7.82*	2.23	-6.07*	-			
Small flies	-2.54	0.83	4.78*	-2.99	-2.97	7.07*	-1.22	4.84*	-		
Hoverflies	-10.87*	-7.50*	-3.55*	-11.32*	-11.30*	-1.26	-9.55*	-3.49*	-8.33*	-	
Wasps	-1.14	2.23	6.19*	-1.59	-1.57	8.47*	0.18	6.24*	1.40	9.73*	-
Number of visits recorded for each functional group of pollinators											
Bombus	-										
Bees	-0.46	-									
Beetles	-4.62*	-4.16*	-								
Big flies	3.12	3.58*	7.74*	-							
Bee flies	1.12	1.58	5.74*	-2.00	-						
Butterflies	-6.47*	-6.01*	-1.85	-9.59*	-7.59*	-					
Others	1.09	1.55	5.71*	-2.03	-0.03	7.56*	-				
Small bees	-5.66*	-5.20*	-1.04	-8.78*	-6.78*	0.81	-6.75*	-			
Small flies	0.01	0.45	4.61*	-3.12	-1.13	6.47*	-1.09	5.66*	-		
Hoverflies	-7.26*	-6.80*	-2.64	-10.38*	-8.38*	-0.79	-8.35*	-1.60	-7.25*	-	
Wasps	0.18	0.64	4.80*	-2.94	-0.94	6.65*	-0.91	5.84*	0.19	7.44*	-

Table S4. Parameter estimates and standard deviation fitted for fixed and random effects by LMM performed to test the effects of life form and flowering plant abundance on number of interactions, total number of visits and number of visits received per flowering plant.

	Number of interactions			Number of visits			Visits per flowering plant		
	Parameter Value	t value	Pr(> t)	Parameter Value	t value	Pr(> t)	Parameter Value	t value	Pr(> t)
Fixed effects									
Intercept		1.21	12.88	<0.001		1.94	14.02	<0.001	
Life form	Shrub	0.53	2.43	0.02		0.54	1.68	0.09	
Plant abundance		0.0009	3.30	<0.01		0.0019	4.41	<0.001	
Random effects									
		SD				SD			SD
Intercept		0.00004				0.00004			0.00014
Population		0.00002				0.00001			0.00001
Residual		0.93977				1.37602			1.62155

SD, standard deviation. Significant effects ($p < 0.05$) are indicated in bold. Missing levels of factors (Life form: Herb) are included on the intercept.

Table S5. Parameter estimates and standard deviation fitted for fixed and random effects by LMM performed to test the effects of habitat type, plant species and flowering plant abundance number of interactions, total number of visits and number of visits received per flowering plant.

		Number of interactions			Number of visits			Visits per flowering plant				
		Parameter	Value	t value	Pr(> t)	Parameter	Value	t value	Pr(> t)	Parameter	Value	
Fixed effects												
Intercept			0.33	0.86	0.39		1.15	2.01	0.04	-2.07	-4.36	<0.001
Habitat type	Encroached		0.14	0.30	0.77		0.15	0.21	0.83	-1.61	-3.04	<0.01
Plant species	Armcae		0.76	1.52	0.13		0.20	0.27	0.78	0.48	0.78	0.44
	Bisint		0.49	0.89	0.37		0.42	0.52	0.61	2.72	3.54	<0.001
	Pilvah		0.60	1.29	0.20		0.85	1.23	0.22	1.10	1.91	0.06
	Jascri		1.00	2.14	0.03		1.34	1.93	0.06	0.83	1.45	0.15
	Jurhum		1.78	3.89	<0.001		2.24	3.32	<0.001	2.15	3.82	<0.01
	Leualp		0.78	1.68	0.09		0.71	1.03	0.30	1.40	2.38	0.02
	Sedbre		0.83	1.80	0.07		1.01	1.47	0.14	0.11	0.19	0.85
	Senpyr		1.57	3.36	<0.001		1.83	2.66	0.01	2.19	3.84	<0.01
	Solvir		1.02	1.95	0.05		0.46	0.59	0.55	2.88	3.98	<0.001
	Thypra		0.95	1.77	0.08		0.91	1.15	0.25	2.81	3.95	<0.001
Plant abundance		0.0009	5.29	<0.001		0.0015	6.01	<0.001	-0.0014	-6.52		<0.001
H. Type:P. species	Encroached:armcae		0.16	0.25	0.80		-0.08	-0.08	0.93	-1.01	-1.27	0.21
	Encroached:bisint		-0.69	-0.95	0.35		-1.00	-0.93	0.35	-2.30	-2.39	0.02
	Encroached:pilvah		0.20	0.33	0.75		-0.02	-0.02	0.98	0.13	0.18	0.86
	Encroached:jascri		-0.18	-0.31	0.76		-0.24	-0.27	0.79	0.56	0.78	0.44
	Encroached:jurhum		-0.52	-0.90	0.37		-0.44	-0.52	0.61	-0.23	-0.32	0.75
	Encroached:leualp		0.13	0.22	0.83		0.12	0.14	0.89	-0.28	-0.38	0.70
	Encroached:sedbre		-0.44	-0.75	0.46		-0.78	-0.88	0.38	-0.15	-0.22	0.83
	Encroached:senpyr		-0.20	-0.34	0.74		-0.44	-0.49	0.63	0.72	0.98	0.33
	Encroached:solvir		-0.05	-0.08	0.94		0.51	0.53	0.60	-1.63	-1.98	0.04
	Encroached:thypra		-0.04	-0.06	0.95		-0.04	-0.04	0.97	0.19	0.21	0.83
Random effects												
Intercept					SD			SD			SD	
Population					0.00005			0.00008			0.15597	
Residual					0.01324			0.35383			0.00191	
					0.98578			1.43571			1.10141	

SD, standard deviation. Significant effects ($p < 0.05$) are indicated in bold. Missing levels of factors (Habitat type: Pasture; Plant species: Silcil) are included on the intercept. Table S1 contains acronyms used for plant species. Silcil was the plant species with the lowest number of interactions and visits measured, which facilitates the comparison among plant species.

Table S6. Z ratio values for Tukey's post-hoc tests used to detect differences between plant species in the number of interactions, total number of visits and number of visits per flowering plant recorded. *significant differences after Tukey's adjustment for a family of 11 means.

	Silcil	Armcae	Bisint	Pilvah	Jascri	Jurhum	Leualp	Sedbre	Senpyr	Solvir	Thypra
Number of interactions recorded for each plant species											
Silcil	-										
Armcae	-2.5	-									
Bisint	-0.39	1.93	-								
Pilvah	-2.32	0.5	-1.7	-							
Jascri	-3	-0.24	-2.32	-0.86	-						
Jurhum	-5.19*	-2.37	-4.31*	-3.41*	-2.49	-					
Leualp	-2.83	-0.04	-2.19	-0.62	0.23	2.78	-				
Sedbre	-2.06	0.8	-1.44	0.35	1.21	3.78*	0.96	-			
Senpyr	-4.82*	-2.08	-4.02*	-2.99	-2.13	0.2	-2.42	-3.34*	-		
Solvir	-2.99	-0.48	-2.39	-1.03	-0.3	1.85	-0.5	-1.33	1.6	-	
Thypra	-2.64	-0.26	-2.12	-0.75	-0.07	1.94	-0.26	-1.02	1.71	0.19	-
Total number of visits for each plant species											
Silcil	-										
Armcae	-0.32	-									
Bisint	0.16	0.47	-								
Pilvah	-1.88	-1.58	-1.92	-							
Jascri	-2.7	-2.55	-2.7	-1.06	-						
Jurhum	-4.66*	-4.36*	-4.48*	-3.33*	-2.19	-					
Leualp	-1.72	-1.42	-1.81	0.18	1.22	3.50*	-				
Sedbre	-1.41	-1.07	-1.46	0.6	1.64	3.95*	0.41	-			
Senpyr	-3.59*	-3.23*	-3.49*	-2.04	-1	1.12	-2.22	-2.63	-		
Solvir	-1.45	-1.12	-1.51	0.29	1.16	3.14*	0.14	-0.22	2.09	-	
Thypra	-1.71	-1.42	-1.79	-0.12	0.72	2.53	-0.26	-0.59	1.57	-0.35	
Number of visits per flowering plant											
Silcil	-										
Armcae	0.07	-									
Bisint	-3.21	-3.36*	-								
Pilvah	-3.17	-3.45*	0.92	-							
Jascri	-3.01	-3.36*	1.02	0.17	-						
Jurhum	-5.72*	-5.99*	-1.09	-3	-3.14	-					
Leualp	-3.39*	-3.64*	0.7	-0.31	-0.47	2.64	-				
Sedbre	-0.09	-0.17	3.48*	3.83*	3.68*	7.06*	4.03*	-			
Senpyr	-6.90*	-7.07*	-2.2	-4.42*	-4.50*	-1.72	-4.10*	-8.29*	-		
Solvir	-4.72*	-4.84*	-0.98	-2.3	-2.41	-0.06	-2.04	-5.31*	1.25	-	
Thypra	-6.37*	-6.52*	-2.58	-4.26*	-4.35*	-2.18	-4.02*	-7.14*	-0.86	-1.78	

CONCLUSIONS

CONCLUSIONS

- 1) Estimated mean dispersal distances of *S. ciliata* and *A. caespitosa* were short in all cases (less than 1 m). Effective seed dispersal in these two species may be further restricted by post-dispersal factors. The limited dispersal range of the two studied species raises uncertainty on their ability to track their climatic niche through migration under ongoing rapid global warming.
- 2) There were significant differences in the kernel function that provided the best fit, as well as in the mean dispersal distance among populations of *A. caespitosa*, while differences in the mean dispersal distance among *S. ciliata* populations were only marginally significant. These results illustrate the complexity of intraspecific variation in the processes underlying recruitment.
- 3) There was a significant positive association between the spatial demographic and genetic structure of the five populations of *S. ciliata* studied. These findings support theoretical studies showing that stronger spatial aggregation of adults generally increases the magnitude of FSGS compared to random individual distributions.
- 4) Population genetic diversity and inbreeding were homogeneous across all populations of *Silene ciliata* studied. As these populations differed in the intensity of FSGS, no relationship between these two variables and the intensity of FSGS was found.
- 5) The aggregated population spatial pattern of *S. ciliata* seems to be the result of environmental heterogeneity and short-distance dispersal of the species together with the availability of safe sites for establishment, with a reduced effect of plant-plant interactions. These findings downplay the relevance of intraspecific interactions in the spatial patterns of this species and outline the importance of including all relevant processes fostering plant co-occurrence to fully understand plant population dynamics.

- 6) The common garden approach confirmed that among-population variation in flowering onset, number of flowerheads and number of viable fruits in *A. caespitosa* is partially genetically based.
- 7) The stronger selection response in favor of early-flowering individuals in the low edge of *A. caespitosa* elevational range did not correspond with greater proportion of early-flowering individuals. Nevertheless, the observed genetic variability within populations in flowering onset indicates that there is adaptive potential to respond to ongoing global warming.
- 8) As predicted by the Functional Biodiversity hypothesis (FBH), pollinator richness increased at encroached pastures. However, pollinator niche breadth did not vary, contrary to the expectation of the Niche Variation hypothesis. As a result, plant-plant competition for shared pollinators strengthened, which can potentially reduce the reproductive success of some plant species.

COAUTORES

COAUTHORS

AFILIACIÓN DE LOS COAUTORES

Jose María Iriondo (Director de tesis)

Departamento de Biología y Geología, Física y Química Inorgánica. ESCET.
Universidad Rey Juan Carlos. Calle Tulipán S/N, 28933, Móstoles, Madrid, España.

Alfredo García Fernández (Capítulos 1, 2 y 3)

Departamento de Biología y Geología, Física y Química Inorgánica. ESCET.
Universidad Rey Juan Carlos. Calle Tulipán S/N, 28933, Móstoles, Madrid, España.

Juan José Robledo Arnuncio (Capítulos 1 y 2)

Dep. Forest Ecology & Genetics, INIA-CIFOR, Ctra. de la Coruña km 7.5, 28040
Madrid, Spain.

Javier Morente López (Capítulos 2 y 5)

Departamento de Biología y Geología, Física y Química Inorgánica. ESCET.
Universidad Rey Juan Carlos. Calle Tulipán S/N, 28933, Móstoles, Madrid, España.

Ángela López (Capítulo 2)

Departamento de Biología y Geología, Física y Química Inorgánica. ESCET.
Universidad Rey Juan Carlos. Calle Tulipán S/N, 28933, Móstoles, Madrid, España.

Marie Roumet (Capítulo 2)

Plant Ecological Genetics, Institute of Integrative Biology (IBZ), ETH Zurich,
Universitätstrasse 16, 8092 Zurich, Switzerland.

Marcelino de la Cruz (Capítulo 3)

Departamento de Biología y Geología, Física y Química Inorgánica. ESCET.
Universidad Rey Juan Carlos. Calle Tulipán S/N, 28933, Móstoles, Madrid, España.

Gema Escribano (Capítulo 3)

Departamento de Biología y Geología, Física y Química Inorgánica. ESCET.
Universidad Rey Juan Carlos. Calle Tulipán S/N, 28933, Móstoles, Madrid, España.

Adrián Escudero (Capítulo 4)

Departamento de Biología y Geología, Física y Química Inorgánica. ESCET.
Universidad Rey Juan Carlos. Calle Tulipán S/N, 28933, Móstoles, Madrid, España.

Raúl García Camacho (Capítulo 4)

Departamento de Biogeografía y Cambio Global, Museo Nacional de Ciencias
Naturales, MNCN-CSIC, Serrano 115 dpto, E-28006 Madrid, Spain.

Cristina García (Capítulo 5)

Research Center in Biodiversity and Genetic Resources, CIBIO/InBIO, Campus
Agrário de Vairão, Rua Padre Armando Quintas, 4485-661 Vairão, Porto, Portugal.

AGRADECIMIENTOS/ACKNOWLEDGMENTS

Quería comenzar agradeciendo a Txema, mi director de tesis, todo su apoyo durante estos años. Me siento muy afortunado por haber trabajado junto a él y su grupo. Por su dedicación, responsabilidad y respeto a la hora de ejercer su profesión ha sido un ejemplo para mí. Me he sentido muy valorado, respetado y protegido. Gracias por todo Txema, recordaré siempre con cariño estos años.

Quería dedicarle un agradecimiento especial a Gema. Nos conocimos en el año 2002 cuando empezamos la Licenciatura de Ciencias Ambientales. Hace ya 12 años de aquello, 12 años desde que comenzamos a escribir nuestra historia juntos. Me siento muy afortunado de haber podido vivir esta experiencia tan próximo tí. Tengo que agradecerte demasiadas cosas, pero la más importante es tu paciencia, ayuda y cariño en los momentos duros de la tesis; pero también tu alegría en los buenos momentos (la mayoría). No podría haber encontrado una persona mejor para compartir mi vida.

Quería agradecerles a Alfredo García, Luis Gimenez y Raúl García su trabajo previo. Partir del conocimiento adquirido a través de sus tesis ha sido un lujo para mí, y continuar con su trabajo una gran responsabilidad. Espero haber estado a su altura. Creo que Alf se merece unas líneas adicionales. Desde el principio me recibió con los brazos abiertos (y teniendo en cuenta lo grande que es eso es mucho recibir). Me enseñó el departamento, las poblaciones y las peculiaridades de las plantas. No se desalentó cuando le pedía continuar un poco más buscando una población a dos mil metros de altura y pendiente arriba, o cuando quería medir otras 200 plantas antes de irme a casa en pleno agosto. Su apoyo y participación en mi trabajo ha sido magnífico. Aunque he de reconocer que no se quiso quedarse un 24 de agosto haciendo 100 fotos a la parcela de Najarra a las siete de la tarde. Menos mal, porque nunca las he utilizado para nada. ¡Muchas gracias compañero del alma!

Mi otro gran compinche ha sido Javier Morente “el divo”. Apareció como un ángel caído del cielo la primavera de 2011. La paliza que se pegó conmigo registrando contactos “planta-polinizador” (o debo de decir visitante floral...) fue majestuosa. En las 6 semanas que duró el muestreo subimos 17 veces al pico de Peñalara y 12 al Pico del Nevero. Lo que supone un desnivel acumulado de más de 31.000 metros. Eso por no contar las largas horas paseando parcela arriba, parcela abajo buscando polinizadores. En los días malos, después de 8 horas de muestreo contamos 40 contactos. ¡Vaya fiestón! Que se lo digan a Samuel que vino un día el pobre y casi se nos queda en el sitio del frío. Le encontramos parapetado detrás de unas ruinas, con un pañuelo en el bastón para que le encontráramos... y eso que Samuel está más

fuerte que el vinagre! Pero lo mejor no fue eso. Después de todo el día teníamos que pinchar los bichos. El momento más memorable fue cuando después de pinchar una avispa del género *Sphex* volvimos de cenar y la vimos dando vueltas sobre sí misma insertada en el alfiler. Los gritos de espanto todavía resuenan en el Valle del Lozoya. Gracias Javi y mucha suerte con lo tuyo. ¡Te está quedando la tesis preciosa!

También quería agradecerle a Pesca y Carlos toda su ayuda estos años. Nuestros otros compañeros del “High Mountain Team” son excepcionales. Con sus tamaños muéstrales *Adrianicus* y sus cachivaches inventados han marcado estilo por la Sierra de Guadarrama. ¡Hasta las monjas se arremolinaban a su lado para observarlos! ¡Y más de un militar con su rifle CETME en ristre ha tenido que claudicar ante sus huestes! De verdad amigos, muchas gracias por toda vuestra ayuda habéis sido fundamentales para mí estos años. ¡SOS QUIERO!

La verdad que cuando veo la lista de coautores me siento muy afortunado. Su ayuda, sugerencias, comentarios y correcciones han sido indispensables. Les doy las gracias a todos. Juanjo Robledo Arnuncio se involucró muchísimo en los dos primeros capítulos de la tesis. Es una de las mentes más brillantes que he conocido y también una gran persona. He disfrutado mucho y aprendido aún más en cada reunión que he tenido con él. También quería agradecerle a Cristina García que me invitara a visitar su grupo en Portugal. He aprendido mucho escribiendo ese último capítulo con ella. Cristina, gracias por poner un poco de cordura en mi trabajo y por regalarme algunas de tus ideas geniales. Dale muchos recuerdos a los compañeros del CIBIO: Jesus, Laura, Alex, Guille, Catarina, Xavier... ¡Comeros un buen *Ballau* en el Doca a mi salud! ¡Nos veremos pronto!

Alex Widmer me acogió en dos ocasiones en su grupo (Plant Ecological Genetics) en el ETH de Zúrich. Además de enriquecerme como científico me permitió conocer a gente maravillosa. Quería además agradecer a Mónica y Samuel sus cuidados durante mis dos estancias en Zúrich. ¡Os veré pronto amigos! Ahora ya seréis tres personas a las que tendré que ir a visitar a Uster. I thank Alex Widmer for hosting me in his group at EHT-Zurich and giving me the opportunity to know him, Ana, Marie, Fernanda, Margot, Kathrin, Mathias and the rest of the group. Ana, thank you for being so good as host and friend, showing me all those important places in Zurich.

El ambiente de trabajo en el departamento es maravilloso. Me gustaría darle las gracias al departamento en su conjunto por ser tan buenas personas. Os quiero un montón a todos. El mundo sin el equipo cañas (ahora conocido como “vecinos”) no hubiera sido lo mismo (ellos saben quienes son). Esas cervezas en el *piguis* han sido la mejor medicina contra todas las penurias que me ha tocado pasar (muy pocas por

suerte). ¡Gracias Juankar por esas cañas bien tiradas! Mi compañera de despacho Luisa me ha socorrido en todo momento. Siempre al quite con un cacho de pan, una galletita y una sonrisa a tiempo. Gracias también por invitarme a patatas fritas del Coop de Zurich. Aunque los dos sabemos que lo hacías para despegarme del ordenador y que nos fuéramos a pasear por la ciudad. Samuel, Sonia, Silvia, Jesús, Juanlu, Milla, Mauricio, Yoli, Patri, Miguel, Enrique ¡Y los que faltáis!... sois tantos que no puedo dedicaros ni una línea a cada uno. De verdad, muchas gracias a todos. He sido muy feliz en este departamento y ha sido gracias a todos vosotros.

Quería también agradecerle a todos los profesos (ahora compañeros) que me dieron clase en Ciencias Ambientales. Pizca (¡un beso!!), Luisiño, Isabel, David, Rosa, Natalia, Marcos, Emilio, M^a Carmen.... ¡Muchas gracias por todo! Disfruté mucho las clases y aprendí mucho. Estoy aquí gracias a vosotros.

Adri, muchas gracias por tu cariño y ayuda durante estos años. Sin ser mi director de tesis siempre has estado dispuesto a echarme una mano y a darme tus maravillosos consejos. Muchas gracias por todo.

Casi las últimas palabras las he reservado para Emilio. Junto a él di mis primeros pasos en ciencia. Con él diseñé mi primer muestreo de campo, escribí mi primer artículo y recibí el primer rechazo de una revista. Muchas gracias por ser mi maestro durante los primeros años. Gracias también por los madrugones los fines de semana para muestrear tejones. Pero sobre todo gracias por ser mi amigo y tan buena persona.

He reservado las últimas líneas para lo más importante que tengo. Mi familia. He llegado hasta aquí gracias al amor, dedicación y apoyo incondicional de mis padres, Felipe y Vicen. A ellos les dedicó la tesis. Siempre me han hecho sentir amado y protegido. Han dado todo lo que tenían por mi vida. Estoy muy orgulloso de ellos. Mis hermanos han sido un ejemplo para mí. A menudo me dicen que me parezco a ellos, y para mí eso es el mejor cumplido que me podrían hacer. Quería daros las gracias a vosotros y a vuestra (mí) familia; Tatiana, Cristina, Helena, Victor, Marta, Concha por todo el cariño que me habéis dado y por haber sabido entender mis ausencias durante estos cuatro años. ¡Os quiero!

¡Gracias a todos por ayudarme a escribir mi historia!

Carlos

Madrid 18 de Septiembre de 2014

